

Table 2.11C Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	891	874	863	920	825	852
12	23	20	18	12	9	9
13	32	33	25	26	17	19
14	41	35	35	30	24	22
15	48	47	41	38	31	29
16	59	56	48	50	37	35
17	65	63	56	54	45	44
18	96	97	87	86	73	69
19	101	95	92	84	72	71
20	99	105	85	88	73	79
21	111	102	92	85	81	71
22	108	93	89	78	77	69
23	96	91	80	73	71	68
24	97	95	76	73	67	65
25	84	93	69	78	62	72
26-29	384	370	296	286	273	263
30-34	415	420	305	298	279	270
35-39	405	397	267	283	242	259
40-44	456	442	280	308	256	275
45-49	441	475	278	298	262	279
50-54	620	627	402	437	382	409
55-59	580	625	323	378	308	355
60-64	480	541	266	288	257	272
65 or Older	450	466	312	317	283	304

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.11D Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.35	0.34	0.34	0.36	0.33	0.34
12	0.58	0.51	0.48	0.32	0.24	0.24
13	0.78	0.74	0.62	0.60	0.43	0.44
14	0.85	0.83	0.75	0.72	0.55	0.55
15	0.92	0.93	0.81	0.79	0.65	0.68
16	1.08	1.04	0.94	0.97	0.80	0.76
17	1.08	1.08	1.02	1.03	0.87	0.89
18	1.14	1.18	1.12	1.18	1.09	1.08
19	1.12	1.23	1.20	1.25	1.20	1.20
20	1.14	1.23	1.30	1.28	1.32	1.31
21	1.18	1.37	1.35	1.33	1.34	1.28
22	1.14	1.13	1.21	1.18	1.25	1.17
23	1.12	1.15	1.26	1.27	1.26	1.25
24	1.11	1.20	1.28	1.34	1.24	1.31
25	1.23	1.21	1.38	1.39	1.31	1.41
26-29	1.12	1.06	1.30	1.25	1.25	1.23
30-34	0.93	1.00	1.13	1.15	1.11	1.10
35-39	1.08	1.10	1.08	1.13	1.03	1.11
40-44	0.99	1.00	1.11	1.14	1.05	1.07
45-49	0.94	0.93	1.00	1.05	0.98	1.02
50-54	1.42	1.37	1.48	1.56	1.41	1.51
55-59	1.49	1.43	1.46	1.60	1.40	1.55
60-64	1.55	1.70	1.53	1.63	1.50	1.55
65 or Older	1.18	1.20	0.82	0.81	0.75	0.78

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.11P Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.3371		0.5467		0.5678	
12	0.2162		0.0631		0.8257	
13	0.4101		0.8357		0.9865	
14	0.0196		0.2514		0.4282	
15	0.8179		0.4242		0.9664	
16	0.0416		0.4548		0.0885	
17	0.2011		0.2063		0.0787	
18	0.8277		0.7473		0.5776	
19	0.7067		0.8157		0.6329	
20	0.7917		0.6667		0.9319	
21	0.1120		0.2619		0.4417	
22	0.0136		0.0646		0.1477	
23	0.3365		0.2417		0.2874	
24	0.7008		0.6464		0.8317	
25	0.6453		1.0000		0.4941	
26-29	0.6029		0.7207		0.6712	
30-34	0.7127		0.1936		0.1034	
35-39	0.0837		0.5823		0.6732	
40-44	0.4710		0.6755		0.7850	
45-49	0.5193		0.4080		0.3342	
50-54	0.4184		0.9863		0.4269	
55-59	0.6504		0.2486		0.3830	
60-64	0.6421		0.5473		0.6301	
65 or Older	0.9443		0.9781		0.4659	

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.12C Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	909	881	836	839	789	781
12	20	17	17	11	5	7
13	25	29	20	23	13	17
14	37	31	31	26	21	20
15	44	42	36	33	27	25
16	52	51	43	43	33	32
17	60	60	48	49	40	39
18	88	89	77	77	65	63
19	91	88	81	75	66	63
20	92	95	79	77	68	68
21	106	93	85	75	74	66
22	103	88	78	71	68	64
23	91	88	74	66	64	59
24	92	90	69	70	63	62
25	80	88	64	69	57	64
26-29	368	355	269	262	247	243
30-34	400	401	276	269	250	252
35-39	378	377	239	255	222	226
40-44	449	419	242	274	228	260
45-49	429	456	261	261	245	249
50-54	606	618	356	410	346	392
55-59	567	611	304	341	290	322
60-64	477	523	235	257	225	243
65 or Older	474	469	282	277	259	266

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.12D Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.36	0.35	0.33	0.33	0.31	0.31
12	0.53	0.43	0.43	0.27	0.13	0.19
13	0.63	0.66	0.51	0.55	0.33	0.41
14	0.79	0.75	0.68	0.64	0.49	0.50
15	0.86	0.89	0.74	0.72	0.58	0.60
16	1.02	1.01	0.88	0.89	0.72	0.70
17	1.06	1.09	0.92	0.98	0.82	0.81
18	1.15	1.24	1.09	1.17	1.03	1.04
19	1.21	1.29	1.27	1.25	1.18	1.16
20	1.25	1.21	1.32	1.23	1.28	1.23
21	1.27	1.39	1.36	1.33	1.31	1.24
22	1.20	1.14	1.24	1.21	1.25	1.19
23	1.17	1.27	1.27	1.28	1.22	1.24
24	1.15	1.28	1.26	1.35	1.20	1.30
25	1.27	1.28	1.35	1.40	1.26	1.35
26-29	1.21	1.05	1.26	1.20	1.19	1.16
30-34	1.01	1.02	1.10	1.10	1.06	1.05
35-39	1.11	1.08	1.03	1.10	0.99	1.04
40-44	1.05	1.07	1.00	1.09	0.96	1.06
45-49	1.04	0.99	0.99	0.98	0.95	0.96
50-54	1.48	1.42	1.37	1.54	1.35	1.48
55-59	1.57	1.53	1.40	1.50	1.36	1.45
60-64	1.61	1.76	1.40	1.49	1.36	1.41
65 or Older	1.25	1.20	0.74	0.71	0.68	0.68

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.12P Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.3822		0.2928		0.4373	
12	0.1490		0.0622		0.5373	
13	0.8448		0.4500		0.2598	
14	0.0545		0.1355		0.6405	
15	0.8042		0.7516		0.9733	
16	0.0080		0.3512		0.1126	
17	0.1509		0.6311		0.2488	
18	0.4775		0.2535		0.3832	
19	0.5779		0.5123		0.9119	
20	0.4712		0.1934		0.1900	
21	0.0529		0.0692		0.3852	
22	0.0090		0.0906		0.2306	
23	0.1978		0.0954		0.0833	
24	0.9340		0.7536		0.7188	
25	0.5275		0.1946		0.1987	
26-29	0.3378		0.8066		0.8110	
30-34	0.8810		0.3644		0.4858	
35-39	0.4569		0.6020		0.4559	
40-44	0.5281		0.2092		0.2884	
45-49	0.3884		0.5984		0.8150	
50-54	0.5187		0.7400		0.7938	
55-59	0.7368		0.5331		0.5514	
60-64	0.3237		0.6088		0.6525	
65 or Older	0.8721		0.8762		0.7782	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.13C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	680	699	331	350	289	313
12	8	9	6	6	3	4
13	17	17	14	14	8	10
14	22	18	16	16	12	9
15	22	24	19	20	13	13
16	31	29	24	23	15	16
17	36	31	30	25	18	17
18	47	48	38	42	26	34
19	57	51	42	37	29	27
20	48	55	36	37	27	33
21	54	52	34	32	25	22
22	57	49	38	33	30	26
23	44	47	29	28	22	23
24	44	44	26	27	20	21
25	46	54	26	32	24	26
26-29	193	185	118	104	90	93
30-34	213	196	115	105	105	88
35-39	239	238	105	131	93	126
40-44	241	227	114	90	106	84
45-49	206	233	75	105	66	96
50-54	289	293	115	94	113	78
55-59	228	243	74	95	65	92
60-64	200	199	88	87	83	69
65 or Older	280	316	99	129	87	120

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.13D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.27	0.28	0.13	0.14	0.11	0.12
12	0.23	0.24	0.16	0.16	0.09	0.11
13	0.43	0.40	0.36	0.33	0.22	0.24
14	0.51	0.45	0.37	0.40	0.27	0.24
15	0.50	0.55	0.43	0.47	0.30	0.30
16	0.67	0.66	0.54	0.53	0.36	0.37
17	0.76	0.70	0.65	0.58	0.40	0.40
18	0.86	0.91	0.72	0.82	0.52	0.69
19	1.06	1.01	0.81	0.76	0.58	0.58
20	1.01	1.03	0.81	0.76	0.62	0.68
21	1.05	1.04	0.73	0.68	0.58	0.48
22	1.13	1.01	0.86	0.75	0.69	0.59
23	0.98	1.02	0.66	0.65	0.52	0.56
24	0.96	1.03	0.63	0.65	0.49	0.51
25	1.14	1.16	0.70	0.75	0.63	0.62
26-29	1.01	0.96	0.67	0.61	0.52	0.55
30-34	0.92	0.86	0.56	0.51	0.51	0.44
35-39	1.00	0.99	0.52	0.64	0.46	0.61
40-44	0.99	0.96	0.54	0.44	0.50	0.41
45-49	0.84	0.90	0.33	0.46	0.29	0.42
50-54	1.19	1.19	0.51	0.43	0.50	0.35
55-59	1.14	1.18	0.40	0.49	0.35	0.47
60-64	1.18	1.16	0.56	0.52	0.53	0.42
65 or Older	0.74	0.81	0.26	0.33	0.23	0.31

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.13P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.8252		0.6963		0.5722	
12	0.8703		0.8988		0.9025	
13	0.7231		0.7456		0.6991	
14	0.2768		0.5652		0.2898	
15	0.6088		0.5671		0.5076	
16	0.5600		0.3814		0.7900	
17	0.5812		0.2892		0.7358	
18	0.6632		0.1691		0.1226	
19	0.9164		0.5329		0.6515	
20	0.9110		0.2779		0.9307	
21	0.8999		0.9934		0.3949	
22	0.0940		0.9557		0.9956	
23	0.5219		0.8406		0.7404	
24	0.5671		0.1025		0.1691	
25	0.6967		0.3185		0.8029	
26-29	0.3587		0.7644		0.1734	
30-34	0.2403		0.2375		0.0835	
35-39	0.5965		0.2349		0.1097	
40-44	0.8941		0.0119		0.0090	
45-49	0.7270		0.0669		0.1002	
50-54	0.7368		0.4198		0.1612	
55-59	0.6220		0.1348		0.0792	
60-64	0.5773		0.8051		0.9021	
65 or Older	0.4321		0.4271		0.2852	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.14C Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	884	929	499	508	358	362
12	11	8	9	5	6	4
13	21	17	16	13	10	9
14	25	23	21	19	13	10
15	35	29	30	24	18	16
16	39	38	31	31	22	19
17	43	41	37	34	25	23
18	72	72	64	64	43	42
19	76	72	65	59	38	39
20	76	75	59	56	42	42
21	80	82	61	55	38	36
22	81	73	56	52	38	35
23	75	69	47	46	32	34
24	73	70	47	44	30	33
25	66	73	41	45	29	30
26-29	296	289	170	168	101	106
30-34	300	302	167	165	126	110
35-39	300	291	143	151	106	103
40-44	304	333	141	171	102	108
45-49	297	318	136	152	104	114
50-54	438	440	203	190	155	141
55-59	395	393	176	184	125	125
60-64	316	341	131	130	108	102
65 or Older	433	429	131	135	88	109

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.14D Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.35	0.37	0.20	0.20	0.14	0.14
12	0.30	0.22	0.23	0.14	0.16	0.11
13	0.53	0.40	0.40	0.30	0.26	0.23
14	0.56	0.56	0.48	0.48	0.30	0.24
15	0.71	0.66	0.64	0.54	0.41	0.39
16	0.83	0.78	0.68	0.67	0.51	0.43
17	0.88	0.85	0.77	0.75	0.54	0.53
18	1.10	1.16	1.04	1.10	0.79	0.81
19	1.13	1.20	1.05	1.10	0.74	0.80
20	1.25	1.26	1.14	1.02	0.93	0.82
21	1.27	1.40	1.15	1.09	0.81	0.80
22	1.25	1.24	1.10	1.08	0.85	0.80
23	1.28	1.26	0.98	0.97	0.74	0.80
24	1.30	1.27	1.02	0.98	0.69	0.77
25	1.40	1.32	1.02	0.99	0.76	0.74
26-29	1.27	1.20	0.94	0.93	0.58	0.61
30-34	1.08	1.08	0.78	0.77	0.62	0.53
35-39	1.15	1.11	0.69	0.69	0.52	0.51
40-44	1.12	1.11	0.66	0.76	0.48	0.51
45-49	1.06	1.08	0.58	0.65	0.45	0.50
50-54	1.51	1.57	0.87	0.82	0.67	0.62
55-59	1.62	1.63	0.91	0.92	0.66	0.63
60-64	1.68	1.71	0.82	0.77	0.68	0.61
65 or Older	1.14	1.10	0.34	0.35	0.23	0.28

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.14P Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.0553		0.3057		0.7154	
12	0.0111		0.0405		0.8952	
13	0.0128		0.1346		0.3171	
14	0.0591		0.3473		0.2510	
15	0.2112		0.0240		0.4825	
16	0.3034		0.4531		0.0064	
17	0.2874		0.0503		0.1666	
18	0.9561		0.6694		0.8329	
19	0.9493		0.1832		0.5468	
20	0.1063		0.1105		0.1923	
21	0.5690		0.2704		0.4145	
22	0.0657		0.8880		0.6175	
23	0.6893		0.4336		0.3882	
24	0.5434		0.4370		0.5057	
25	0.8581		0.2093		0.5075	
26-29	0.7868		0.8288		0.5460	
30-34	0.4290		0.1472		0.0596	
35-39	0.4979		0.4821		0.6936	
40-44	0.0121		0.1864		0.4408	
45-49	0.2521		0.4139		0.1204	
50-54	0.1708		0.2994		0.4040	
55-59	0.2582		0.7197		0.9886	
60-64	0.6292		0.3614		0.5140	
65 or Older	0.4183		0.6193		0.2374	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.15C Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	670	713	907	966	957	990
12	28	20	23	17	14	11
13	36	39	30	33	17	20
14	50	44	41	37	25	24
15	65	60	59	54	36	36
16	73	67	63	61	43	42
17	74	74	67	69	51	51
18	112	114	104	107	89	84
19	118	107	112	101	92	86
20	113	117	107	111	89	96
21	132	130	130	128	120	112
22	117	111	115	108	107	94
23	106	108	102	106	95	95
24	112	104	107	99	96	92
25	94	111	89	108	82	98
26-29	426	415	408	387	361	355
30-34	464	453	429	417	383	387
35-39	463	472	427	444	370	408
40-44	537	492	479	455	427	398
45-49	495	504	447	463	403	393
50-54	703	673	615	596	534	536
55-59	646	680	568	583	499	513
60-64	536	633	459	517	418	451
65 or Older	410	443	535	501	508	487

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.15D Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.27	0.28	0.36	0.38	0.38	0.39
12	0.68	0.51	0.58	0.43	0.37	0.27
13	0.85	0.87	0.72	0.76	0.43	0.48
14	0.97	0.95	0.85	0.85	0.57	0.59
15	1.04	1.04	1.00	1.01	0.76	0.77
16	1.11	1.05	1.06	1.06	0.89	0.90
17	1.06	1.08	1.04	1.11	0.93	1.01
18	1.07	1.07	1.11	1.19	1.24	1.19
19	1.00	0.98	1.13	1.07	1.26	1.24
20	0.94	0.98	1.13	1.05	1.29	1.26
21	0.84	0.82	0.92	0.97	1.25	1.29
22	0.67	0.80	0.87	1.00	1.23	1.23
23	0.73	0.64	1.02	0.87	1.14	1.21
24	0.63	0.77	0.81	0.99	1.19	1.24
25	1.05	0.69	1.22	0.94	1.40	1.19
26-29	0.73	0.69	0.97	1.01	1.15	1.20
30-34	0.66	0.67	0.95	0.95	1.12	1.07
35-39	0.82	0.83	1.07	1.05	1.16	1.24
40-44	0.64	0.72	0.96	1.01	1.16	1.11
45-49	0.63	0.60	0.94	1.00	1.10	1.09
50-54	1.05	0.94	1.53	1.46	1.59	1.60
55-59	1.02	1.08	1.58	1.72	1.64	1.81
60-64	1.10	1.26	1.76	1.84	1.89	1.88
65 or Older	1.08	1.14	1.41	1.29	1.34	1.25

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.15P Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.4415		0.4558		0.8716	
12	0.0089		0.1601		0.0732	
13	0.4114		0.7357		0.8318	
14	0.0421		0.1230		0.8449	
15	0.2602		0.5419		0.1646	
16	0.0090		0.2877		0.2942	
17	0.0614		0.1054		0.1421	
18	0.7665		0.6593		0.6173	
19	0.1869		0.2830		0.2994	
20	0.4504		0.6574		0.6138	
21	0.5570		0.9544		0.9699	
22	0.0216		0.0147		0.1139	
23	0.0748		0.0097		0.4498	
24	0.4300		0.1800		0.4437	
25	0.0344		0.0536		0.2013	
26-29	0.3634		0.6737		0.5340	
30-34	0.6238		0.6427		0.3297	
35-39	0.4537		0.6288		0.5318	
40-44	0.3558		0.4064		0.9831	
45-49	0.1739		0.6505		0.9974	
50-54	0.9074		0.4044		0.2314	
55-59	0.9414		0.1619		0.2328	
60-64	0.5490		0.9912		0.6437	
65 or Older	0.8555		0.7837		0.6191	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.16C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2009 and 2010

Age Category	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	957	990	785	754	406	421
12	14	11	10	5	2	1
13	17	20	13	11	4	3
14	25	24	18	17	5	7
15	36	36	27	27	13	13
16	43	42	31	32	14	14
17	51	51	42	41	22	19
18	89	84	74	68	34	32
19	92	86	81	71	47	41
20	89	96	74	77	46	50
21	120	112	92	88	45	52
22	107	94	86	73	50	45
23	95	95	72	71	40	39
24	96	92	75	70	38	38
25	82	98	63	69	31	41
26-29	361	355	249	260	120	135
30-34	383	387	264	283	139	139
35-39	370	408	238	258	118	128
40-44	427	398	258	245	121	135
45-49	403	393	247	250	123	136
50-54	534	536	310	306	176	167
55-59	499	513	282	277	150	152
60-64	418	451	189	225	101	98
65 or Older	508	487	320	263	130	122

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.16D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Standard Errors of Percentages, 2009 and 2010

Age Category	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.38	0.39	0.31	0.30	0.16	0.17
12	0.37	0.27	0.27	0.13	0.06	0.03
13	0.43	0.48	0.33	0.26	0.11	0.07
14	0.57	0.59	0.42	0.42	0.12	0.17
15	0.76	0.77	0.59	0.62	0.30	0.30
16	0.89	0.90	0.70	0.73	0.32	0.33
17	0.93	1.01	0.80	0.85	0.49	0.44
18	1.24	1.19	1.13	1.09	0.63	0.62
19	1.26	1.24	1.30	1.17	0.93	0.83
20	1.29	1.26	1.29	1.24	1.00	0.97
21	1.25	1.29	1.28	1.41	0.90	1.06
22	1.23	1.23	1.27	1.23	1.05	0.96
23	1.14	1.21	1.28	1.30	0.91	0.91
24	1.19	1.24	1.32	1.32	0.84	0.89
25	1.40	1.19	1.39	1.29	0.83	0.97
26-29	1.15	1.20	1.10	1.20	0.68	0.75
30-34	1.12	1.07	1.10	1.08	0.68	0.66
35-39	1.16	1.24	1.04	1.07	0.58	0.62
40-44	1.16	1.11	1.00	0.96	0.57	0.62
45-49	1.10	1.09	0.94	0.97	0.53	0.59
50-54	1.59	1.60	1.21	1.25	0.76	0.74
55-59	1.64	1.81	1.31	1.25	0.79	0.76
60-64	1.89	1.88	1.15	1.29	0.64	0.59
65 or Older	1.34	1.25	0.84	0.68	0.34	0.31

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.16P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Age Category	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.8716		0.1811		0.5738	
12	0.0732		0.0373		0.6287	
13	0.8318		0.1702		0.3261	
14	0.8449		0.7606		0.3837	
15	0.1646		0.6670		0.2264	
16	0.2942		0.4745		0.8232	
17	0.1421		0.0308		0.0127	
18	0.6173		0.1942		0.8629	
19	0.2994		0.4295		0.7070	
20	0.6138		0.3542		0.9868	
21	0.9699		0.8743		0.4136	
22	0.1139		0.0033		0.2642	
23	0.4498		0.8608		0.4528	
24	0.4437		0.8395		0.4122	
25	0.2013		0.3902		0.2821	
26-29	0.5340		0.9259		0.5175	
30-34	0.3297		0.7805		0.8058	
35-39	0.5318		0.4061		0.6152	
40-44	0.9831		0.7142		0.4221	
45-49	0.9974		0.5949		0.5226	
50-54	0.2314		0.8837		0.7154	
55-59	0.2328		0.2227		0.5113	
60-64	0.6437		0.4530		0.5345	
65 or Older	0.6191		0.0394		0.1877	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.