

Table 2.1C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	891	874	863	920	825	852
Cigarettes	909	881	836	839	789	781
Smokeless Tobacco	680	699	331	350	289	313
Cigars	884	929	499	508	358	362
Pipe Tobacco ²	703	693	--	--	151	147
ALCOHOL	670	713	907	966	957	990
Binge Alcohol Use ³	--	--	--	--	785	754
Heavy Alcohol Use ³	--	--	--	--	406	421

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.1D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Percentages, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.35	0.34	0.34	0.36	0.33	0.34
Cigarettes	0.36	0.35	0.33	0.33	0.31	0.31
Smokeless Tobacco	0.27	0.28	0.13	0.14	0.11	0.12
Cigars	0.35	0.37	0.20	0.20	0.14	0.14
Pipe Tobacco ²	0.28	0.27	--	--	0.06	0.06
ALCOHOL	0.27	0.28	0.36	0.38	0.38	0.39
Binge Alcohol Use ³	--	--	--	--	0.31	0.30
Heavy Alcohol Use ³	--	--	--	--	0.16	0.17

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.1P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.3371		0.5467		0.5678	
Cigarettes	0.3822		0.2928		0.4373	
Smokeless Tobacco	0.8252		0.6963		0.5722	
Cigars	0.0553		0.3057		0.7154	
Pipe Tobacco ²	0.2523		--		0.8161	
ALCOHOL	0.4415		0.4558		0.8716	
Binge Alcohol Use ³	--		--		0.1811	
Heavy Alcohol Use ³	--		--		0.5738	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.2C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Numbers in Thousands, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	40	39	32	29	19	21
Cigarettes	33	33	27	26	14	18
Smokeless Tobacco	19	19	15	15	9	11
Cigars	24	19	18	14	12	10
Pipe Tobacco ²	11	9	--	--	6	6
ALCOHOL	47	44	37	37	22	22
Binge Alcohol Use ³	--	--	--	--	17	12
Heavy Alcohol Use ³	--	--	--	--	5	3

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.2D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Percentages, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.49	0.46	0.40	0.35	0.25	0.26
Cigarettes	0.41	0.40	0.34	0.31	0.18	0.23
Smokeless Tobacco	0.26	0.24	0.20	0.19	0.12	0.14
Cigars	0.30	0.23	0.23	0.17	0.15	0.13
Pipe Tobacco ²	0.15	0.12	--	--	0.07	0.07
ALCOHOL	0.54	0.52	0.46	0.45	0.28	0.28
Binge Alcohol Use ³	--	--	--	--	0.22	0.15
Heavy Alcohol Use ³	--	--	--	--	0.06	0.04

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.2P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.1802		0.2655		0.8860	
Cigarettes	0.5118		0.7370		0.1923	
Smokeless Tobacco	0.7241		0.8005		0.6808	
Cigars	0.0007		0.0239		0.3525	
Pipe Tobacco ²	0.1987		--		0.8025	
ALCOHOL	0.0448		0.3251		0.4171	
Binge Alcohol Use ³	--		--		0.0264	
Heavy Alcohol Use ³	--		--		0.2793	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.3C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Numbers in Thousands, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	64	58	55	49	40	38
Cigarettes	57	51	48	42	34	33
Smokeless Tobacco	32	31	25	26	18	16
Cigars	44	38	38	32	23	19
Pipe Tobacco ²	19	15	--	--	13	8
ALCOHOL	86	75	74	67	46	43
Binge Alcohol Use ³	--	--	--	--	32	32
Heavy Alcohol Use ³	--	--	--	--	14	14

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.3D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Percentages, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.64	0.62	0.57	0.55	0.44	0.45
Cigarettes	0.59	0.57	0.52	0.48	0.38	0.40
Smokeless Tobacco	0.36	0.37	0.29	0.31	0.20	0.20
Cigars	0.46	0.44	0.42	0.38	0.26	0.23
Pipe Tobacco ²	0.22	0.18	--	--	0.15	0.10
ALCOHOL	0.74	0.70	0.68	0.66	0.50	0.49
Binge Alcohol Use ³	--	--	--	--	0.36	0.37
Heavy Alcohol Use ³	--	--	--	--	0.17	0.17

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.3P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.0934		0.1856		0.6204	
Cigarettes	0.1589		0.2321		0.8068	
Smokeless Tobacco	0.7613		0.4199		0.8830	
Cigars	0.0340		0.0228		0.2334	
Pipe Tobacco ²	0.1304		--		0.0751	
ALCOHOL	0.0323		0.1529		0.3099	
Binge Alcohol Use ³	--		--		0.5954	
Heavy Alcohol Use ³	--		--		0.4663	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.4C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Numbers in Thousands, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	87	88	74	77	59	58
Cigarettes	80	81	64	70	53	52
Smokeless Tobacco	48	43	38	34	24	23
Cigars	58	55	48	46	32	30
Pipe Tobacco ²	28	24	--	--	16	14
ALCOHOL	105	105	93	96	67	68
Binge Alcohol Use ³	--	--	--	--	53	54
Heavy Alcohol Use ³	--	--	--	--	26	24

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.4D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Percentages, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.75	0.78	0.70	0.74	0.60	0.60
Cigarettes	0.74	0.77	0.64	0.71	0.56	0.54
Smokeless Tobacco	0.51	0.48	0.42	0.39	0.27	0.27
Cigars	0.62	0.56	0.52	0.49	0.36	0.34
Pipe Tobacco ²	0.31	0.28	--	--	0.18	0.16
ALCOHOL	0.75	0.76	0.74	0.76	0.64	0.69
Binge Alcohol Use ³	--	--	--	--	0.55	0.57
Heavy Alcohol Use ³	--	--	--	--	0.29	0.28

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.4P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.0202		0.1576		0.0152	
Cigarettes	0.0042		0.3273		0.0539	
Smokeless Tobacco	0.4077		0.1606		0.9280	
Cigars	0.1285		0.0461		0.0039	
Pipe Tobacco ²	0.3040		--		0.0354	
ALCOHOL	0.0010		0.0485		0.0697	
Binge Alcohol Use ³	--		--		0.0333	
Heavy Alcohol Use ³	--		--		0.0275	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.5C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Numbers in Thousands, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	193	198	171	167	137	139
Cigarettes	175	180	152	145	124	122
Smokeless Tobacco	98	96	74	76	51	61
Cigars	141	141	117	112	74	72
Pipe Tobacco ²	51	62	--	--	24	31
ALCOHOL	233	232	219	218	187	178
Binge Alcohol Use ³	--	--	--	--	156	141
Heavy Alcohol Use ³	--	--	--	--	86	79

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.5D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Percentages, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.67	0.74	0.73	0.74	0.72	0.72
Cigarettes	0.71	0.77	0.74	0.72	0.71	0.70
Smokeless Tobacco	0.59	0.59	0.48	0.50	0.34	0.42
Cigars	0.66	0.72	0.61	0.63	0.46	0.46
Pipe Tobacco ²	0.36	0.42	--	--	0.18	0.22
ALCOHOL	0.63	0.60	0.68	0.65	0.82	0.74
Binge Alcohol Use ³	--	--	--	--	0.81	0.70
Heavy Alcohol Use ³	--	--	--	--	0.56	0.51

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.5P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.9635		0.6366		0.8880	
Cigarettes	0.3983		0.0985		0.2508	
Smokeless Tobacco	0.6570		0.5886		0.2494	
Cigars	0.4188		0.1512		0.7536	
Pipe Tobacco ²	0.0090		--		0.0253	
ALCOHOL	0.3285		0.4144		0.4119	
Binge Alcohol Use ³	--		--		0.1462	
Heavy Alcohol Use ³	--		--		0.9788	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.6C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Numbers in Thousands, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	278	262	215	209	193	178
Cigarettes	261	243	197	180	168	157
Smokeless Tobacco	128	121	77	74	60	56
Cigars	209	192	133	121	84	78
Pipe Tobacco ²	85	83	--	--	32	28
ALCOHOL	321	331	312	321	283	279
Binge Alcohol Use ³	--	--	--	--	213	197
Heavy Alcohol Use ³	--	--	--	--	107	110

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.6D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Percentages, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.54	0.59	0.60	0.63	0.62	0.61
Cigarettes	0.57	0.63	0.63	0.63	0.59	0.59
Smokeless Tobacco	0.50	0.49	0.34	0.33	0.28	0.26
Cigars	0.62	0.59	0.51	0.49	0.37	0.36
Pipe Tobacco ²	0.37	0.38	--	--	0.16	0.14
ALCOHOL	0.39	0.36	0.48	0.48	0.60	0.61
Binge Alcohol Use ³	--	--	--	--	0.66	0.64
Heavy Alcohol Use ³	--	--	--	--	0.45	0.47

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.6P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.0230		0.0417		0.0720	
Cigarettes	0.0064		0.0034		0.0179	
Smokeless Tobacco	0.6348		0.3458		0.8147	
Cigars	0.1203		0.4415		0.6510	
Pipe Tobacco ²	0.2200		--		0.1186	
ALCOHOL	0.5378		0.7168		0.8816	
Binge Alcohol Use ³	--		--		0.2903	
Heavy Alcohol Use ³	--		--		0.6916	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.7C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Numbers in Thousands, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	268	270	314	317	307	306
Cigarettes	288	273	301	305	287	291
Smokeless Tobacco	247	237	161	152	137	132
Cigars	298	297	220	221	158	149
Pipe Tobacco ²	171	171	--	--	71	52
ALCOHOL	180	180	247	263	294	306
Binge Alcohol Use ³	--	--	--	--	285	301
Heavy Alcohol Use ³	--	--	--	--	175	186

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.7D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Percentages, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.74	0.74	0.87	0.87	0.85	0.84
Cigarettes	0.79	0.75	0.83	0.84	0.79	0.80
Smokeless Tobacco	0.68	0.65	0.44	0.42	0.38	0.36
Cigars	0.82	0.81	0.61	0.61	0.43	0.41
Pipe Tobacco ²	0.47	0.47	--	--	0.20	0.14
ALCOHOL	0.50	0.49	0.68	0.72	0.81	0.84
Binge Alcohol Use ³	--	--	--	--	0.79	0.82
Heavy Alcohol Use ³	--	--	--	--	0.48	0.51

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.7P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.9163		0.4920		0.3731	
Cigarettes	0.5760		0.3800		0.7004	
Smokeless Tobacco	0.8788		0.5713		0.8359	
Cigars	0.7059		0.2346		0.3174	
Pipe Tobacco ²	0.8949		--		0.3938	
ALCOHOL	0.8073		0.5242		0.8107	
Binge Alcohol Use ³	--		--		0.9138	
Heavy Alcohol Use ³	--		--		0.8097	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.8C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Numbers in Thousands, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	772	770	700	793	684	747
Cigarettes	793	782	676	729	655	688
Smokeless Tobacco	607	619	249	275	233	253
Cigars	795	823	377	427	293	307
Pipe Tobacco ²	656	659	--	--	122	131
ALCOHOL	598	648	846	866	869	875
Binge Alcohol Use ³	--	--	--	--	678	624
Heavy Alcohol Use ³	--	--	--	--	332	347

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.8D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Percentages, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.49	0.49	0.44	0.50	0.43	0.47
Cigarettes	0.50	0.49	0.43	0.46	0.42	0.43
Smokeless Tobacco	0.39	0.39	0.16	0.17	0.15	0.16
Cigars	0.51	0.52	0.24	0.27	0.19	0.19
Pipe Tobacco ²	0.42	0.42	--	--	0.08	0.08
ALCOHOL	0.38	0.41	0.54	0.55	0.55	0.55
Binge Alcohol Use ³	--	--	--	--	0.43	0.39
Heavy Alcohol Use ³	--	--	--	--	0.21	0.22

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.8P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.5894		0.8594		0.8891	
Cigarettes	0.6653		0.9935		0.9221	
Smokeless Tobacco	0.8988		0.5512		0.6123	
Cigars	0.1406		0.6813		0.5265	
Pipe Tobacco ²	0.1233		--		0.3328	
ALCOHOL	0.7423		0.6958		0.9159	
Binge Alcohol Use ³	--		--		0.2970	
Heavy Alcohol Use ³	--		--		0.5602	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.9C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Numbers in Thousands, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	523	506	627	667	598	627
Cigarettes	586	540	598	579	561	538
Smokeless Tobacco	594	611	316	335	282	301
Cigars	646	672	448	446	333	321
Pipe Tobacco ²	628	605	--	--	143	137
ALCOHOL	393	394	571	627	650	676
Binge Alcohol Use ³	--	--	--	--	605	578
Heavy Alcohol Use ³	--	--	--	--	349	353

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.9D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Percentages, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.43	0.41	0.51	0.54	0.49	0.51
Cigarettes	0.48	0.44	0.49	0.47	0.46	0.44
Smokeless Tobacco	0.49	0.49	0.26	0.27	0.23	0.24
Cigars	0.53	0.54	0.37	0.36	0.27	0.26
Pipe Tobacco ²	0.51	0.49	--	--	0.12	0.11
ALCOHOL	0.32	0.32	0.47	0.51	0.53	0.55
Binge Alcohol Use ³	--	--	--	--	0.49	0.47
Heavy Alcohol Use ³	--	--	--	--	0.29	0.29

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.9P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.7080		0.7523		0.7282	
Cigarettes	0.9014		0.8717		0.8614	
Smokeless Tobacco	0.8109		0.8021		0.7651	
Cigars	0.0878		0.2901		0.5629	
Pipe Tobacco ²	0.4810		--		0.7857	
ALCOHOL	0.4949		0.0728		0.7611	
Binge Alcohol Use ³	--		--		0.3018	
Heavy Alcohol Use ³	--		--		0.6326	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.10C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Numbers in Thousands, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	652	649	520	572	501	535
Cigarettes	653	657	512	559	493	526
Smokeless Tobacco	249	245	73	89	51	78
Cigars	477	486	203	206	129	153
Pipe Tobacco ²	232	225	--	--	37	54
ALCOHOL	521	538	664	654	683	634
Binge Alcohol Use ³	--	--	--	--	441	441
Heavy Alcohol Use ³	--	--	--	--	192	201

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.10D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Percentages, 2009 and 2010

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.50	0.50	0.40	0.44	0.39	0.41
Cigarettes	0.50	0.50	0.39	0.43	0.38	0.40
Smokeless Tobacco	0.19	0.19	0.06	0.07	0.04	0.06
Cigars	0.37	0.37	0.16	0.16	0.10	0.12
Pipe Tobacco ²	0.18	0.17	--	--	0.03	0.04
ALCOHOL	0.40	0.41	0.51	0.50	0.53	0.49
Binge Alcohol Use ³	--	--	--	--	0.34	0.34
Heavy Alcohol Use ³	--	--	--	--	0.15	0.15

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.10P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOBACCO PRODUCTS¹	0.2882		0.1582		0.1724	
Cigarettes	0.2579		0.1440		0.1688	
Smokeless Tobacco	0.1649		0.6749		0.2941	
Cigars	0.1628		0.7691		0.7359	
Pipe Tobacco ²	0.0775		--		0.1415	
ALCOHOL	0.6010		0.5581		0.9702	
Binge Alcohol Use ³	--		--		0.3155	
Heavy Alcohol Use ³	--		--		0.6968	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.