

Table 2.57C (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	681	700	334	351	291	314
GEOGRAPHIC DIVISION						
Northeast	237	250	109	100	100	79
New England	111	129	42	39	37	29
Middle Atlantic	211	213	101	92	94	74
Midwest	276	286	127	135	108	115
East North Central	214	226	96	95	80	85
West North Central	173	178	83	99	67	78
South	435	471	247	247	212	226
South Atlantic	318	343	186	167	153	149
East South Central	173	215	84	124	82	120
West South Central	243	236	148	132	134	116
West	339	365	154	186	137	169
Mountain	201	203	105	142	97	131
Pacific	270	300	112	116	96	105
COUNTY TYPE						
Large Metro	571	596	206	250	177	217
Small Metro	483	511	220	218	193	186
250K – 1 Mil. Pop.	394	404	172	178	149	151
< 250K Pop.	300	323	141	125	125	109
Nonmetro	406	426	203	197	173	180
Urbanized	256	284	116	120	84	106
Less Urbanized	290	315	151	124	143	112
Completely Rural	170	155	77	94	53	88

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.57D (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.27	0.28	0.13	0.14	0.12	0.12
GEOGRAPHIC DIVISION						
Northeast	0.51	0.54	0.24	0.21	0.22	0.17
New England	0.91	1.05	0.34	0.32	0.30	0.24
Middle Atlantic	0.62	0.62	0.30	0.27	0.27	0.22
Midwest	0.50	0.52	0.23	0.24	0.20	0.21
East North Central	0.56	0.59	0.25	0.25	0.21	0.22
West North Central	1.03	1.06	0.49	0.59	0.40	0.47
South	0.47	0.51	0.27	0.27	0.23	0.24
South Atlantic	0.66	0.70	0.38	0.34	0.31	0.30
East South Central	1.15	1.43	0.56	0.82	0.54	0.80
West South Central	0.85	0.82	0.52	0.46	0.47	0.40
West	0.58	0.62	0.26	0.32	0.23	0.29
Mountain	1.13	1.12	0.59	0.79	0.54	0.73
Pacific	0.67	0.74	0.28	0.29	0.24	0.26
COUNTY TYPE						
Large Metro	0.36	0.37	0.14	0.17	0.13	0.15
Small Metro	0.52	0.49	0.28	0.26	0.25	0.22
250K – 1 Mil. Pop.	0.61	0.59	0.32	0.32	0.28	0.27
< 250K Pop.	0.93	0.93	0.55	0.45	0.48	0.39
Nonmetro	0.70	0.79	0.43	0.44	0.38	0.41
Urbanized	1.03	1.15	0.57	0.64	0.42	0.57
Less Urbanized	1.10	1.16	0.69	0.57	0.66	0.52
Completely Rural	1.75	2.43	1.42	1.92	1.04	1.86

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.57P (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.7936		0.7466		0.6285	
GEOGRAPHIC DIVISION						
Northeast	0.5817		0.9679		0.8766	
New England	0.2954		0.8780		0.7222	
Middle Atlantic	0.9680		0.9808		0.9890	
Midwest	0.2950		0.4961		0.6297	
East North Central	0.5654		0.8864		0.9199	
West North Central	0.3284		0.2447		0.3972	
South	0.4226		0.7126		0.6032	
South Atlantic	0.3294		0.9053		0.5377	
East South Central	0.3502		0.3412		0.2372	
West South Central	0.3259		0.7777		0.3575	
West	0.3227		0.5346		0.5315	
Mountain	0.7712		0.1929		0.5637	
Pacific	0.3210		0.6455		0.7791	
COUNTY TYPE						
Large Metro	0.5664		0.3695		0.4362	
Small Metro	0.6084		0.8370		0.9140	
250K – 1 Mil. Pop.	0.8480		0.4605		0.5573	
< 250K Pop.	0.5781		0.5566		0.5830	
Nonmetro	0.5444		0.6627		0.9047	
Urbanized	0.5888		0.5302		0.1094	
Less Urbanized	0.1745		0.0431		0.0405	
Completely Rural	0.7799		0.2167		0.0744	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.58C (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	60	58	48	46	32	30
GEOGRAPHIC DIVISION						
Northeast	18	19	17	17	10	11
New England	8	9	7	9	3	6
Middle Atlantic	16	17	15	15	10	9
Midwest	24	22	20	18	14	12
East North Central	17	16	15	13	11	9
West North Central	16	16	14	13	9	8
South	42	43	32	30	23	22
South Atlantic	32	32	25	23	18	16
East South Central	16	16	13	12	10	9
West South Central	21	20	14	16	10	12
West	31	25	24	23	13	13
Mountain	14	15	10	14	7	8
Pacific	27	20	22	17	11	11
COUNTY TYPE						
Large Metro	43	38	31	33	18	21
Small Metro	38	34	31	25	19	18
250K – 1 Mil. Pop.	30	26	25	21	15	14
< 250K Pop.	24	22	19	16	12	11
Nonmetro	36	34	28	23	20	17
Urbanized	26	20	21	14	14	9
Less Urbanized	23	27	18	18	13	13
Completely Rural	11	9	9	8	6	7

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.58D (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.25	0.24	0.19	0.19	0.13	0.12
GEOGRAPHIC DIVISION						
Northeast	0.42	0.45	0.39	0.41	0.24	0.26
New England	0.73	0.86	0.66	0.82	0.30	0.54
Middle Atlantic	0.51	0.53	0.47	0.47	0.30	0.29
Midwest	0.44	0.41	0.37	0.34	0.26	0.22
East North Central	0.45	0.44	0.39	0.35	0.29	0.24
West North Central	1.01	0.98	0.86	0.81	0.54	0.49
South	0.47	0.48	0.35	0.34	0.26	0.25
South Atlantic	0.70	0.72	0.54	0.51	0.40	0.36
East South Central	1.10	1.14	0.92	0.85	0.71	0.65
West South Central	0.70	0.65	0.48	0.52	0.33	0.40
West	0.52	0.42	0.41	0.39	0.22	0.22
Mountain	0.75	0.82	0.57	0.81	0.38	0.43
Pacific	0.67	0.49	0.53	0.43	0.27	0.26
COUNTY TYPE						
Large Metro	0.30	0.28	0.23	0.25	0.14	0.16
Small Metro	0.43	0.41	0.37	0.32	0.24	0.23
250K – 1 Mil. Pop.	0.49	0.47	0.43	0.39	0.28	0.27
< 250K Pop.	0.88	0.80	0.69	0.58	0.44	0.43
Nonmetro	0.76	0.77	0.62	0.55	0.47	0.41
Urbanized	1.29	1.08	1.07	0.79	0.74	0.51
Less Urbanized	1.08	1.21	0.88	0.83	0.66	0.63
Completely Rural	1.91	2.44	1.67	2.19	1.27	1.90

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.58P (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.2809		0.4888		0.8642	
GEOGRAPHIC DIVISION						
Northeast	0.3417		0.5872		0.3641	
New England	0.1469		0.1375		0.1577	
Middle Atlantic	0.8025		0.7752		0.8063	
Midwest	0.5035		0.4921		0.3379	
East North Central	0.4317		0.3138		0.4101	
West North Central	0.8972		0.9723		0.5950	
South	0.3217		0.5478		0.8372	
South Atlantic	0.7162		0.6138		0.9180	
East South Central	0.9244		0.7972		0.8591	
West South Central	0.1859		0.8322		0.6759	
West	0.3942		0.6867		0.5792	
Mountain	0.8365		0.3136		0.9601	
Pacific	0.2775		0.2869		0.5378	
COUNTY TYPE						
Large Metro	0.2910		0.7235		0.6253	
Small Metro	0.9903		0.5178		0.9200	
250K – 1 Mil. Pop.	0.7136		0.4137		0.9452	
< 250K Pop.	0.6554		0.9538		0.7988	
Nonmetro	0.5829		0.3291		0.5360	
Urbanized	0.2590		0.1221		0.2661	
Less Urbanized	0.8328		0.9255		0.9055	
Completely Rural	0.9028		0.7366		0.9971	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.59C (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	138	137	102	102	78	81
GEOGRAPHIC DIVISION						
Northeast	49	48	35	33	23	25
New England	21	23	14	16	11	13
Middle Atlantic	43	42	32	28	21	22
Midwest	49	54	42	40	35	32
East North Central	39	44	34	32	28	25
West North Central	29	32	24	24	22	19
South	95	92	73	70	54	58
South Atlantic	76	62	57	47	43	38
East South Central	28	42	24	32	18	30
West South Central	51	52	39	40	29	33
West	67	68	49	49	37	39
Mountain	34	29	25	21	20	17
Pacific	57	62	42	44	30	35
COUNTY TYPE						
Large Metro	98	114	65	71	46	54
Small Metro	105	110	73	73	54	52
250K – 1 Mil. Pop.	72	75	48	49	36	37
< 250K Pop.	74	79	55	52	40	38
Nonmetro	111	79	70	57	51	48
Urbanized	97	49	59	29	43	20
Less Urbanized	57	62	38	49	28	43
Completely Rural	17	21	12	14	9	11

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.59D (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.41	0.40	0.30	0.30	0.23	0.24
GEOGRAPHIC DIVISION						
Northeast	0.79	0.80	0.57	0.54	0.38	0.42
New England	1.33	1.42	0.90	0.97	0.68	0.80
Middle Atlantic	0.96	0.96	0.70	0.64	0.46	0.49
Midwest	0.66	0.72	0.57	0.54	0.48	0.42
East North Central	0.76	0.87	0.68	0.63	0.55	0.49
West North Central	1.30	1.34	1.05	1.00	0.96	0.81
South	0.79	0.74	0.60	0.56	0.45	0.46
South Atlantic	1.24	0.99	0.94	0.75	0.70	0.61
East South Central	1.48	2.13	1.25	1.60	0.95	1.49
West South Central	1.28	1.27	0.98	0.97	0.72	0.80
West	0.83	0.84	0.61	0.60	0.45	0.49
Mountain	1.42	1.15	1.04	0.85	0.84	0.70
Pacific	1.01	1.11	0.75	0.78	0.53	0.63
COUNTY TYPE						
Large Metro	0.49	0.52	0.35	0.36	0.25	0.28
Small Metro	0.78	0.71	0.61	0.53	0.46	0.42
250K – 1 Mil. Pop.	0.86	0.83	0.63	0.60	0.49	0.47
< 250K Pop.	1.43	1.29	1.25	0.98	0.93	0.80
Nonmetro	0.94	1.09	0.77	0.90	0.64	0.83
Urbanized	1.44	1.33	1.20	0.95	1.00	0.74
Less Urbanized	1.43	1.76	1.12	1.54	0.92	1.45
Completely Rural	3.72	4.64	3.15	2.99	2.45	2.56

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.59P (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.8966		0.3404		0.3834	
GEOGRAPHIC DIVISION						
Northeast	0.6253		0.3052		0.2757	
New England	0.7921		0.1439		0.0483	
Middle Atlantic	0.4919		0.6758		0.8969	
Midwest	0.5336		0.8221		0.9506	
East North Central	0.7376		0.8218		0.9678	
West North Central	0.5769		0.9818		0.8350	
South	0.7022		0.8340		0.9119	
South Atlantic	0.6032		0.8133		0.6807	
East South Central	0.7222		0.4421		0.9468	
West South Central	0.7831		0.2353		0.5185	
West	0.9866		0.5355		0.3725	
Mountain	0.1335		0.5682		0.7179	
Pacific	0.4760		0.7206		0.2312	
COUNTY TYPE						
Large Metro	0.1439		0.0989		0.4259	
Small Metro	0.1553		0.7230		0.8533	
250K – 1 Mil. Pop.	0.2911		0.8018		0.6171	
< 250K Pop.	0.2846		0.4061		0.3755	
Nonmetro	1.0000		0.4734		0.1336	
Urbanized	0.7831		0.8723		0.5202	
Less Urbanized	0.8820		0.2732		0.0191	
Completely Rural	0.6470		0.7168		0.8979	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.60C (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	657	677	307	333	274	302
GEOGRAPHIC DIVISION						
Northeast	232	247	100	93	95	75
New England	107	127	37	34	34	26
Middle Atlantic	206	212	94	88	88	70
Midwest	270	276	122	127	105	109
East North Central	206	219	89	91	76	82
West North Central	172	169	80	90	67	73
South	419	465	231	238	203	219
South Atlantic	307	337	170	158	142	145
East South Central	170	211	83	119	78	115
West South Central	238	231	138	126	127	111
West	333	355	140	175	129	162
Mountain	200	204	94	138	91	129
Pacific	263	289	103	105	91	97
COUNTY TYPE						
Large Metro	541	571	189	230	167	205
Small Metro	452	476	195	196	176	170
250K – 1 Mil. Pop.	373	383	157	165	140	141
< 250K Pop.	273	294	122	104	110	96
Nonmetro	374	389	185	178	162	165
Urbanized	222	258	96	111	71	101
Less Urbanized	271	281	139	106	137	98
Completely Rural	160	142	74	84	50	80

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.60D (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.34	0.35	0.16	0.17	0.14	0.15
GEOGRAPHIC DIVISION						
Northeast	0.64	0.68	0.28	0.26	0.26	0.21
New England	1.13	1.33	0.39	0.36	0.36	0.28
Middle Atlantic	0.78	0.79	0.35	0.33	0.33	0.26
Midwest	0.64	0.65	0.29	0.30	0.25	0.26
East North Central	0.70	0.74	0.30	0.31	0.26	0.28
West North Central	1.34	1.31	0.63	0.70	0.52	0.56
South	0.59	0.65	0.33	0.33	0.29	0.31
South Atlantic	0.81	0.88	0.45	0.41	0.37	0.38
East South Central	1.46	1.81	0.71	1.02	0.67	0.98
West South Central	1.11	1.06	0.64	0.58	0.59	0.51
West	0.75	0.79	0.32	0.39	0.29	0.36
Mountain	1.47	1.49	0.69	1.00	0.67	0.94
Pacific	0.86	0.93	0.34	0.34	0.30	0.31
COUNTY TYPE						
Large Metro	0.45	0.48	0.18	0.21	0.16	0.19
Small Metro	0.65	0.63	0.33	0.32	0.29	0.28
250K – 1 Mil. Pop.	0.77	0.75	0.39	0.40	0.35	0.34
< 250K Pop.	1.12	1.18	0.63	0.52	0.56	0.47
Nonmetro	0.90	0.95	0.54	0.52	0.48	0.49
Urbanized	1.32	1.42	0.68	0.78	0.50	0.72
Less Urbanized	1.41	1.40	0.86	0.66	0.85	0.61
Completely Rural	2.11	2.79	1.72	2.18	1.25	2.13

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.60P (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.8349		0.9037		0.7980	
GEOGRAPHIC DIVISION						
Northeast	0.5645		0.7105		0.5667	
New England	0.3855		0.4297		0.1673	
Middle Atlantic	0.8672		0.9499		0.9637	
Midwest	0.2401		0.4551		0.6803	
East North Central	0.5324		0.8592		0.8690	
West North Central	0.2649		0.1990		0.4202	
South	0.4975		0.7206		0.6463	
South Atlantic	0.3722		0.8291		0.4663	
East South Central	0.3761		0.2286		0.2381	
West South Central	0.4039		0.5177		0.2401	
West	0.2927		0.5893		0.6432	
Mountain	0.6242		0.2667		0.5244	
Pacific	0.3413		0.6939		0.9806	
COUNTY TYPE						
Large Metro	0.7094		0.7351		0.6077	
Small Metro	0.7588		0.7797		0.9341	
250K – 1 Mil. Pop.	0.9850		0.4487		0.6669	
< 250K Pop.	0.6561		0.5932		0.6788	
Nonmetro	0.5951		0.7245		0.9345	
Urbanized	0.5191		0.2684		0.0364	
Less Urbanized	0.1883		0.0222		0.0096	
Completely Rural	0.7844		0.2040		0.0639	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.61C (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	675	693	333	345	288	310
GEOGRAPHIC DIVISION						
Northeast	235	248	107	98	98	79
New England	109	127	41	37	37	28
Middle Atlantic	210	213	99	91	91	73
Midwest	274	284	127	133	106	114
East North Central	213	225	94	94	79	85
West North Central	173	175	83	96	67	77
South	434	467	249	243	213	221
South Atlantic	317	341	188	162	153	146
East South Central	171	213	83	123	81	118
West South Central	247	232	148	130	133	113
West	338	364	152	183	136	168
Mountain	202	205	104	141	96	131
Pacific	268	299	110	115	95	104
COUNTY TYPE						
Large Metro	563	590	203	245	175	213
Small Metro	479	503	219	212	190	182
250K – 1 Mil. Pop.	390	398	170	174	148	148
< 250K Pop.	296	316	141	119	123	105
Nonmetro	397	416	201	190	172	176
Urbanized	249	276	113	118	81	104
Less Urbanized	284	308	149	118	142	109
Completely Rural	165	151	75	90	51	85

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.61D (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.30	0.30	0.15	0.15	0.13	0.14
GEOGRAPHIC DIVISION						
Northeast	0.56	0.59	0.25	0.23	0.23	0.19
New England	0.99	1.13	0.37	0.33	0.33	0.25
Middle Atlantic	0.68	0.68	0.32	0.29	0.29	0.24
Midwest	0.55	0.57	0.25	0.27	0.21	0.23
East North Central	0.61	0.65	0.27	0.27	0.23	0.24
West North Central	1.15	1.15	0.55	0.63	0.44	0.51
South	0.52	0.56	0.30	0.29	0.26	0.26
South Atlantic	0.72	0.77	0.43	0.36	0.35	0.33
East South Central	1.26	1.57	0.61	0.90	0.59	0.87
West South Central	0.97	0.90	0.58	0.50	0.52	0.44
West	0.65	0.69	0.29	0.35	0.26	0.32
Mountain	1.26	1.26	0.65	0.87	0.60	0.81
Pacific	0.74	0.81	0.30	0.31	0.26	0.28
COUNTY TYPE						
Large Metro	0.40	0.41	0.16	0.19	0.14	0.17
Small Metro	0.57	0.54	0.31	0.28	0.27	0.24
250K – 1 Mil. Pop.	0.68	0.64	0.36	0.35	0.31	0.30
< 250K Pop.	1.01	1.01	0.61	0.48	0.52	0.42
Nonmetro	0.78	0.86	0.48	0.48	0.42	0.44
Urbanized	1.14	1.26	0.63	0.70	0.45	0.62
Less Urbanized	1.21	1.28	0.76	0.61	0.74	0.57
Completely Rural	1.92	2.57	1.57	2.04	1.14	1.97

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.61P (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.8226		0.6745		0.6229	
GEOGRAPHIC DIVISION						
Northeast	0.6456		0.9684		0.7831	
New England	0.3493		0.8767		0.5623	
Middle Atlantic	0.9959		0.9799		0.9641	
Midwest	0.3021		0.5533		0.6986	
East North Central	0.5892		0.7801		0.8470	
West North Central	0.3138		0.2376		0.4231	
South	0.4614		0.6596		0.6212	
South Atlantic	0.3374		0.8564		0.5324	
East South Central	0.3450		0.3168		0.2367	
West South Central	0.3816		0.7937		0.3317	
West	0.3003		0.4949		0.5036	
Mountain	0.7972		0.2333		0.5659	
Pacific	0.2817		0.7829		0.7284	
COUNTY TYPE						
Large Metro	0.5251		0.3928		0.4702	
Small Metro	0.5709		0.7708		0.9154	
250K – 1 Mil. Pop.	0.8323		0.3964		0.5662	
< 250K Pop.	0.5344		0.5487		0.5894	
Nonmetro	0.5686		0.7501		0.8553	
Urbanized	0.5507		0.4060		0.0823	
Less Urbanized	0.1821		0.0418		0.0397	
Completely Rural	0.7569		0.2192		0.0684	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.