

**Table 2.62C (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	876	929	501	510	361	363
GEOGRAPHIC DIVISION						
Northeast	377	365	223	213	154	151
New England	194	176	104	112	80	89
Middle Atlantic	321	315	197	180	131	121
Midwest	334	352	221	220	153	159
East North Central	276	287	177	173	120	124
West North Central	174	203	129	133	95	97
South	541	602	295	316	210	235
South Atlantic	430	450	230	237	169	171
East South Central	209	223	92	123	75	94
West South Central	278	301	159	166	111	135
West	466	508	261	256	183	161
Mountain	228	237	140	117	96	76
Pacific	408	447	220	229	158	143
COUNTY TYPE						
Large Metro	1,013	1,046	421	449	287	298
Small Metro	697	800	295	322	198	197
250K – 1 Mil. Pop.	590	646	258	253	161	151
< 250K Pop.	448	481	179	202	134	133
Nonmetro	545	499	211	213	147	147
Urbanized	387	363	158	126	108	71
Less Urbanized	349	382	135	173	93	126
Completely Rural	235	163	57	49	35	38

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.62D (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.35	0.37	0.20	0.20	0.14	0.14
GEOGRAPHIC DIVISION						
Northeast	0.81	0.78	0.48	0.46	0.33	0.32
New England	1.59	1.44	0.85	0.91	0.66	0.73
Middle Atlantic	0.94	0.92	0.58	0.53	0.38	0.35
Midwest	0.60	0.64	0.40	0.40	0.28	0.29
East North Central	0.72	0.74	0.46	0.45	0.31	0.32
West North Central	1.04	1.21	0.77	0.79	0.57	0.58
South	0.59	0.65	0.32	0.34	0.23	0.25
South Atlantic	0.89	0.92	0.47	0.48	0.35	0.35
East South Central	1.40	1.48	0.61	0.81	0.50	0.63
West South Central	0.98	1.04	0.56	0.57	0.39	0.47
West	0.80	0.86	0.45	0.44	0.31	0.27
Mountain	1.28	1.31	0.79	0.65	0.54	0.42
Pacific	1.01	1.10	0.54	0.56	0.39	0.35
COUNTY TYPE						
Large Metro	0.50	0.55	0.29	0.29	0.21	0.21
Small Metro	0.62	0.61	0.36	0.36	0.25	0.24
250K – 1 Mil. Pop.	0.74	0.78	0.45	0.43	0.30	0.28
< 250K Pop.	1.20	1.03	0.65	0.65	0.49	0.47
Nonmetro	0.83	0.77	0.47	0.45	0.34	0.34
Urbanized	1.38	1.17	0.75	0.63	0.54	0.39
Less Urbanized	1.20	1.18	0.63	0.70	0.46	0.55
Completely Rural	2.34	2.51	1.19	1.18	0.73	0.93

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.62P (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.0631		0.2988		0.7452	
GEOGRAPHIC DIVISION						
Northeast	0.6001		0.4284		0.7950	
New England	0.7072		0.6143		0.9366	
Middle Atlantic	0.6896		0.2309		0.7971	
Midwest	0.8136		0.3747		0.1450	
East North Central	0.9530		0.8783		0.1308	
West North Central	0.7311		0.0750		0.6595	
South	0.6903		0.2910		0.6917	
South Atlantic	0.8140		0.3489		0.7469	
East South Central	0.6968		0.0535		0.2740	
West South Central	0.4029		0.4569		0.6247	
West	0.0135		0.0770		0.0057	
Mountain	0.0564		0.3568		0.2420	
Pacific	0.0848		0.1372		0.0128	
COUNTY TYPE						
Large Metro	0.0973		0.2181		0.2833	
Small Metro	0.8335		0.9612		0.5506	
250K – 1 Mil. Pop.	0.6021		0.7815		0.3793	
< 250K Pop.	0.7093		0.6345		0.8867	
Nonmetro	0.0671		0.7536		0.6064	
Urbanized	0.8904		0.9887		0.6275	
Less Urbanized	0.2454		0.8326		0.4168	
Completely Rural	0.0036		0.4342		0.5345	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.63C (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	74	65	61	55	39	36
GEOGRAPHIC DIVISION						
Northeast	29	26	25	23	17	14
New England	14	14	13	11	11	7
Middle Atlantic	25	22	22	20	13	13
Midwest	27	24	23	21	17	14
East North Central	23	20	19	18	14	12
West North Central	15	14	13	11	10	8
South	44	41	37	35	25	25
South Atlantic	33	29	28	24	20	19
East South Central	14	16	12	13	9	10
West South Central	24	25	19	22	13	12
West	42	36	38	30	20	17
Mountain	17	16	14	14	9	8
Pacific	39	33	35	27	18	15
COUNTY TYPE						
Large Metro	62	55	52	46	31	27
Small Metro	44	39	35	31	23	21
250K – 1 Mil. Pop.	36	33	29	27	20	18
< 250K Pop.	25	22	21	17	12	10
Nonmetro	29	29	24	23	17	14
Urbanized	21	16	19	13	13	7
Less Urbanized	19	24	15	18	11	11
Completely Rural	9	7	6	6	4	5

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.63D (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.30	0.27	0.25	0.22	0.16	0.15
GEOGRAPHIC DIVISION						
Northeast	0.68	0.62	0.59	0.55	0.40	0.34
New England	1.28	1.27	1.15	1.04	1.02	0.63
Middle Atlantic	0.80	0.70	0.68	0.64	0.40	0.40
Midwest	0.51	0.45	0.43	0.39	0.32	0.26
East North Central	0.61	0.53	0.51	0.47	0.37	0.31
West North Central	0.92	0.87	0.81	0.68	0.63	0.49
South	0.49	0.46	0.41	0.39	0.28	0.28
South Atlantic	0.72	0.65	0.62	0.53	0.43	0.42
East South Central	0.97	1.10	0.80	0.93	0.60	0.70
West South Central	0.79	0.82	0.64	0.74	0.42	0.40
West	0.72	0.63	0.64	0.52	0.34	0.30
Mountain	0.93	0.87	0.76	0.79	0.50	0.46
Pacific	0.95	0.82	0.85	0.66	0.44	0.38
COUNTY TYPE						
Large Metro	0.44	0.38	0.37	0.33	0.23	0.20
Small Metro	0.51	0.45	0.42	0.38	0.29	0.26
250K – 1 Mil. Pop.	0.63	0.57	0.52	0.49	0.37	0.35
< 250K Pop.	0.84	0.75	0.71	0.63	0.45	0.41
Nonmetro	0.61	0.69	0.54	0.55	0.40	0.35
Urbanized	0.93	0.87	0.91	0.74	0.66	0.44
Less Urbanized	0.89	1.13	0.74	0.81	0.55	0.52
Completely Rural	2.07	2.00	1.43	1.88	1.01	1.58

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.63P (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.0002		0.0001		0.0005	
GEOGRAPHIC DIVISION						
Northeast	0.2681		0.3095		0.3714	
New England	0.9880		0.3870		0.3608	
Middle Atlantic	0.1865		0.4926		0.6784	
Midwest	0.0014		0.0005		0.0011	
East North Central	0.0007		0.0032		0.0036	
West North Central	0.5771		0.0717		0.1212	
South	0.0421		0.1980		0.1074	
South Atlantic	0.0961		0.3376		0.4721	
East South Central	0.3250		0.8679		0.9280	
West South Central	0.4324		0.3399		0.0291	
West	0.1413		0.0288		0.1276	
Mountain	0.2632		0.2248		0.4979	
Pacific	0.2635		0.0622		0.1676	
COUNTY TYPE						
Large Metro	0.0009		0.0074		0.0084	
Small Metro	0.9160		0.0972		0.2541	
250K – 1 Mil. Pop.	0.4714		0.7132		0.8212	
< 250K Pop.	0.3552		0.0157		0.0574	
Nonmetro	0.0016		0.0201		0.0151	
Urbanized	0.0100		0.0337		0.0018	
Less Urbanized	0.0521		0.0712		0.2200	
Completely Rural	0.3843		0.5535		0.4345	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.64C (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	165	159	137	134	99	101
GEOGRAPHIC DIVISION						
Northeast	68	66	54	54	41	38
New England	34	33	25	27	22	20
Middle Atlantic	58	57	48	46	34	32
Midwest	62	60	53	51	41	41
East North Central	50	50	44	43	34	34
West North Central	36	35	29	29	22	26
South	105	103	86	89	62	73
South Atlantic	88	87	69	72	47	60
East South Central	31	35	27	31	20	26
West South Central	44	50	42	47	34	35
West	88	89	74	71	50	44
Mountain	43	42	29	29	20	22
Pacific	77	77	67	65	46	38
COUNTY TYPE						
Large Metro	180	189	121	121	78	78
Small Metro	151	183	103	124	61	73
250K – 1 Mil. Pop.	110	127	71	88	45	50
< 250K Pop.	104	133	77	88	42	56
Nonmetro	155	97	108	69	62	48
Urbanized	142	70	95	48	51	31
Less Urbanized	67	70	51	52	34	34
Completely Rural	23	24	17	14	11	12

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.64D (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.49	0.47	0.41	0.39	0.29	0.30
GEOGRAPHIC DIVISION						
Northeast	1.11	1.09	0.89	0.89	0.66	0.63
New England	2.15	2.06	1.56	1.65	1.36	1.26
Middle Atlantic	1.29	1.28	1.06	1.05	0.75	0.72
Midwest	0.85	0.81	0.73	0.69	0.56	0.55
East North Central	0.98	0.97	0.86	0.85	0.66	0.65
West North Central	1.61	1.47	1.29	1.24	0.96	1.09
South	0.87	0.83	0.72	0.72	0.51	0.59
South Atlantic	1.44	1.38	1.14	1.13	0.77	0.94
East South Central	1.61	1.75	1.41	1.54	1.05	1.30
West South Central	1.10	1.22	1.06	1.13	0.85	0.86
West	1.09	1.10	0.91	0.88	0.62	0.55
Mountain	1.77	1.70	1.20	1.15	0.82	0.87
Pacific	1.35	1.37	1.19	1.16	0.82	0.69
COUNTY TYPE						
Large Metro	0.67	0.66	0.54	0.53	0.40	0.40
Small Metro	0.82	0.81	0.68	0.68	0.48	0.52
250K – 1 Mil. Pop.	0.95	0.95	0.76	0.82	0.59	0.60
< 250K Pop.	1.46	1.53	1.30	1.26	0.81	1.04
Nonmetro	1.21	1.05	1.06	1.00	0.76	0.81
Urbanized	1.83	1.50	1.66	1.44	1.08	1.15
Less Urbanized	1.47	1.62	1.35	1.51	1.10	1.17
Completely Rural	4.14	4.12	3.80	2.90	2.76	2.80

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.64P (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.1035		0.1228		0.6074	
GEOGRAPHIC DIVISION						
Northeast	0.9174		0.8349		0.6052	
New England	0.9443		0.8775		0.6844	
Middle Atlantic	0.9647		0.7699		0.4090	
Midwest	0.9027		0.7350		0.4536	
East North Central	0.5427		0.6149		0.5490	
West North Central	0.4918		0.8706		0.6455	
South	0.1334		0.1448		0.7944	
South Atlantic	0.0884		0.2323		0.7400	
East South Central	0.5435		0.6475		0.9158	
West South Central	0.5198		0.1742		0.9222	
West	0.2247		0.2669		0.0827	
Mountain	0.9373		0.6028		0.9322	
Pacific	0.1473		0.1451		0.0568	
COUNTY TYPE						
Large Metro	0.0286		0.0300		0.1337	
Small Metro	0.5534		0.9809		0.5476	
250K – 1 Mil. Pop.	0.6601		0.3346		0.2640	
< 250K Pop.	0.1354		0.2374		0.6300	
Nonmetro	0.3625		0.9164		0.3981	
Urbanized	0.2570		0.7277		0.5175	
Less Urbanized	0.5811		0.7705		0.5106	
Completely Rural	0.8262		0.3193		0.9880	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.65C (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	843	896	458	482	341	340
GEOGRAPHIC DIVISION						
Northeast	361	352	202	201	145	143
New England	184	172	93	103	73	83
Middle Atlantic	310	302	180	172	126	116
Midwest	323	339	202	202	144	146
East North Central	269	276	161	158	113	115
West North Central	171	197	118	123	88	89
South	524	578	264	296	196	221
South Atlantic	409	432	196	221	153	159
East South Central	209	217	82	112	70	86
West South Central	272	294	150	162	107	131
West	450	494	244	239	176	156
Mountain	217	238	138	110	92	70
Pacific	394	432	202	213	152	139
COUNTY TYPE						
Large Metro	962	987	384	412	273	277
Small Metro	658	736	265	284	179	181
250K – 1 Mil. Pop.	552	606	236	228	148	144
< 250K Pop.	423	431	154	172	120	114
Nonmetro	496	460	169	187	129	129
Urbanized	340	327	114	108	91	61
Less Urbanized	320	346	115	147	84	110
Completely Rural	223	152	51	44	33	35

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.65D (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.44	0.46	0.24	0.25	0.18	0.17
GEOGRAPHIC DIVISION						
Northeast	1.01	0.97	0.56	0.55	0.40	0.39
New England	1.94	1.80	0.98	1.08	0.77	0.87
Middle Atlantic	1.17	1.13	0.68	0.64	0.48	0.44
Midwest	0.76	0.80	0.48	0.47	0.34	0.34
East North Central	0.91	0.93	0.55	0.53	0.38	0.39
West North Central	1.33	1.53	0.92	0.95	0.69	0.69
South	0.74	0.81	0.37	0.41	0.28	0.31
South Atlantic	1.08	1.13	0.52	0.58	0.40	0.42
East South Central	1.79	1.86	0.70	0.96	0.60	0.74
West South Central	1.27	1.35	0.70	0.74	0.50	0.60
West	1.02	1.10	0.55	0.53	0.40	0.35
Mountain	1.59	1.73	1.01	0.80	0.67	0.51
Pacific	1.29	1.39	0.66	0.69	0.49	0.45
COUNTY TYPE						
Large Metro	0.63	0.70	0.35	0.35	0.26	0.26
Small Metro	0.78	0.76	0.43	0.43	0.30	0.29
250K – 1 Mil. Pop.	0.94	0.97	0.55	0.52	0.36	0.35
< 250K Pop.	1.50	1.29	0.76	0.79	0.60	0.55
Nonmetro	1.03	0.95	0.51	0.53	0.39	0.39
Urbanized	1.72	1.46	0.77	0.75	0.64	0.45
Less Urbanized	1.49	1.44	0.73	0.82	0.55	0.65
Completely Rural	2.75	2.93	1.37	1.35	0.87	1.07

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.65P (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.1502		0.8067		0.8804	
GEOGRAPHIC DIVISION						
Northeast	0.6130		0.4615		0.7944	
New England	0.6653		0.5040		0.7621	
Middle Atlantic	0.7339		0.2219		0.9027	
Midwest	0.9667		0.5680		0.1033	
East North Central	0.7746		0.5428		0.0842	
West North Central	0.6279		0.0620		0.6491	
South	0.9516		0.1063		0.5265	
South Atlantic	0.5166		0.1547		0.6182	
East South Central	0.6527		0.0482		0.2635	
West South Central	0.4679		0.7733		0.7873	
West	0.0271		0.2234		0.0346	
Mountain	0.0533		0.3179		0.2380	
Pacific	0.1575		0.4289		0.0811	
COUNTY TYPE						
Large Metro	0.2519		0.6670		0.6412	
Small Metro	0.8580		0.9964		0.6666	
250K – 1 Mil. Pop.	0.7638		0.5515		0.5815	
< 250K Pop.	0.9120		0.3910		0.9939	
Nonmetro	0.0773		0.8150		0.4305	
Urbanized	0.8747		0.5506		0.9288	
Less Urbanized	0.2918		0.9412		0.3543	
Completely Rural	0.0029		0.4091		0.6847	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.66C (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	859	921	497	504	358	358
GEOGRAPHIC DIVISION						
Northeast	372	362	220	210	150	149
New England	189	173	100	110	76	88
Middle Atlantic	319	314	197	178	129	120
Midwest	332	349	218	217	153	157
East North Central	275	284	174	170	120	122
West North Central	173	204	127	133	95	97
South	535	597	293	315	210	232
South Atlantic	425	445	227	235	169	167
East South Central	208	221	91	122	74	94
West South Central	272	300	157	165	110	134
West	457	507	260	252	181	163
Mountain	223	239	139	113	96	76
Pacific	400	446	219	226	155	145
COUNTY TYPE						
Large Metro	1,004	1,036	416	439	283	293
Small Metro	689	791	290	316	194	195
250K – 1 Mil. Pop.	584	639	254	247	158	150
< 250K Pop.	443	473	176	198	132	131
Nonmetro	539	493	207	207	145	145
Urbanized	383	356	155	121	107	70
Less Urbanized	344	377	131	168	92	123
Completely Rural	232	160	55	47	34	37

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.66D (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.38	0.40	0.22	0.22	0.16	0.16
GEOGRAPHIC DIVISION						
Northeast	0.88	0.86	0.52	0.50	0.36	0.35
New England	1.71	1.55	0.90	0.98	0.69	0.79
Middle Atlantic	1.03	1.01	0.63	0.57	0.42	0.39
Midwest	0.67	0.70	0.44	0.43	0.31	0.31
East North Central	0.79	0.82	0.50	0.49	0.35	0.35
West North Central	1.15	1.34	0.84	0.88	0.63	0.64
South	0.64	0.71	0.35	0.37	0.25	0.28
South Atlantic	0.97	1.00	0.52	0.53	0.38	0.38
East South Central	1.54	1.62	0.67	0.90	0.54	0.69
West South Central	1.07	1.16	0.62	0.64	0.43	0.52
West	0.87	0.96	0.50	0.48	0.35	0.31
Mountain	1.39	1.47	0.87	0.70	0.60	0.47
Pacific	1.10	1.22	0.60	0.62	0.43	0.39
COUNTY TYPE						
Large Metro	0.55	0.61	0.32	0.31	0.23	0.23
Small Metro	0.67	0.66	0.39	0.39	0.27	0.26
250K – 1 Mil. Pop.	0.80	0.85	0.49	0.47	0.32	0.31
< 250K Pop.	1.30	1.12	0.72	0.71	0.54	0.51
Nonmetro	0.91	0.85	0.51	0.49	0.37	0.37
Urbanized	1.51	1.30	0.82	0.69	0.60	0.43
Less Urbanized	1.32	1.29	0.69	0.76	0.51	0.60
Completely Rural	2.47	2.67	1.28	1.25	0.80	1.00

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.66P (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.0921		0.5446		0.9730	
GEOGRAPHIC DIVISION						
Northeast	0.6265		0.4880		0.7245	
New England	0.6778		0.5349		0.8458	
Middle Atlantic	0.7362		0.2559		0.7680	
Midwest	0.9614		0.5846		0.0733	
East North Central	0.8616		0.6421		0.0695	
West North Central	0.7311		0.1013		0.5501	
South	0.7683		0.2252		0.5608	
South Atlantic	0.7455		0.2924		0.6836	
East South Central	0.7328		0.0489		0.2629	
West South Central	0.4182		0.5310		0.7815	
West	0.0154		0.1321		0.0090	
Mountain	0.0568		0.4075		0.2678	
Pacific	0.0937		0.2148		0.0190	
COUNTY TYPE						
Large Metro	0.1425		0.3483		0.4060	
Small Metro	0.9119		0.8276		0.4768	
250K – 1 Mil. Pop.	0.6949		0.8039		0.3697	
< 250K Pop.	0.7278		0.4766		0.9951	
Nonmetro	0.1053		0.9795		0.4304	
Urbanized	0.9560		0.7936		0.8897	
Less Urbanized	0.3444		0.9947		0.3457	
Completely Rural	0.0024		0.3640		0.6053	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.