

**Table 2.67C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	656	715	896	967	957	990
GEOGRAPHIC DIVISION						
Northeast	261	274	390	362	430	403
New England	104	133	167	169	187	204
Middle Atlantic	238	237	350	317	386	340
Midwest	251	254	367	343	379	380
East North Central	204	221	312	285	318	311
West North Central	141	127	193	211	209	239
South	454	470	600	653	638	658
South Atlantic	362	352	469	493	500	495
East South Central	163	213	212	281	203	266
West South Central	228	225	297	322	326	353
West	355	415	501	551	513	549
Mountain	149	183	210	262	212	246
Pacific	321	373	455	480	467	492
COUNTY TYPE						
Large Metro	1,862	1,846	1,645	1,627	1,404	1,389
Small Metro	1,349	1,458	1,101	1,233	947	1,023
250K – 1 Mil. Pop.	1,147	1,185	947	1,006	812	831
< 250K Pop.	886	859	713	718	591	604
Nonmetro	946	980	757	819	649	685
Urbanized	694	760	613	654	513	554
Less Urbanized	657	723	491	576	409	461
Completely Rural	411	337	274	250	237	203

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.67D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.26	0.28	0.36	0.38	0.38	0.39
GEOGRAPHIC DIVISION						
Northeast	0.56	0.59	0.84	0.78	0.93	0.87
New England	0.86	1.09	1.37	1.38	1.54	1.67
Middle Atlantic	0.70	0.69	1.02	0.93	1.13	0.99
Midwest	0.45	0.46	0.67	0.62	0.69	0.69
East North Central	0.53	0.57	0.81	0.74	0.83	0.81
West North Central	0.84	0.76	1.16	1.25	1.25	1.42
South	0.49	0.51	0.65	0.70	0.69	0.71
South Atlantic	0.75	0.72	0.97	1.01	1.03	1.01
East South Central	1.09	1.42	1.41	1.86	1.36	1.77
West South Central	0.80	0.78	1.04	1.12	1.14	1.22
West	0.61	0.71	0.86	0.94	0.88	0.93
Mountain	0.83	1.02	1.18	1.45	1.19	1.37
Pacific	0.79	0.92	1.13	1.18	1.16	1.21
COUNTY TYPE						
Large Metro	0.38	0.41	0.52	0.53	0.56	0.56
Small Metro	0.47	0.50	0.65	0.65	0.69	0.70
250K – 1 Mil. Pop.	0.53	0.54	0.80	0.75	0.88	0.83
< 250K Pop.	0.89	1.00	1.14	1.27	1.13	1.30
Nonmetro	0.63	0.69	0.90	0.96	0.89	0.95
Urbanized	1.01	0.97	1.46	1.59	1.49	1.54
Less Urbanized	0.96	1.04	1.27	1.35	1.24	1.28
Completely Rural	1.87	2.20	2.68	2.96	2.71	2.89

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.67P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.4262		0.4509		0.8783	
GEOGRAPHIC DIVISION						
Northeast	0.9208		0.6104		0.4221	
New England	0.5953		0.6523		0.4723	
Middle Atlantic	0.6844		0.7324		0.5978	
Midwest	0.6014		0.5949		0.9775	
East North Central	0.8268		0.2342		0.3807	
West North Central	0.5277		0.4112		0.1965	
South	0.9005		0.4605		1.0000	
South Atlantic	0.5101		0.6711		0.5211	
East South Central	0.9204		0.9842		0.7773	
West South Central	0.3874		0.4100		0.2083	
West	0.0959		0.6750		0.3799	
Mountain	0.0881		0.3282		0.2627	
Pacific	0.2740		0.9888		0.6670	
COUNTY TYPE						
Large Metro	0.9300		0.4707		0.8254	
Small Metro	0.1894		0.8054		0.8491	
250K – 1 Mil. Pop.	0.2209		0.5633		0.7634	
< 250K Pop.	0.5391		0.2613		0.4801	
Nonmetro	0.7096		0.7358		0.8764	
Urbanized	0.9760		0.2504		0.7724	
Less Urbanized	0.9030		0.7145		0.9265	
Completely Rural	0.4435		0.3681		0.7760	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.68C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	112	108	103	104	79	80
GEOGRAPHIC DIVISION						
Northeast	42	40	39	37	32	31
New England	19	19	18	17	15	14
Middle Atlantic	37	35	35	33	28	28
Midwest	41	40	38	38	29	30
East North Central	34	33	32	30	24	22
West North Central	24	24	21	24	17	21
South	67	72	60	68	46	53
South Atlantic	53	52	47	51	34	41
East South Central	22	23	21	22	15	16
West South Central	36	40	34	39	27	28
West	63	62	58	60	43	45
Mountain	28	27	28	25	19	19
Pacific	56	55	51	54	38	40
COUNTY TYPE						
Large Metro	128	118	109	105	69	73
Small Metro	102	87	82	74	54	46
250K – 1 Mil. Pop.	80	71	64	61	42	36
< 250K Pop.	65	52	52	43	34	28
Nonmetro	60	61	50	55	31	36
Urbanized	44	43	37	39	20	23
Less Urbanized	41	46	34	41	22	27
Completely Rural	19	15	15	13	9	9

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.68D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.45	0.44	0.42	0.43	0.32	0.33
GEOGRAPHIC DIVISION						
Northeast	0.97	0.93	0.91	0.88	0.74	0.73
New England	1.67	1.73	1.59	1.56	1.34	1.26
Middle Atlantic	1.16	1.10	1.09	1.05	0.89	0.88
Midwest	0.76	0.75	0.71	0.70	0.54	0.56
East North Central	0.89	0.87	0.85	0.80	0.62	0.58
West North Central	1.48	1.52	1.29	1.52	1.06	1.32
South	0.74	0.80	0.67	0.76	0.52	0.59
South Atlantic	1.18	1.16	1.04	1.14	0.74	0.91
East South Central	1.54	1.62	1.45	1.51	1.02	1.08
West South Central	1.19	1.34	1.13	1.28	0.91	0.94
West	1.07	1.06	0.99	1.03	0.72	0.77
Mountain	1.56	1.52	1.56	1.37	1.07	1.09
Pacific	1.37	1.37	1.25	1.35	0.93	1.00
COUNTY TYPE						
Large Metro	0.67	0.64	0.61	0.62	0.46	0.50
Small Metro	0.79	0.76	0.72	0.73	0.55	0.53
250K – 1 Mil. Pop.	0.92	0.92	0.82	0.90	0.65	0.62
< 250K Pop.	1.37	1.29	1.32	1.25	1.01	0.98
Nonmetro	0.95	1.01	0.89	0.96	0.68	0.79
Urbanized	1.54	1.50	1.41	1.41	1.00	0.96
Less Urbanized	1.42	1.51	1.33	1.50	1.01	1.28
Completely Rural	2.64	3.92	2.16	3.41	1.76	2.60

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.68P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.0000		0.0011		0.0088	
GEOGRAPHIC DIVISION						
Northeast	0.2817		0.5774		0.3564	
New England	0.9802		0.7992		0.5759	
Middle Atlantic	0.2198		0.4541		0.1945	
Midwest	0.0005		0.0012		0.0141	
East North Central	0.0000		0.0001		0.0002	
West North Central	0.6142		0.9674		0.6072	
South	0.0725		0.3130		0.0958	
South Atlantic	0.8661		0.8349		0.4672	
East South Central	0.1414		0.2286		0.4078	
West South Central	0.0237		0.1889		0.1232	
West	0.0010		0.0497		0.6129	
Mountain	0.1694		0.2331		0.3313	
Pacific	0.0023		0.1089		0.9140	
COUNTY TYPE						
Large Metro	0.0000		0.0022		0.0788	
Small Metro	0.2526		0.3833		0.0521	
250K – 1 Mil. Pop.	0.5585		0.8781		0.4429	
< 250K Pop.	0.2416		0.2258		0.0327	
Nonmetro	0.0696		0.2535		0.4717	
Urbanized	0.0079		0.0261		0.3765	
Less Urbanized	0.8536		0.6536		0.6413	
Completely Rural	0.8950		0.9079		0.7477	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.69C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	121	114	142	138	174	169
GEOGRAPHIC DIVISION						
Northeast	37	44	47	54	69	68
New England	16	22	23	30	39	36
Middle Atlantic	34	38	41	46	57	58
Midwest	49	45	57	50	71	64
East North Central	38	36	44	39	58	50
West North Central	32	28	36	32	41	41
South	77	76	95	93	113	111
South Atlantic	64	61	77	74	90	86
East South Central	27	26	31	32	35	40
West South Central	33	38	43	44	55	55
West	74	63	80	79	96	93
Mountain	38	33	40	39	43	45
Pacific	63	53	68	68	86	80
COUNTY TYPE						
Large Metro	298	344	287	330	246	275
Small Metro	247	300	236	284	207	243
250K – 1 Mil. Pop.	193	219	181	203	155	179
< 250K Pop.	164	201	156	193	141	164
Nonmetro	253	150	247	139	214	121
Urbanized	230	114	227	106	197	92
Less Urbanized	119	109	110	101	92	85
Completely Rural	35	37	32	33	26	28

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.69D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.36	0.34	0.42	0.41	0.52	0.50
GEOGRAPHIC DIVISION						
Northeast	0.60	0.72	0.77	0.90	1.13	1.13
New England	0.99	1.36	1.46	1.82	2.45	2.22
Middle Atlantic	0.75	0.86	0.91	1.03	1.26	1.31
Midwest	0.67	0.60	0.78	0.67	0.97	0.85
East North Central	0.75	0.69	0.86	0.77	1.13	0.98
West North Central	1.40	1.17	1.59	1.38	1.80	1.72
South	0.64	0.62	0.79	0.75	0.94	0.89
South Atlantic	1.04	0.97	1.26	1.18	1.48	1.36
East South Central	1.44	1.33	1.61	1.61	1.86	2.04
West South Central	0.83	0.93	1.08	1.08	1.37	1.33
West	0.91	0.78	0.99	0.98	1.18	1.14
Mountain	1.58	1.31	1.67	1.57	1.76	1.80
Pacific	1.10	0.95	1.19	1.21	1.51	1.42
COUNTY TYPE						
Large Metro	0.49	0.52	0.59	0.61	0.73	0.72
Small Metro	0.63	0.55	0.68	0.69	0.88	0.87
250K – 1 Mil. Pop.	0.83	0.67	0.87	0.86	1.07	1.03
< 250K Pop.	1.00	0.97	1.14	1.18	1.51	1.60
Nonmetro	0.88	0.78	1.15	0.94	1.47	1.18
Urbanized	1.23	1.17	1.84	1.38	2.40	1.68
Less Urbanized	1.21	1.18	1.32	1.54	1.63	1.76
Completely Rural	2.73	2.37	3.22	3.16	3.41	4.36

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.69P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.8202		0.8268		0.5434	
GEOGRAPHIC DIVISION						
Northeast	0.3803		0.2786		0.3758	
New England	0.1116		0.0289		0.6165	
Middle Atlantic	0.8701		0.9795		0.4368	
Midwest	0.8523		0.2727		0.7153	
East North Central	0.6749		0.6807		0.6504	
West North Central	0.7833		0.1985		0.1812	
South	0.2547		0.4974		0.4352	
South Atlantic	0.4502		0.5152		0.9891	
East South Central	0.9416		0.8184		0.5615	
West South Central	0.2259		0.6207		0.2894	
West	0.1724		0.6160		0.7543	
Mountain	0.2587		0.1583		0.3065	
Pacific	0.3272		0.8979		0.8921	
COUNTY TYPE						
Large Metro	0.6748		0.7255		0.8300	
Small Metro	0.5615		0.9907		0.8013	
250K – 1 Mil. Pop.	0.1690		0.6156		0.5959	
< 250K Pop.	0.3520		0.4851		0.2778	
Nonmetro	0.5133		0.1588		0.3092	
Urbanized	0.8682		0.5508		0.9125	
Less Urbanized	0.5434		0.2449		0.1510	
Completely Rural	0.9873		0.5974		0.6493	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.70C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	615	680	870	925	931	946
GEOGRAPHIC DIVISION						
Northeast	247	259	386	354	422	394
New England	92	124	163	165	174	196
Middle Atlantic	228	225	348	310	381	334
Midwest	230	233	356	333	366	357
East North Central	191	204	305	275	307	293
West North Central	124	122	186	207	204	231
South	441	455	582	620	611	621
South Atlantic	349	332	445	464	467	466
East South Central	159	210	208	273	197	256
West South Central	222	219	292	316	321	344
West	324	385	473	522	483	516
Mountain	117	169	185	245	195	228
Pacific	300	348	436	457	441	466
COUNTY TYPE						
Large Metro	1,762	1,725	1,549	1,518	1,333	1,317
Small Metro	1,213	1,319	993	1,112	867	932
250K – 1 Mil. Pop.	1,032	1,083	855	924	748	771
< 250K Pop.	787	769	635	634	528	540
Nonmetro	842	887	658	736	575	631
Urbanized	588	681	511	581	438	501
Less Urbanized	587	649	436	511	371	422
Completely Rural	379	309	246	230	218	188

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.70D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.32	0.35	0.45	0.47	0.48	0.48
GEOGRAPHIC DIVISION						
Northeast	0.69	0.71	1.07	0.98	1.17	1.09
New England	0.97	1.30	1.72	1.73	1.83	2.06
Middle Atlantic	0.86	0.84	1.31	1.16	1.44	1.25
Midwest	0.54	0.55	0.84	0.78	0.86	0.84
East North Central	0.64	0.69	1.03	0.93	1.04	0.99
West North Central	0.96	0.95	1.45	1.61	1.59	1.80
South	0.62	0.63	0.82	0.87	0.86	0.87
South Atlantic	0.92	0.87	1.17	1.21	1.23	1.22
East South Central	1.37	1.80	1.79	2.34	1.69	2.20
West South Central	1.03	1.01	1.36	1.45	1.49	1.58
West	0.73	0.86	1.07	1.16	1.09	1.15
Mountain	0.86	1.23	1.36	1.78	1.44	1.66
Pacific	0.98	1.12	1.42	1.47	1.44	1.50
COUNTY TYPE						
Large Metro	0.47	0.50	0.65	0.67	0.70	0.70
Small Metro	0.59	0.61	0.84	0.82	0.88	0.87
250K – 1 Mil. Pop.	0.64	0.67	1.04	0.95	1.13	1.03
< 250K Pop.	1.14	1.26	1.47	1.62	1.42	1.66
Nonmetro	0.75	0.82	1.08	1.16	1.07	1.16
Urbanized	1.22	1.17	1.73	1.92	1.75	1.87
Less Urbanized	1.16	1.25	1.58	1.66	1.52	1.58
Completely Rural	2.24	2.75	3.14	3.73	3.16	3.54

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.70P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.8182		0.5975		0.9800	
GEOGRAPHIC DIVISION						
Northeast	0.7778		0.7176		0.3318	
New England	0.6809		0.8330		0.4589	
Middle Atlantic	0.5902		0.7661		0.4712	
Midwest	0.3282		0.5233		0.9876	
East North Central	0.4011		0.1286		0.2250	
West North Central	0.5980		0.2681		0.1044	
South	0.7978		0.5142		0.8504	
South Atlantic	0.5488		0.6600		0.5064	
East South Central	0.9837		0.9967		0.6820	
West South Central	0.1496		0.5174		0.3063	
West	0.1144		0.7351		0.3192	
Mountain	0.0538		0.2445		0.1735	
Pacific	0.3604		0.8750		0.6480	
COUNTY TYPE						
Large Metro	0.4197		0.6050		0.9047	
Small Metro	0.1285		0.7109		0.8579	
250K – 1 Mil. Pop.	0.0889		0.7117		0.8958	
< 250K Pop.	0.6777		0.2820		0.6248	
Nonmetro	0.9340		0.9559		0.6531	
Urbanized	0.8117		0.3660		0.7443	
Less Urbanized	0.9009		0.5452		0.7910	
Completely Rural	0.3766		0.4250		0.8436	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.71C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	629	694	883	953	945	974
GEOGRAPHIC DIVISION						
Northeast	254	263	392	359	431	400
New England	97	126	167	168	183	200
Middle Atlantic	233	228	352	315	388	338
Midwest	237	242	363	339	376	371
East North Central	198	210	310	281	316	304
West North Central	128	126	190	209	207	237
South	451	461	598	647	634	648
South Atlantic	359	338	465	487	491	489
East South Central	162	212	212	280	201	263
West South Central	226	224	296	324	331	352
West	330	393	479	536	495	538
Mountain	126	168	192	252	202	239
Pacific	303	356	438	470	452	484
COUNTY TYPE						
Large Metro	1,824	1,815	1,613	1,606	1,390	1,381
Small Metro	1,307	1,424	1,069	1,209	929	1,011
250K – 1 Mil. Pop.	1,114	1,156	921	985	800	822
< 250K Pop.	856	836	692	701	579	595
Nonmetro	925	951	741	791	643	671
Urbanized	676	736	600	631	508	542
Less Urbanized	640	702	479	558	405	453
Completely Rural	399	329	264	244	233	200

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.71D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.28	0.30	0.39	0.42	0.42	0.42
GEOGRAPHIC DIVISION						
Northeast	0.60	0.62	0.93	0.85	1.02	0.95
New England	0.88	1.13	1.51	1.50	1.65	1.79
Middle Atlantic	0.75	0.73	1.14	1.01	1.25	1.09
Midwest	0.48	0.48	0.73	0.68	0.75	0.74
East North Central	0.57	0.60	0.90	0.81	0.91	0.87
West North Central	0.85	0.83	1.26	1.37	1.38	1.56
South	0.54	0.55	0.72	0.77	0.76	0.77
South Atlantic	0.81	0.76	1.06	1.09	1.11	1.10
East South Central	1.20	1.56	1.56	2.05	1.49	1.93
West South Central	0.89	0.87	1.16	1.25	1.30	1.36
West	0.63	0.74	0.91	1.01	0.95	1.02
Mountain	0.79	1.03	1.20	1.55	1.26	1.47
Pacific	0.83	0.97	1.21	1.28	1.24	1.32
COUNTY TYPE						
Large Metro	0.41	0.44	0.56	0.58	0.62	0.61
Small Metro	0.51	0.52	0.71	0.71	0.76	0.76
250K – 1 Mil. Pop.	0.56	0.57	0.89	0.82	0.97	0.90
< 250K Pop.	0.97	1.06	1.25	1.37	1.24	1.41
Nonmetro	0.66	0.73	0.97	1.04	0.97	1.03
Urbanized	1.08	1.04	1.57	1.71	1.59	1.68
Less Urbanized	1.01	1.10	1.38	1.47	1.35	1.39
Completely Rural	2.02	2.51	2.92	3.39	2.96	3.25

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.71P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.7868		0.5900		0.9436	
GEOGRAPHIC DIVISION						
Northeast	0.8853		0.6060		0.4256	
New England	0.5165		0.6022		0.5284	
Middle Atlantic	0.6125		0.7540		0.5670	
Midwest	0.3300		0.4207		0.9289	
East North Central	0.3810		0.1200		0.2710	
West North Central	0.6488		0.3717		0.1619	
South	0.9629		0.4844		0.9423	
South Atlantic	0.4688		0.6173		0.5160	
East South Central	0.9766		0.9538		0.7488	
West South Central	0.2260		0.4901		0.2441	
West	0.1930		0.7922		0.3557	
Mountain	0.1037		0.3793		0.2646	
Pacific	0.4656		0.9063		0.6273	
COUNTY TYPE						
Large Metro	0.4690		0.6485		0.8796	
Small Metro	0.1578		0.7766		0.8741	
250K – 1 Mil. Pop.	0.1553		0.6333		0.8060	
< 250K Pop.	0.5808		0.2900		0.5595	
Nonmetro	0.8551		0.7950		0.8543	
Urbanized	0.8383		0.2773		0.8167	
Less Urbanized	0.9696		0.6880		0.9828	
Completely Rural	0.3872		0.4107		0.8436	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.72C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	957	990	783	752	406	422
GEOGRAPHIC DIVISION						
Northeast	430	403	335	301	176	172
New England	187	204	157	148	94	82
Middle Atlantic	386	340	297	264	148	152
Midwest	379	380	308	334	180	176
East North Central	318	311	253	281	143	143
West North Central	209	239	175	182	113	104
South	638	658	493	452	259	264
South Atlantic	500	495	389	356	201	213
East South Central	203	266	166	152	102	88
West South Central	326	353	253	238	133	127
West	513	549	383	425	192	243
Mountain	212	246	183	206	86	122
Pacific	467	492	336	373	172	209
COUNTY TYPE						
Large Metro	1,404	1,389	776	738	316	349
Small Metro	947	1,023	537	533	251	294
250K – 1 Mil. Pop.	812	831	444	412	210	233
< 250K Pop.	591	604	334	335	146	178
Nonmetro	649	685	380	405	191	165
Urbanized	513	554	279	296	150	107
Less Urbanized	409	461	259	258	112	119
Completely Rural	237	203	107	104	48	60

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.72D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.38	0.39	0.31	0.30	0.16	0.17
GEOGRAPHIC DIVISION						
Northeast	0.93	0.87	0.72	0.65	0.38	0.37
New England	1.54	1.67	1.29	1.21	0.77	0.67
Middle Atlantic	1.13	0.99	0.87	0.77	0.43	0.44
Midwest	0.69	0.69	0.56	0.60	0.33	0.32
East North Central	0.83	0.81	0.66	0.73	0.37	0.37
West North Central	1.25	1.42	1.05	1.08	0.68	0.62
South	0.69	0.71	0.54	0.49	0.28	0.28
South Atlantic	1.03	1.01	0.80	0.73	0.41	0.43
East South Central	1.36	1.77	1.11	1.01	0.68	0.59
West South Central	1.14	1.22	0.89	0.83	0.47	0.44
West	0.88	0.93	0.66	0.72	0.33	0.41
Mountain	1.19	1.37	1.03	1.14	0.48	0.68
Pacific	1.16	1.21	0.83	0.91	0.43	0.51
COUNTY TYPE						
Large Metro	0.56	0.56	0.46	0.43	0.23	0.24
Small Metro	0.69	0.70	0.54	0.53	0.31	0.34
250K – 1 Mil. Pop.	0.88	0.83	0.65	0.63	0.38	0.41
< 250K Pop.	1.13	1.30	0.96	0.94	0.53	0.62
Nonmetro	0.89	0.95	0.71	0.75	0.42	0.36
Urbanized	1.49	1.54	1.11	1.15	0.74	0.54
Less Urbanized	1.24	1.28	1.02	0.97	0.54	0.52
Completely Rural	2.71	2.89	1.86	1.77	0.98	1.26

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.72P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.8783		0.1412		0.4869	
GEOGRAPHIC DIVISION						
Northeast	0.4221		0.9546		0.2107	
New England	0.4723		0.9902		0.5923	
Middle Atlantic	0.5978		0.9452		0.2534	
Midwest	0.9775		0.1274		0.2249	
East North Central	0.3807		0.3518		0.2999	
West North Central	0.1965		0.1678		0.5267	
South	1.0000		0.0727		0.5781	
South Atlantic	0.5211		0.3439		0.7305	
East South Central	0.7773		0.4448		0.2928	
West South Central	0.2083		0.1103		0.4015	
West	0.3799		0.6472		0.1997	
Mountain	0.2627		0.5949		0.4475	
Pacific	0.6670		0.7990		0.2928	
COUNTY TYPE						
Large Metro	0.8254		0.3992		0.7816	
Small Metro	0.8491		0.2826		0.5266	
250K – 1 Mil. Pop.	0.7634		0.8205		0.6616	
< 250K Pop.	0.4801		0.1521		0.6522	
Nonmetro	0.8764		0.5063		0.7933	
Urbanized	0.7724		0.8084		0.7767	
Less Urbanized	0.9265		0.3727		0.7024	
Completely Rural	0.7760		0.7024		0.5019	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.73C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	79	80	62	62	30	28
GEOGRAPHIC DIVISION						
Northeast	32	31	27	24	13	13
New England	15	14	12	12	7	5
Middle Atlantic	28	28	25	21	11	12
Midwest	29	30	23	23	12	9
East North Central	24	22	19	17	10	7
West North Central	17	21	13	16	7	6
South	46	53	35	38	16	14
South Atlantic	34	41	26	27	12	10
East South Central	15	16	12	13	7	4
West South Central	27	28	21	20	10	9
West	43	45	34	35	17	18
Mountain	19	19	14	18	7	9
Pacific	38	40	31	30	15	16
COUNTY TYPE						
Large Metro	69	73	53	54	24	23
Small Metro	54	46	39	32	16	13
250K – 1 Mil. Pop.	42	36	30	25	13	11
< 250K Pop.	34	28	25	19	9	6
Nonmetro	31	36	23	24	11	11
Urbanized	20	23	14	15	7	7
Less Urbanized	22	27	16	18	7	7
Completely Rural	9	9	7	8	3	2

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.73D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.32	0.33	0.25	0.25	0.12	0.11
GEOGRAPHIC DIVISION						
Northeast	0.74	0.73	0.64	0.58	0.31	0.30
New England	1.34	1.26	1.12	1.11	0.66	0.45
Middle Atlantic	0.89	0.88	0.77	0.68	0.35	0.38
Midwest	0.54	0.56	0.43	0.42	0.22	0.18
East North Central	0.62	0.58	0.51	0.44	0.26	0.19
West North Central	1.06	1.32	0.79	1.00	0.43	0.40
South	0.52	0.59	0.39	0.42	0.18	0.16
South Atlantic	0.74	0.91	0.57	0.61	0.26	0.23
East South Central	1.02	1.08	0.80	0.87	0.48	0.30
West South Central	0.91	0.94	0.70	0.67	0.33	0.30
West	0.72	0.77	0.58	0.60	0.28	0.31
Mountain	1.07	1.09	0.79	0.99	0.40	0.50
Pacific	0.93	1.00	0.76	0.74	0.37	0.39
COUNTY TYPE						
Large Metro	0.46	0.50	0.38	0.39	0.18	0.18
Small Metro	0.55	0.53	0.43	0.39	0.20	0.17
250K – 1 Mil. Pop.	0.65	0.62	0.52	0.45	0.24	0.22
< 250K Pop.	1.01	0.98	0.78	0.66	0.36	0.24
Nonmetro	0.68	0.79	0.55	0.56	0.27	0.26
Urbanized	1.00	0.96	0.72	0.74	0.41	0.43
Less Urbanized	1.01	1.28	0.83	0.84	0.39	0.36
Completely Rural	1.76	2.60	1.50	2.29	0.74	0.75

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.73P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.0088		0.0032		0.0106	
GEOGRAPHIC DIVISION						
Northeast	0.3564		0.1678		0.2077	
New England	0.5759		0.9248		0.0141	
Middle Atlantic	0.1945		0.1048		0.8682	
Midwest	0.0141		0.0091		0.0417	
East North Central	0.0002		0.0000		0.0160	
West North Central	0.6072		0.3475		0.7529	
South	0.0958		0.1505		0.0475	
South Atlantic	0.4672		0.8572		0.5332	
East South Central	0.4078		0.3388		0.0905	
West South Central	0.1232		0.0809		0.1131	
West	0.6129		0.3050		0.6999	
Mountain	0.3313		0.6078		0.5061	
Pacific	0.9140		0.3762		0.9232	
COUNTY TYPE						
Large Metro	0.0788		0.0410		0.1042	
Small Metro	0.0521		0.0140		0.0257	
250K – 1 Mil. Pop.	0.4429		0.1274		0.3827	
< 250K Pop.	0.0327		0.0366		0.0054	
Nonmetro	0.4717		0.8071		0.5540	
Urbanized	0.3765		0.7651		0.3748	
Less Urbanized	0.6413		0.5573		0.9055	
Completely Rural	0.7477		0.2818		0.8168	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.74C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	174	169	182	168	128	127
GEOGRAPHIC DIVISION						
Northeast	69	68	72	67	53	47
New England	39	36	39	33	26	26
Middle Atlantic	57	58	61	57	46	39
Midwest	71	64	71	74	46	49
East North Central	58	50	57	61	39	40
West North Central	41	41	40	40	24	28
South	113	111	113	114	89	90
South Atlantic	90	86	90	92	75	79
East South Central	35	40	35	40	22	25
West South Central	55	55	57	57	40	37
West	96	93	96	87	53	64
Mountain	43	45	41	41	23	24
Pacific	86	80	86	76	47	59
COUNTY TYPE						
Large Metro	246	275	192	195	93	104
Small Metro	207	243	163	185	92	107
250K – 1 Mil. Pop.	155	179	119	128	58	66
< 250K Pop.	141	164	114	132	69	86
Nonmetro	214	121	159	94	81	48
Urbanized	197	92	142	72	73	34
Less Urbanized	92	85	75	67	34	35
Completely Rural	26	28	22	20	10	11

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.74D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.52	0.50	0.54	0.49	0.38	0.37
GEOGRAPHIC DIVISION						
Northeast	1.13	1.13	1.17	1.10	0.87	0.78
New England	2.45	2.22	2.45	2.06	1.65	1.61
Middle Atlantic	1.26	1.31	1.34	1.30	1.01	0.88
Midwest	0.97	0.85	0.96	0.98	0.63	0.65
East North Central	1.13	0.98	1.12	1.19	0.76	0.78
West North Central	1.80	1.72	1.77	1.68	1.08	1.20
South	0.94	0.89	0.94	0.91	0.74	0.73
South Atlantic	1.48	1.36	1.47	1.46	1.22	1.26
East South Central	1.86	2.04	1.82	2.02	1.14	1.28
West South Central	1.37	1.33	1.43	1.38	0.99	0.90
West	1.18	1.14	1.18	1.08	0.65	0.79
Mountain	1.76	1.80	1.68	1.66	0.96	0.98
Pacific	1.51	1.42	1.52	1.36	0.83	1.05
COUNTY TYPE						
Large Metro	0.73	0.72	0.75	0.67	0.48	0.49
Small Metro	0.88	0.87	0.91	0.96	0.70	0.77
250K – 1 Mil. Pop.	1.07	1.03	1.07	1.05	0.70	0.77
< 250K Pop.	1.51	1.60	1.68	1.90	1.43	1.68
Nonmetro	1.47	1.18	1.31	1.15	1.02	0.76
Urbanized	2.40	1.68	2.07	1.77	1.73	1.14
Less Urbanized	1.63	1.76	1.69	1.68	1.17	1.12
Completely Rural	3.41	4.36	3.49	3.60	2.73	3.07

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.74P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.5434		0.0542		0.6412	
GEOGRAPHIC DIVISION						
Northeast	0.3758		0.3619		0.6995	
New England	0.6165		0.9885		0.7019	
Middle Atlantic	0.4368		0.2710		0.8463	
Midwest	0.7153		0.3301		0.6649	
East North Central	0.6504		0.0548		0.2380	
West North Central	0.1812		0.3059		0.3039	
South	0.4352		0.1350		0.6918	
South Atlantic	0.9891		0.2091		0.8925	
East South Central	0.5615		0.2542		0.4361	
West South Central	0.2894		0.8495		0.9421	
West	0.7543		0.6783		0.6851	
Mountain	0.3065		0.9786		0.2113	
Pacific	0.8921		0.6680		0.9583	
COUNTY TYPE						
Large Metro	0.8300		0.4014		0.6797	
Small Metro	0.8013		0.0341		0.1531	
250K – 1 Mil. Pop.	0.5959		0.1871		0.5241	
< 250K Pop.	0.2778		0.0790		0.1470	
Nonmetro	0.3092		0.6529		0.9965	
Urbanized	0.9125		0.5518		0.7346	
Less Urbanized	0.1510		0.5316		0.9675	
Completely Rural	0.6493		0.0606		0.4631	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.75C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	931	946	751	715	377	398
GEOGRAPHIC DIVISION						
Northeast	422	394	308	294	166	161
New England	174	196	139	144	87	73
Middle Atlantic	381	334	276	258	140	144
Midwest	366	357	289	317	176	162
East North Central	307	293	235	268	138	133
West North Central	204	231	169	171	111	92
South	611	621	468	420	238	242
South Atlantic	467	466	362	325	180	191
East South Central	197	256	160	143	99	84
West South Central	321	344	246	228	127	123
West	483	516	376	399	182	229
Mountain	195	228	173	186	83	116
Pacific	441	466	334	354	163	198
COUNTY TYPE						
Large Metro	1,333	1,317	716	686	295	321
Small Metro	867	932	485	459	231	265
250K – 1 Mil. Pop.	748	771	404	367	199	218
< 250K Pop.	528	540	295	284	131	152
Nonmetro	575	631	320	372	163	156
Urbanized	438	501	213	258	118	98
Less Urbanized	371	422	231	235	103	110
Completely Rural	218	188	95	94	46	57

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.75D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.48	0.48	0.39	0.37	0.19	0.20
GEOGRAPHIC DIVISION						
Northeast	1.17	1.09	0.86	0.81	0.46	0.44
New England	1.83	2.06	1.47	1.51	0.92	0.76
Middle Atlantic	1.44	1.25	1.04	0.96	0.53	0.54
Midwest	0.86	0.84	0.68	0.75	0.41	0.38
East North Central	1.04	0.99	0.79	0.90	0.47	0.45
West North Central	1.59	1.80	1.31	1.33	0.87	0.72
South	0.86	0.87	0.66	0.59	0.34	0.34
South Atlantic	1.23	1.22	0.96	0.85	0.47	0.50
East South Central	1.69	2.20	1.38	1.22	0.85	0.72
West South Central	1.49	1.58	1.15	1.05	0.59	0.57
West	1.09	1.15	0.85	0.89	0.41	0.51
Mountain	1.44	1.66	1.28	1.35	0.61	0.84
Pacific	1.44	1.50	1.09	1.14	0.53	0.64
COUNTY TYPE						
Large Metro	0.70	0.70	0.57	0.54	0.28	0.29
Small Metro	0.88	0.87	0.69	0.63	0.38	0.41
250K – 1 Mil. Pop.	1.13	1.03	0.84	0.76	0.48	0.50
< 250K Pop.	1.42	1.66	1.21	1.14	0.64	0.71
Nonmetro	1.07	1.16	0.82	0.91	0.48	0.45
Urbanized	1.75	1.87	1.20	1.39	0.82	0.67
Less Urbanized	1.52	1.58	1.23	1.20	0.65	0.65
Completely Rural	3.16	3.54	2.16	2.04	1.17	1.50

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.75P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.9800		0.3485		0.6433	
GEOGRAPHIC DIVISION						
Northeast	0.3318		0.6974		0.1916	
New England	0.4589		0.9454		0.5678	
Middle Atlantic	0.4712		0.6405		0.2378	
Midwest	0.9876		0.2052		0.2704	
East North Central	0.2250		0.7548		0.5068	
West North Central	0.1044		0.0645		0.3581	
South	0.8504		0.1291		0.6787	
South Atlantic	0.5064		0.4364		0.6964	
East South Central	0.6820		0.6164		0.4025	
West South Central	0.3063		0.1356		0.4495	
West	0.3192		0.5132		0.1378	
Mountain	0.1735		0.5647		0.2880	
Pacific	0.6480		0.6452		0.2678	
COUNTY TYPE						
Large Metro	0.9047		0.5757		0.7611	
Small Metro	0.8579		0.4953		0.7678	
250K – 1 Mil. Pop.	0.8958		0.9687		0.7479	
< 250K Pop.	0.6248		0.2592		0.9701	
Nonmetro	0.6531		0.6599		0.8956	
Urbanized	0.7443		0.8345		0.8096	
Less Urbanized	0.7910		0.5218		0.7738	
Completely Rural	0.8436		0.6678		0.4548	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.76C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	945	974	784	748	404	421
GEOGRAPHIC DIVISION						
Northeast	431	400	333	302	175	171
New England	183	200	154	147	92	82
Middle Atlantic	388	338	296	265	148	151
Midwest	376	371	307	332	180	175
East North Central	316	304	251	278	142	142
West North Central	207	237	176	181	114	103
South	634	648	494	453	260	265
South Atlantic	491	489	390	358	202	214
East South Central	201	263	165	150	101	88
West South Central	331	352	254	237	132	127
West	495	538	384	418	192	241
Mountain	202	239	181	204	86	122
Pacific	452	484	339	366	172	208
COUNTY TYPE						
Large Metro	1,390	1,381	771	733	313	347
Small Metro	929	1,011	527	527	249	294
250K – 1 Mil. Pop.	800	822	438	407	208	233
< 250K Pop.	579	595	328	330	145	178
Nonmetro	643	671	376	399	190	164
Urbanized	508	542	276	290	149	106
Less Urbanized	405	453	256	254	110	118
Completely Rural	233	200	105	102	47	59

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.76D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.42	0.42	0.34	0.33	0.18	0.18
GEOGRAPHIC DIVISION						
Northeast	1.02	0.95	0.79	0.71	0.42	0.40
New England	1.65	1.79	1.39	1.32	0.83	0.74
Middle Atlantic	1.25	1.09	0.95	0.85	0.48	0.49
Midwest	0.75	0.74	0.62	0.66	0.36	0.35
East North Central	0.91	0.87	0.72	0.80	0.41	0.41
West North Central	1.38	1.56	1.17	1.19	0.75	0.68
South	0.76	0.77	0.59	0.54	0.31	0.31
South Atlantic	1.11	1.10	0.89	0.80	0.46	0.48
East South Central	1.49	1.93	1.22	1.10	0.75	0.64
West South Central	1.30	1.36	1.00	0.92	0.52	0.49
West	0.95	1.02	0.73	0.79	0.37	0.46
Mountain	1.26	1.47	1.13	1.25	0.54	0.75
Pacific	1.24	1.32	0.93	1.00	0.47	0.57
COUNTY TYPE						
Large Metro	0.62	0.61	0.51	0.48	0.25	0.26
Small Metro	0.76	0.76	0.60	0.58	0.34	0.38
250K – 1 Mil. Pop.	0.97	0.90	0.72	0.70	0.42	0.45
< 250K Pop.	1.24	1.41	1.06	1.04	0.59	0.68
Nonmetro	0.97	1.03	0.78	0.82	0.47	0.40
Urbanized	1.59	1.68	1.21	1.27	0.82	0.59
Less Urbanized	1.35	1.39	1.12	1.07	0.59	0.58
Completely Rural	2.96	3.25	2.04	1.90	1.09	1.36

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.76P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.9436		0.1921		0.5806	
GEOGRAPHIC DIVISION						
Northeast	0.4256		0.8940		0.2383	
New England	0.5284		0.9661		0.6994	
Middle Atlantic	0.5670		0.8662		0.2492	
Midwest	0.9289		0.1694		0.2662	
East North Central	0.2710		0.4970		0.3618	
West North Central	0.1619		0.1365		0.5314	
South	0.9423		0.0862		0.6476	
South Atlantic	0.5160		0.3376		0.7191	
East South Central	0.7488		0.4761		0.3300	
West South Central	0.2441		0.1405		0.4589	
West	0.3557		0.6072		0.1932	
Mountain	0.2646		0.5732		0.4285	
Pacific	0.6273		0.7608		0.2922	
COUNTY TYPE						
Large Metro	0.8796		0.4767		0.8627	
Small Metro	0.8741		0.3399		0.5860	
250K – 1 Mil. Pop.	0.8060		0.8670		0.6738	
< 250K Pop.	0.5595		0.1935		0.7483	
Nonmetro	0.8543		0.5048		0.8129	
Urbanized	0.8167		0.7731		0.7857	
Less Urbanized	0.9828		0.4192		0.7293	
Completely Rural	0.8436		0.8080		0.5211	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.