

Table 2.77C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	300	291	272	265	212	204
GENDER						
Male	202	198	186	180	150	142
Female	175	169	157	155	125	117
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	265	251	243	230	193	182
White	228	217	213	201	171	159
Black or African American	102	97	89	82	63	61
American Indian or Alaska Native	14	14	11	11	7	9
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	49	49	43	44	28	31
Two or More Races	31	30	25	26	22	18
Hispanic or Latino	140	137	120	124	82	85
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	147	145	139	135	117	108
Female, White, Not Hispanic	134	129	124	120	96	96
Male, Black, Not Hispanic	62	65	56	55	42	43
Female, Black, Not Hispanic	69	59	59	52	44	35
Male, Hispanic	102	93	90	85	61	63
Female, Hispanic	78	76	68	68	49	47

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.77D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.44	0.44	0.44	0.43	0.43	0.41
GENDER						
Male	0.57	0.60	0.58	0.60	0.58	0.55
Female	0.60	0.58	0.58	0.58	0.55	0.52
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.50	0.48	0.49	0.47	0.48	0.47
White	0.57	0.54	0.57	0.54	0.57	0.54
Black or African American	1.08	1.00	1.04	0.96	0.91	0.89
American Indian or Alaska Native	4.11	3.58	4.19	3.52	3.12	3.66
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.02	2.36	1.98	2.25	1.54	1.82
Two or More Races	2.68	2.33	2.72	2.35	2.51	2.03
Hispanic or Latino	0.97	1.01	0.96	1.00	0.81	0.84
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.70	0.75	0.72	0.75	0.74	0.71
Female, White, Not Hispanic	0.75	0.73	0.74	0.73	0.71	0.71
Male, Black, Not Hispanic	1.41	1.46	1.36	1.42	1.26	1.31
Female, Black, Not Hispanic	1.57	1.34	1.48	1.31	1.27	1.08
Male, Hispanic	1.33	1.33	1.35	1.36	1.17	1.22
Female, Hispanic	1.35	1.34	1.34	1.34	1.14	1.04

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.77P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.0006		0.0282		0.1032	
GENDER						
Male	0.0025		0.0984		0.8062	
Female	0.0438		0.1148		0.0246	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0026		0.0310		0.1414	
White	0.0020		0.0051		0.1384	
Black or African American	0.5830		0.6131		0.9793	
American Indian or Alaska Native	0.8651		0.8176		0.8700	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.5129		0.7518		0.7845	
Two or More Races	0.9820		0.4758		0.3015	
Hispanic or Latino	0.0985		0.5157		0.5194	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.0423		0.0923		0.6881	
Female, White, Not Hispanic	0.0085		0.0138		0.0629	
Male, Black, Not Hispanic	0.4886		0.9809		0.9812	
Female, Black, Not Hispanic	0.9356		0.4390		0.9720	
Male, Hispanic	0.0135		0.2265		0.8079	
Female, Hispanic	0.9276		0.7659		0.1683	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.78C Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	2,166	2,205	1,872	1,949	1,645	1,675
GENDER						
Male	1,358	1,415	1,189	1,239	1,095	1,134
Female	1,290	1,270	1,132	1,111	982	906
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2,043	2,099	1,759	1,858	1,540	1,601
White	1,822	1,899	1,591	1,691	1,400	1,461
Black or African American	715	708	554	582	478	492
American Indian or Alaska Native	111	*	76	57	58	50
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	426	421	352	378	300	328
Two or More Races	144	175	119	147	108	117
Hispanic or Latino	711	707	597	603	498	484
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	1,129	1,176	971	1,034	895	953
Female, White, Not Hispanic	1,105	1,121	996	997	864	816
Male, Black, Not Hispanic	435	423	369	358	329	325
Female, Black, Not Hispanic	450	475	362	384	298	317
Male, Hispanic	489	509	436	454	385	381
Female, Hispanic	413	409	324	322	256	255

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.78D Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.30	0.32	0.42	0.44	0.44	0.45
GENDER						
Male	0.36	0.35	0.55	0.59	0.62	0.63
Female	0.45	0.48	0.59	0.58	0.60	0.56
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.30	0.34	0.44	0.46	0.46	0.47
White	0.28	0.32	0.47	0.50	0.50	0.52
Black or African American	1.20	1.08	1.39	1.37	1.37	1.34
American Indian or Alaska Native	3.73	*	5.28	5.06	5.05	4.51
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.45	2.46	2.39	2.68	2.17	2.50
Two or More Races	1.99	2.78	3.25	3.65	3.49	3.75
Hispanic or Latino	0.95	0.96	1.25	1.28	1.19	1.22
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.32	0.35	0.64	0.68	0.70	0.73
Female, White, Not Hispanic	0.44	0.48	0.65	0.66	0.71	0.67
Male, Black, Not Hispanic	1.29	1.39	1.99	1.86	2.07	1.94
Female, Black, Not Hispanic	1.74	1.39	1.81	1.81	1.63	1.74
Male, Hispanic	1.13	1.10	1.59	1.75	1.70	1.76
Female, Hispanic	1.48	1.47	1.63	1.66	1.50	1.51

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.78P Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.8731		0.6394		0.9945	
GENDER						
Male	0.9779		0.1136		0.7266	
Female	0.8066		0.4390		0.7569	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.6239		0.6455		0.9242	
White	0.4032		0.6914		0.9415	
Black or African American	0.4305		0.9674		0.9402	
American Indian or Alaska Native	*		0.7678		0.8628	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.9512		0.7583		0.7986	
Two or More Races	0.2631		0.7125		0.6396	
Hispanic or Latino	0.4657		0.9277		0.8035	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.4811		0.3744		0.8075	
Female, White, Not Hispanic	0.5736		0.7692		0.8845	
Male, Black, Not Hispanic	0.5068		0.3312		0.4832	
Female, Black, Not Hispanic	0.1441		0.4672		0.6059	
Male, Hispanic	0.4008		0.4401		0.8180	
Female, Hispanic	0.7486		0.5035		0.4807	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.79C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Demographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	212	204	174	162	91	84
GENDER						
Male	150	142	127	119	76	67
Female	125	117	93	90	39	44
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	193	182	157	146	84	79
White	171	159	143	130	82	74
Black or African American	63	61	45	43	15	16
American Indian or Alaska Native	7	9	6	8	3	6
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	1
Asian	28	31	22	20	7	7
Two or More Races	22	18	17	15	10	10
Hispanic or Latino	82	85	68	68	29	30
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	117	108	104	93	68	57
Female, White, Not Hispanic	96	96	72	75	34	41
Male, Black, Not Hispanic	42	43	28	31	14	14
Female, Black, Not Hispanic	44	35	31	24	7	7
Male, Hispanic	61	63	54	54	25	27
Female, Hispanic	49	47	38	37	16	13

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.79D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Demographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.43	0.41	0.38	0.35	0.22	0.21
GENDER						
Male	0.58	0.55	0.53	0.51	0.36	0.32
Female	0.55	0.52	0.45	0.43	0.21	0.23
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.48	0.47	0.42	0.40	0.26	0.24
White	0.57	0.54	0.51	0.48	0.34	0.31
Black or African American	0.91	0.89	0.70	0.68	0.27	0.28
American Indian or Alaska Native	3.12	3.66	2.58	3.46	1.45	2.74
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	1.37
Asian	1.54	1.82	1.27	1.25	0.42	0.42
Two or More Races	2.51	2.03	2.10	1.70	1.32	1.30
Hispanic or Latino	0.81	0.84	0.76	0.72	0.37	0.38
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.74	0.71	0.71	0.67	0.53	0.46
Female, White, Not Hispanic	0.71	0.71	0.57	0.60	0.30	0.36
Male, Black, Not Hispanic	1.26	1.31	0.91	1.00	0.47	0.50
Female, Black, Not Hispanic	1.27	1.08	0.97	0.80	0.24	0.24
Male, Hispanic	1.17	1.22	1.11	1.12	0.62	0.65
Female, Hispanic	1.14	1.04	0.98	0.88	0.44	0.34

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.79P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Demographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.1032		0.0286		0.4051	
GENDER						
Male	0.8062		0.3267		0.5435	
Female	0.0246		0.0142		0.4766	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1414		0.0594		0.3708	
White	0.1384		0.0854		0.4046	
Black or African American	0.9793		0.8311		0.8929	
American Indian or Alaska Native	0.8700		0.7232		0.7095	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.7845		0.4468		1.0000	
Two or More Races	0.3015		0.2333		0.5865	
Hispanic or Latino	0.5194		0.2835		0.8517	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.6881		0.2897		0.3334	
Female, White, Not Hispanic	0.0629		0.1037		0.8423	
Male, Black, Not Hispanic	0.9812		0.6664		0.9093	
Female, Black, Not Hispanic	0.9720		0.4236		0.9749	
Male, Hispanic	0.8079		0.5872		0.5122	
Female, Hispanic	0.1683		0.2262		0.3847	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.80C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Demographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	1,645	1,675	920	910	399	445
GENDER						
Male	1,095	1,134	711	708	347	375
Female	982	906	481	470	189	203
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	1,540	1,601	827	843	374	421
White	1,400	1,461	754	752	356	374
Black or African American	478	492	296	285	123	138
American Indian or Alaska Native	58	50	35	40	21	17
Native Hawaiian or Other Pacific Islander	*	*	*	*	10	*
Asian	300	328	114	147	30	72
Two or More Races	108	117	70	62	29	31
Hispanic or Latino	498	484	348	369	138	137
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	895	953	575	587	300	318
Female, White, Not Hispanic	864	816	407	383	176	163
Male, Black, Not Hispanic	329	325	225	210	109	114
Female, Black, Not Hispanic	298	317	168	171	50	77
Male, Hispanic	385	381	307	311	133	117
Female, Hispanic	256	255	158	174	39	69

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.80D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Demographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.44	0.45	0.36	0.34	0.18	0.19
GENDER						
Male	0.62	0.63	0.58	0.54	0.33	0.33
Female	0.60	0.56	0.39	0.39	0.17	0.18
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.46	0.47	0.38	0.36	0.20	0.21
White	0.50	0.52	0.43	0.40	0.23	0.23
Black or African American	1.37	1.34	1.06	1.00	0.50	0.54
American Indian or Alaska Native	5.05	4.51	3.18	3.73	2.13	1.75
Native Hawaiian or Other Pacific Islander	*	*	*	*	1.58	*
Asian	2.17	2.50	1.15	1.35	0.32	0.69
Two or More Races	3.49	3.75	2.76	2.61	1.31	1.29
Hispanic or Latino	1.19	1.22	1.00	1.05	0.48	0.46
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.70	0.73	0.67	0.66	0.40	0.42
Female, White, Not Hispanic	0.71	0.67	0.49	0.46	0.23	0.21
Male, Black, Not Hispanic	2.07	1.94	1.85	1.70	1.00	1.00
Female, Black, Not Hispanic	1.63	1.74	1.14	1.13	0.37	0.56
Male, Hispanic	1.70	1.76	1.68	1.67	0.89	0.77
Female, Hispanic	1.50	1.51	1.04	1.13	0.28	0.49

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.80P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Demographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.9945		0.3157		0.6656	
GENDER						
Male	0.7266		0.3606		0.6915	
Female	0.7569		0.5679		0.8041	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.9242		0.2028		0.7149	
White	0.9415		0.1441		0.5630	
Black or African American	0.9402		0.9538		0.9358	
American Indian or Alaska Native	0.8628		0.5841		0.4965	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.7986		0.3619		0.1551	
Two or More Races	0.6396		0.5209		0.8157	
Hispanic or Latino	0.8035		0.7227		0.8058	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.8075		0.5183		0.9229	
Female, White, Not Hispanic	0.8845		0.1029		0.1692	
Male, Black, Not Hispanic	0.4832		0.4193		0.5111	
Female, Black, Not Hispanic	0.6059		0.3921		0.3844	
Male, Hispanic	0.8180		0.8049		0.2579	
Female, Hispanic	0.4807		0.2522		0.0649	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.81C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	302	291	273	266	212	204
GEOGRAPHIC DIVISION						
Northeast	131	112	124	106	98	87
New England	60	49	60	46	49	35
Middle Atlantic	116	100	109	94	85	79
Midwest	113	116	106	107	85	81
East North Central	100	98	94	90	76	65
West North Central	53	62	49	57	39	47
South	178	201	166	181	132	141
South Atlantic	138	156	128	142	99	104
East South Central	61	66	55	57	41	44
West South Central	103	107	94	97	73	84
West	168	146	149	134	107	97
Mountain	63	77	55	64	35	48
Pacific	154	124	136	117	99	85
COUNTY TYPE						
Large Metro	218	225	200	206	146	154
Small Metro	188	186	167	166	130	128
250K – 1 Mil. Pop.	143	137	124	121	93	91
< 250K Pop.	127	126	115	115	92	90
Nonmetro	137	122	126	111	96	85
Urbanized	108	79	100	73	74	53
Less Urbanized	90	99	80	90	61	69
Completely Rural	32	31	27	26	19	19

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.81D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.45	0.44	0.44	0.44	0.43	0.41
GEOGRAPHIC DIVISION						
Northeast	0.97	0.96	1.02	0.97	1.03	0.97
New England	1.78	1.77	2.05	1.89	2.07	1.72
Middle Atlantic	1.14	1.13	1.18	1.12	1.19	1.15
Midwest	0.78	0.78	0.80	0.80	0.78	0.75
East North Central	0.94	0.91	0.97	0.92	0.96	0.82
West North Central	1.41	1.49	1.40	1.58	1.31	1.57
South	0.76	0.77	0.74	0.77	0.72	0.77
South Atlantic	1.20	1.15	1.17	1.17	1.08	1.11
East South Central	1.55	1.66	1.48	1.62	1.44	1.57
West South Central	1.24	1.34	1.20	1.35	1.16	1.38
West	0.99	1.01	1.04	1.01	0.89	0.83
Mountain	1.55	1.41	1.58	1.39	1.26	1.33
Pacific	1.23	1.31	1.31	1.31	1.14	1.04
COUNTY TYPE						
Large Metro	0.62	0.64	0.62	0.64	0.58	0.59
Small Metro	0.75	0.76	0.76	0.76	0.73	0.75
250K – 1 Mil. Pop.	0.91	0.91	0.87	0.91	0.82	0.85
< 250K Pop.	1.27	1.32	1.44	1.36	1.42	1.45
Nonmetro	1.06	1.05	1.11	1.08	1.10	1.03
Urbanized	1.74	1.55	1.87	1.68	1.83	1.44
Less Urbanized	1.43	1.51	1.47	1.59	1.52	1.65
Completely Rural	2.44	3.88	2.31	3.37	2.30	2.75

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.81P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.0003		0.0168		0.0687	
GEOGRAPHIC DIVISION						
Northeast	0.2079		0.3163		0.2020	
New England	0.4053		0.1853		0.3096	
Middle Atlantic	0.3203		0.7110		0.3597	
Midwest	0.0046		0.0722		0.1995	
East North Central	0.0011		0.0184		0.0337	
West North Central	0.7422		0.8371		0.4557	
South	0.2043		0.6913		0.2695	
South Atlantic	0.9195		0.8679		0.8284	
East South Central	0.3622		0.5902		0.4420	
West South Central	0.1022		0.5356		0.2095	
West	0.0330		0.0861		0.9125	
Mountain	0.7298		0.6193		0.9912	
Pacific	0.0264		0.0922		0.8983	
COUNTY TYPE						
Large Metro	0.0010		0.0750		0.3085	
Small Metro	0.4048		0.3236		0.1291	
250K – 1 Mil. Pop.	0.8633		0.6805		0.6153	
< 250K Pop.	0.2313		0.2924		0.0762	
Nonmetro	0.0866		0.1910		0.4795	
Urbanized	0.0236		0.0807		0.7862	
Less Urbanized	0.7693		0.8221		0.6037	
Completely Rural	0.9755		0.6388		0.7062	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.82C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	2,165	2,217	1,875	1,960	1,647	1,681
GEOGRAPHIC DIVISION						
Northeast	910	844	786	744	703	681
New England	404	452	355	419	328	415
Middle Atlantic	817	710	702	613	625	539
Midwest	763	849	686	748	618	671
East North Central	585	683	540	602	488	532
West North Central	480	486	423	439	376	397
South	1,382	1,462	1,195	1,307	1,045	1,107
South Atlantic	1,022	1,118	878	1,014	759	856
East South Central	471	479	358	403	297	342
West South Central	730	836	656	741	596	630
West	1,159	1,178	1,030	1,041	894	883
Mountain	486	603	408	518	364	417
Pacific	1,053	999	945	894	814	772
COUNTY TYPE						
Large Metro	1,801	1,786	1,589	1,578	1,372	1,368
Small Metro	1,272	1,381	1,042	1,174	909	985
250K – 1 Mil. Pop.	1,084	1,132	897	970	784	811
< 250K Pop.	828	803	672	669	561	571
Nonmetro	891	931	705	779	619	663
Urbanized	642	723	565	624	483	537
Less Urbanized	617	677	458	537	391	441
Completely Rural	389	323	255	240	227	196

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.82D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.29	0.32	0.41	0.44	0.44	0.45
GEOGRAPHIC DIVISION						
Northeast	0.64	0.66	0.99	0.90	1.09	1.00
New England	0.90	1.21	1.57	1.60	1.72	1.88
Middle Atlantic	0.80	0.77	1.21	1.07	1.33	1.15
Midwest	0.49	0.50	0.76	0.72	0.79	0.78
East North Central	0.59	0.63	0.94	0.85	0.95	0.91
West North Central	0.88	0.86	1.32	1.46	1.45	1.66
South	0.57	0.58	0.75	0.81	0.80	0.81
South Atlantic	0.85	0.80	1.10	1.14	1.16	1.15
East South Central	1.26	1.66	1.65	2.19	1.58	2.04
West South Central	0.93	0.91	1.23	1.32	1.36	1.43
West	0.66	0.79	0.97	1.07	1.00	1.07
Mountain	0.79	1.12	1.25	1.64	1.32	1.54
Pacific	0.88	1.03	1.28	1.35	1.31	1.39
COUNTY TYPE						
Large Metro	0.43	0.46	0.59	0.61	0.65	0.65
Small Metro	0.54	0.56	0.76	0.76	0.81	0.81
250K – 1 Mil. Pop.	0.59	0.62	0.94	0.88	1.03	0.95
< 250K Pop.	1.03	1.12	1.33	1.47	1.30	1.51
Nonmetro	0.70	0.77	1.02	1.10	1.02	1.09
Urbanized	1.13	1.09	1.63	1.82	1.66	1.78
Less Urbanized	1.07	1.17	1.45	1.55	1.42	1.47
Completely Rural	2.14	2.59	3.03	3.55	3.08	3.37

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.82P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.8764		0.6512		0.9840	
GEOGRAPHIC DIVISION						
Northeast	0.8189		0.6733		0.3661	
New England	0.7247		0.8572		0.3656	
Middle Atlantic	0.6532		0.7029		0.5766	
Midwest	0.2749		0.4445		0.9396	
East North Central	0.3682		0.1216		0.2509	
West North Central	0.5227		0.3569		0.1444	
South	0.8494		0.4984		0.8493	
South Atlantic	0.5376		0.6554		0.4804	
East South Central	1.0000		0.9804		0.7223	
West South Central	0.1751		0.4912		0.3002	
West	0.1586		0.8379		0.3164	
Mountain	0.0646		0.3354		0.2125	
Pacific	0.4486		0.8272		0.6065	
COUNTY TYPE						
Large Metro	0.4251		0.6418		0.8973	
Small Metro	0.1923		0.8865		0.9881	
250K – 1 Mil. Pop.	0.1489		0.5758		0.7768	
< 250K Pop.	0.7209		0.3428		0.6819	
Nonmetro	0.8545		0.8361		0.8556	
Urbanized	0.8398		0.2992		0.8706	
Less Urbanized	0.9869		0.7060		0.9787	
Completely Rural	0.3876		0.3528		0.7620	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.83C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	212	204	174	161	91	84
GEOGRAPHIC DIVISION						
Northeast	98	87	77	71	33	36
New England	49	35	40	29	18	16
Middle Atlantic	85	79	65	64	28	32
Midwest	85	81	72	62	38	33
East North Central	76	65	65	49	34	25
West North Central	39	47	32	38	15	21
South	132	141	109	113	65	59
South Atlantic	99	104	82	83	53	47
East South Central	41	44	34	39	18	23
West South Central	73	84	61	68	33	31
West	107	97	86	77	40	37
Mountain	35	48	29	37	14	15
Pacific	99	85	80	67	37	34
COUNTY TYPE						
Large Metro	146	154	116	118	58	58
Small Metro	130	128	105	101	57	54
250K – 1 Mil. Pop.	93	91	72	69	33	35
< 250K Pop.	92	90	75	74	45	41
Nonmetro	96	85	83	68	44	35
Urbanized	74	53	64	42	35	20
Less Urbanized	61	69	52	55	26	28
Completely Rural	19	19	16	16	7	6

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.83D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.43	0.41	0.38	0.35	0.22	0.21
GEOGRAPHIC DIVISION						
Northeast	1.03	0.97	0.89	0.87	0.45	0.49
New England	2.07	1.72	1.80	1.53	0.94	0.87
Middle Atlantic	1.19	1.15	1.02	1.04	0.51	0.59
Midwest	0.78	0.75	0.71	0.64	0.41	0.37
East North Central	0.96	0.82	0.88	0.70	0.52	0.40
West North Central	1.31	1.57	1.09	1.34	0.58	0.78
South	0.72	0.77	0.65	0.67	0.44	0.39
South Atlantic	1.08	1.11	0.98	0.95	0.71	0.62
East South Central	1.44	1.57	1.29	1.47	0.73	0.96
West South Central	1.16	1.38	1.10	1.23	0.68	0.61
West	0.89	0.83	0.78	0.70	0.40	0.39
Mountain	1.26	1.33	1.04	1.12	0.46	0.51
Pacific	1.14	1.04	1.02	0.88	0.54	0.52
COUNTY TYPE						
Large Metro	0.58	0.59	0.49	0.49	0.28	0.28
Small Metro	0.73	0.75	0.68	0.66	0.44	0.41
250K – 1 Mil. Pop.	0.82	0.85	0.74	0.72	0.41	0.42
< 250K Pop.	1.42	1.45	1.32	1.33	0.95	0.90
Nonmetro	1.10	1.03	1.02	0.91	0.63	0.52
Urbanized	1.83	1.44	1.71	1.37	1.08	0.70
Less Urbanized	1.52	1.65	1.39	1.40	0.84	0.82
Completely Rural	2.30	2.75	2.14	2.53	1.12	1.20

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.83P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.0687		0.0106		0.3111	
GEOGRAPHIC DIVISION						
Northeast	0.2020		0.2748		0.9188	
New England	0.3096		0.1794		0.0898	
Middle Atlantic	0.3597		0.6055		0.3305	
Midwest	0.1995		0.0306		0.2200	
East North Central	0.0337		0.0010		0.0577	
West North Central	0.4557		0.4034		0.5504	
South	0.2695		0.1119		0.6541	
South Atlantic	0.8284		0.3014		0.6401	
East South Central	0.4420		0.5722		0.9503	
West South Central	0.2095		0.3150		0.9029	
West	0.9125		0.6035		0.5482	
Mountain	0.9912		0.9420		0.4735	
Pacific	0.8983		0.5828		0.7138	
COUNTY TYPE						
Large Metro	0.3085		0.1561		0.4978	
Small Metro	0.1291		0.0508		0.2241	
250K – 1 Mil. Pop.	0.6153		0.2080		0.8658	
< 250K Pop.	0.0762		0.1091		0.0878	
Nonmetro	0.4795		0.2376		0.6915	
Urbanized	0.7862		0.4660		0.9433	
Less Urbanized	0.6037		0.4801		0.4502	
Completely Rural	0.7062		0.5486		0.4023	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.84C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	1,647	1,681	924	910	400	446
GEOGRAPHIC DIVISION						
Northeast	703	681	419	350	186	174
New England	328	415	188	175	99	80
Middle Atlantic	625	539	376	303	157	155
Midwest	618	671	355	394	187	180
East North Central	488	532	284	331	148	145
West North Central	376	397	210	216	118	104
South	1,045	1,107	573	521	250	265
South Atlantic	759	856	402	381	186	205
East South Central	297	342	190	171	103	92
West South Central	596	630	341	305	133	139
West	894	883	446	520	193	265
Mountain	364	417	217	250	86	130
Pacific	814	772	388	454	173	230
COUNTY TYPE						
Large Metro	1,372	1,368	750	721	305	339
Small Metro	909	985	511	500	238	281
250K – 1 Mil. Pop.	784	811	427	397	206	227
< 250K Pop.	561	571	314	305	133	164
Nonmetro	619	663	351	392	176	159
Urbanized	483	537	248	285	135	103
Less Urbanized	391	441	244	247	106	113
Completely Rural	227	196	100	99	47	58

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.84D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.44	0.45	0.36	0.34	0.18	0.19
GEOGRAPHIC DIVISION						
Northeast	1.09	1.00	0.82	0.76	0.43	0.42
New England	1.72	1.88	1.44	1.37	0.86	0.74
Middle Atlantic	1.33	1.15	0.99	0.91	0.50	0.51
Midwest	0.79	0.78	0.64	0.70	0.38	0.37
East North Central	0.95	0.91	0.75	0.84	0.43	0.43
West North Central	1.45	1.66	1.23	1.25	0.80	0.71
South	0.80	0.81	0.62	0.56	0.31	0.32
South Atlantic	1.16	1.15	0.90	0.83	0.45	0.48
East South Central	1.58	2.04	1.28	1.14	0.79	0.67
West South Central	1.36	1.43	1.06	0.96	0.53	0.53
West	1.00	1.07	0.78	0.83	0.38	0.47
Mountain	1.32	1.54	1.18	1.31	0.56	0.79
Pacific	1.31	1.39	1.00	1.05	0.50	0.59
COUNTY TYPE						
Large Metro	0.65	0.65	0.53	0.50	0.26	0.27
Small Metro	0.81	0.81	0.63	0.60	0.35	0.39
250K – 1 Mil. Pop.	1.03	0.95	0.76	0.73	0.44	0.47
< 250K Pop.	1.30	1.51	1.11	1.06	0.59	0.69
Nonmetro	1.02	1.09	0.80	0.86	0.47	0.42
Urbanized	1.66	1.78	1.21	1.33	0.81	0.62
Less Urbanized	1.42	1.47	1.17	1.13	0.61	0.61
Completely Rural	3.08	3.37	2.12	1.95	1.14	1.42

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.84P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	0.9840		0.2817		0.5918	
GEOGRAPHIC DIVISION						
Northeast	0.3661		0.8113		0.1913	
New England	0.3656		0.7906		0.7939	
Middle Atlantic	0.5766		0.8938		0.1689	
Midwest	0.9396		0.2214		0.3022	
East North Central	0.2509		0.6511		0.4682	
West North Central	0.1444		0.1232		0.4642	
South	0.8493		0.1225		0.6280	
South Atlantic	0.4804		0.4286		0.6533	
East South Central	0.7223		0.4869		0.2868	
West South Central	0.3002		0.1627		0.4109	
West	0.3164		0.6022		0.1609	
Mountain	0.2125		0.5881		0.3981	
Pacific	0.6065		0.7460		0.2550	
COUNTY TYPE						
Large Metro	0.8973		0.5255		0.8715	
Small Metro	0.9881		0.4445		0.6527	
250K – 1 Mil. Pop.	0.7768		0.9703		0.6630	
< 250K Pop.	0.6819		0.2420		0.8892	
Nonmetro	0.8556		0.6092		0.7163	
Urbanized	0.8706		0.8624		0.7440	
Less Urbanized	0.9787		0.4368		0.5744	
Completely Rural	0.7620		0.6198		0.4372	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.