

**Table 5.14C Substance Dependence for Specific Substances in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2009 and 2010**

<b>Past Year Dependence</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
<b>ILLCIT DRUGS<sup>1,2</sup></b>	183	204	33	34	67	76	167	183
Marijuana and Hashish	117	154	29	29	59	64	94	135
Cocaine	86	82	7	6	21	20	82	80
Heroin	68	49	4	3	14	17	67	45
Hallucinogens	34	23	10	12	16	16	28	11
Inhalants	13	17	6	6	8	5	9	15
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	115	132	15	18	36	40	108	124
Pain Relievers	99	108	14	16	33	38	94	99
Tranquilizers	50	75	5	8	12	14	48	72
Stimulants <sup>2</sup>	38	41	8	7	16	17	33	36
Sedatives	33	36	3	4	11	5	31	35
<b>ALCOHOL</b>	294	304	31	30	77	81	284	289
<b>BOTH ILLICIT DRUGS AND ALCOHOL<sup>1,2</sup></b>	100	88	15	16	37	40	89	74
<b>ILLCIT DRUGS OR ALCOHOL<sup>1,2</sup></b>	340	364	42	42	94	103	324	340

\*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>2</sup> Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 5.14D Substance Dependence for Specific Substances in the Past Year, by Age Group: Standard Errors of Percentages, 2009 and 2010**

<b>Past Year Dependence</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
<b>ILLICIT DRUGS<sup>1,2</sup></b>	0.07	0.08	0.13	0.14	0.20	0.22	0.09	0.09
Marijuana and Hashish	0.05	0.06	0.12	0.12	0.18	0.19	0.05	0.07
Cocaine	0.03	0.03	0.03	0.02	0.06	0.06	0.04	0.04
Heroin	0.03	0.02	0.02	0.01	0.04	0.05	0.03	0.02
Hallucinogens	0.01	0.01	0.04	0.05	0.05	0.05	0.01	0.01
Inhalants	0.01	0.01	0.03	0.02	0.02	0.02	0.00	0.01
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.05	0.05	0.06	0.07	0.11	0.12	0.06	0.06
Pain Relievers	0.04	0.04	0.06	0.07	0.10	0.11	0.05	0.05
Tranquilizers	0.02	0.03	0.02	0.03	0.04	0.04	0.02	0.04
Stimulants <sup>2</sup>	0.02	0.02	0.03	0.03	0.05	0.05	0.02	0.02
Sedatives	0.01	0.01	0.01	0.02	0.03	0.02	0.02	0.02
<b>ALCOHOL</b>	0.12	0.12	0.13	0.12	0.23	0.24	0.15	0.15
<b>BOTH ILLICIT DRUGS AND ALCOHOL<sup>1,2</sup></b>	0.04	0.03	0.06	0.07	0.11	0.12	0.05	0.04
<b>ILLICIT DRUGS OR ALCOHOL<sup>1,2</sup></b>	0.14	0.14	0.17	0.17	0.28	0.30	0.17	0.17

\*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>2</sup> Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 5.14P Substance Dependence for Specific Substances in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

<b>Past Year Dependence</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
<b>ILLCIT DRUGS<sup>1,2</sup></b>	0.7406		0.2853		0.9738		0.5579	
Marijuana and Hashish	0.4456		0.6142		0.9340		0.4710	
Cocaine	0.5196		0.7512		0.1190		0.7830	
Heroin	0.6998		0.4012		0.2242		0.5009	
Hallucinogens	0.4499		0.3073		0.7705		0.1905	
Inhalants	0.8246		0.5600		0.3270		0.5519	
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.8698		0.1220		0.6682		0.8524	
Pain Relievers	0.9701		0.1636		0.7906		0.7295	
Tranquilizers	0.6407		0.2284		0.6231		0.8065	
Stimulants <sup>2</sup>	0.4979		0.2440		0.9065		0.6412	
Sedatives	0.7848		0.3271		0.4107		0.7000	
<b>ALCOHOL</b>	0.1221		0.1201		0.7236		0.1640	
<b>BOTH ILLICIT DRUGS AND ALCOHOL<sup>1,2</sup></b>	0.1218		0.9323		0.3538		0.1805	
<b>ILLCIT DRUGS OR ALCOHOL<sup>1,2</sup></b>	0.2707		0.8044		0.9526		0.2296	

\*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>2</sup> Estimates in these designated rows do not include data from new methamphetamine items added in 2005 and 2006.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.