

Table 6.11A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2009 and 2010

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
ILLICIT DRUGS²	5,687	5,385	7,597	7,366	4,010 ^b	4,939	4,520	4,933
Marijuana and Hashish	4,727	4,578	6,160	5,880	3,105 ^b	3,844	2,726	3,072
Cocaine	832	643	513	498	192	155	101	170
Crack	174	112	147	124	103	70	69	72
Heroin	59	51	42	47	34	70	60	70
Hallucinogens	554	546	439	400	144	130	121	132
LSD	88	78	41	47	9	16	20	13
PCP	34 ^a	4	12	7	4	*	*	24
Ecstasy	336	306	272	237	101	90	51	62
Inhalants	114	111	106	163	120	201	219	211
Nonmedical Use of Psychotherapeutics ^{3,4}	1,790	1,591	2,248	2,060	1,076 ^a	1,424	1,839	1,892
Pain Relievers	1,349	1,236	1,702	1,497	724 ^a	997	1,482	1,370
OxyContin [®]	153	135	188	192	26 ^a	90	144	147
Tranquilizers	607	456	626	676	348	511	429	517
Stimulants ⁴	457	371	465	336	165	133	203	237
Methamphetamine ⁴	163 ^a	72	172	115	80	66	87	100
Sedatives	103	71	103	125	54	29	111	148
ILLICIT DRUGS OTHER THAN MARIJUANA²	2,633	2,335	2,904	2,712	1,415	1,706	2,205	2,263

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.11B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2009 and 2010

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
ILLCIT DRUGS²	33.2	31.8	17.9	17.7	5.6 ^b	6.8	3.7	4.0
Marijuana and Hashish	27.6	27.0	14.5	14.1	4.4 ^a	5.3	2.2	2.5
Cocaine	4.9	3.8	1.2	1.2	0.3	0.2	0.1	0.1
Crack	1.0	0.7	0.3	0.3	0.1	0.1	0.1	0.1
Heroin	0.3	0.3	0.1	0.1	0.0	0.1	0.0	0.1
Hallucinogens	3.2	3.2	1.0	1.0	0.2	0.2	0.1	0.1
LSD	0.5	0.5	0.1	0.1	0.0	0.0	0.0	0.0
PCP	0.2 ^a	0.0	0.0	0.0	0.0	*	*	0.0
Ecstasy	2.0	1.8	0.6	0.6	0.1	0.1	0.0	0.1
Inhalants	0.7	0.7	0.3	0.4	0.2	0.3	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{3,4}	10.5	9.4	5.3	4.9	1.5	2.0	1.5	1.5
Pain Relievers	7.9	7.3	4.0	3.6	1.0	1.4	1.2	1.1
OxyContin [®]	0.9	0.8	0.4	0.5	0.0 ^a	0.1	0.1	0.1
Tranquilizers	3.5	2.7	1.5	1.6	0.5	0.7	0.4	0.4
Stimulants ⁴	2.7	2.2	1.1	0.8	0.2	0.2	0.2	0.2
Methamphetamine ⁴	1.0 ^a	0.4	0.4	0.3	0.1	0.1	0.1	0.1
Sedatives	0.6	0.4	0.2	0.3	0.1	0.0	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	15.4	13.8	6.8	6.5	2.0	2.3	1.8	1.9

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.12A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2009 and 2010

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
ILLICIT DRUGS²	362 ^a	289	697	740	327	363	1,085	1,075
Marijuana and Hashish	329 ^a	256	602	658	244	292	624	591
Cocaine	22	22	32	22	4	10	12	7
Crack	5	3	3	0	*	*	*	*
Heroin	7	2	1	3	*	1	5	2
Hallucinogens	72	52	80	83	17	30	48	50
LSD	18	17	6	15	2	7	9	6
PCP	7	2	3	2	1	*	2	3
Ecstasy	34	22	51	57	7	21	24	25
Inhalants	27	24	43	36	36	34	138	169
Nonmedical Use of Psychotherapeutics ^{3,4}	120	86	176	204	99	69	377	365
Pain Relievers	99	78	144	169	83	56	328	307
OxyContin [®]	29	13	19	17	7	2	14	14
Tranquilizers	45	28	48	44	21	10	29	51
Stimulants ⁴	37	23	37	42	14	10	40	32
Methamphetamine ⁴	10	5	10	11	4	2	12	6
Sedatives	6	5	6	10	3	*	25	20
ILLICIT DRUGS OTHER THAN MARIJUANA²	176 ^a	128	260	291	132	123	551	553

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.12B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2009 and 2010

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
ILLICIT DRUGS²	69.9	70.6	42.1 ^b	49.3	22.6	26.0	5.2	5.1
Marijuana and Hashish	63.6	62.5	36.3 ^b	43.8	16.8 ^a	20.9	3.0	2.8
Cocaine	4.2	5.3	2.0	1.4	0.3	0.7	0.1	0.0
Crack	1.1	0.8	0.2	0.0	*	*	*	*
Heroin	1.3	0.5	0.1	0.2	*	0.1	0.0	0.0
Hallucinogens	13.9	12.7	4.8	5.5	1.1	2.2	0.2	0.2
LSD	3.5	4.2	0.4	1.0	0.1	0.5	0.0	0.0
PCP	1.4	0.4	0.2	0.1	0.1	*	0.0	0.0
Ecstasy	6.5	5.4	3.1	3.8	0.5	1.5	0.1	0.1
Inhalants	5.3	5.8	2.6	2.4	2.5	2.4	0.7	0.8
Nonmedical Use of Psychotherapeutics ^{3,4}	23.2	21.1	10.6	13.6	6.8	4.9	1.8	1.7
Pain Relievers	19.2	19.2	8.7	11.3	5.7	4.0	1.6	1.5
OxyContin [®]	5.6	3.2	1.1	1.2	0.5	0.2	0.1	0.1
Tranquilizers	8.7	6.8	2.9	3.0	1.4	0.7	0.1	0.2
Stimulants ⁴	7.1	5.6	2.2	2.8	1.0	0.7	0.2	0.2
Methamphetamine ⁴	2.0	1.3	0.6	0.7	0.3	0.2	0.1	0.0
Sedatives	1.2	1.2	0.4	0.7	0.2	*	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA²	34.0	31.3	15.7	19.4	9.1	8.8	2.6	2.6

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.13A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2009 and 2010

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
ILLICIT DRUGS²	2,293	2,260	2,911	2,737	1,004 ^b	1,246	916 ^a	1,085
Marijuana and Hashish	2,006	1,999	2,550	2,346	872 ^b	1,106	654 ^b	842
Cocaine	295	282	129	159	25	33	14	23
Crack	13	29	15	31	8	9	4	3
Heroin	20	30	19	38	7	13	20	8
Hallucinogens	288	321	226	225	56	83	46	44
LSD	55	57	25	32	5	9	11	6
PCP	4	3	*	2	*	*	1	*
Ecstasy	161	178	151	133	42	67	23	28
Inhalants	44	39	54	57	21	23	29	16
Nonmedical Use of Psychotherapeutics ^{3,4}	780 ^a	645	791	736	226	283	322	340
Pain Relievers	575 ^a	468	593	565	169	227	264	259
OxyContin [®]	60	55	83	105	12	25	21 ^a	52
Tranquilizers	262	201	219	223	61	89	74	53
Stimulants ⁴	225	168	124	149	40	31	32	62
Methamphetamine ⁴	37	15	33	47	11	11	3	10
Sedatives	34	20	22	15	16	4	12	19
ILLICIT DRUGS OTHER THAN MARIJUANA²	1,083	1,006	1,041	954	293	365	377	385

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.13B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2009 and 2010

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
ILLICIT DRUGS²	49.7	48.7	31.0	29.8	14.8 ^b	17.5	7.1 ^a	8.3
Marijuana and Hashish	43.5	43.1	27.2	25.5	12.9 ^b	15.5	5.1 ^b	6.4
Cocaine	6.4	6.1	1.4	1.7	0.4	0.5	0.1	0.2
Crack	0.3	0.6	0.2	0.3	0.1	0.1	0.0	0.0
Heroin	0.4	0.6	0.2	0.4	0.1	0.2	0.2	0.1
Hallucinogens	6.3	6.9	2.4	2.5	0.8	1.2	0.4	0.3
LSD	1.2	1.2	0.3	0.3	0.1	0.1	0.1	0.0
PCP	0.1	0.1	*	0.0	*	*	0.0	*
Ecstasy	3.5	3.8	1.6	1.5	0.6	0.9	0.2	0.2
Inhalants	1.0	0.8	0.6	0.6	0.3	0.3	0.2	0.1
Nonmedical Use of Psychotherapeutics ^{3,4}	16.9 ^b	13.9	8.4	8.0	3.3	4.0	2.5	2.6
Pain Relievers	12.5 ^a	10.1	6.3	6.1	2.5	3.2	2.1	2.0
OxyContin [®]	1.3	1.2	0.9	1.1	0.2	0.4	0.2 ^a	0.4
Tranquilizers	5.7	4.3	2.3	2.4	0.9	1.2	0.6	0.4
Stimulants ⁴	4.9	3.6	1.3	1.6	0.6	0.4	0.3	0.5
Methamphetamine ⁴	0.8	0.3	0.4	0.5	0.2	0.2	0.0	0.1
Sedatives	0.7	0.4	0.2	0.2	0.2	0.1	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA²	23.5	21.7	11.1	10.4	4.3	5.1	2.9	2.9

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.14A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2009 and 2010

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
ILLCIT DRUGS²	3,032	2,836	3,988	3,889	2,679 ^a	3,329	2,519	2,774
Marijuana and Hashish	2,391	2,324	3,009	2,877	1,989	2,445	1,447	1,639
Cocaine	515	339	351	318	163	113	76	140
Crack	155	80	130	93	95	61	65	69
Heroin	32	19	22	6	27	56	35	60
Hallucinogens	194	174	132	92	71	17	27	38
LSD	15	4	10	*	*	*	*	*
PCP	23	*	9	3	3	*	*	21
Ecstasy	141	106	71	47	52	*	*	10
Inhalants	42	49	10 ^b	69	63	144	52	26
Nonmedical Use of Psychotherapeutics ^{3,4}	890	860	1,281	1,120	751	1,072	1,140	1,188
Pain Relievers	675	690	965	762	473	714	890	805
OxyContin [®]	64	67	86	70	7 ^a	62	109	81
Tranquilizers	300	227	358	408	266	412	326	412
Stimulants ⁴	196	180	304 ^a	144	111	91	130	143
Methamphetamine ⁴	116	51	129	56	65	52	73	85
Sedatives	63	47	74	101	36	25	74	109
ILLCIT DRUGS OTHER THAN MARIJUANA²	1,374	1,201	1,603	1,467	990	1,218	1,277	1,325

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.14B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2009 and 2010

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
ILLICIT DRUGS²	25.3	23.9	12.7	12.6	4.3 ^a	5.2	2.9	3.1
Marijuana and Hashish	19.9	19.6	9.6	9.3	3.2	3.8	1.7	1.9
Cocaine	4.3	2.9	1.1	1.0	0.3	0.2	0.1	0.2
Crack	1.3	0.7	0.4	0.3	0.2	0.1	0.1	0.1
Heroin	0.3	0.2	0.1	0.0	0.0	0.1	0.0	0.1
Hallucinogens	1.6	1.5	0.4	0.3	0.1	0.0	0.0	0.0
LSD	0.1	0.0	0.0	*	*	*	*	*
PCP	0.2	*	0.0	0.0	0.0	*	*	0.0
Ecstasy	1.2	0.9	0.2	0.2	0.1	*	*	0.0
Inhalants	0.4	0.4	0.0 ^b	0.2	0.1	0.2	0.1	0.0
Nonmedical Use of Psychotherapeutics ^{3,4}	7.4	7.2	4.1	3.6	1.2	1.7	1.3	1.3
Pain Relievers	5.6	5.8	3.1	2.5	0.8	1.1	1.0	0.9
OxyContin [®]	0.5	0.6	0.3	0.2	0.0 ^a	0.1	0.1	0.1
Tranquilizers	2.5	1.9	1.1	1.3	0.4	0.6	0.4	0.5
Stimulants ⁴	1.6	1.5	1.0 ^a	0.5	0.2	0.1	0.1	0.2
Methamphetamine ⁴	1.0	0.4	0.4	0.2	0.1	0.1	0.1	0.1
Sedatives	0.5	0.4	0.2	0.3	0.1	0.0	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA²	11.5	10.1	5.1	4.7	1.6	1.9	1.5	1.5

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.15A Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Numbers in Thousands, 2009 and 2010

Drug	Total (2009)	Total (2010)	Aged 12-20 (2009)	Aged 12-20 (2010)	Aged 21+ (2009)	Aged 21+ (2010)
ILLICIT DRUGS¹	17,294	17,689	3,911	3,899	13,382	13,790
Marijuana or Hashish	13,992	14,302	3,460	3,460	10,533	10,842
Cocaine (Including Crack)	1,536	1,296	202	203	1,334	1,093
Heroin	134	169	32	45	102	124
Hallucinogens	1,137	1,076	466	456	670	620
Inhalants	341	475	171	134	170 ^a	340
Pain Relievers ²	3,776	3,730	867	786	2,908	2,945
Tranquilizers ²	1,581	1,643	311	259	1,269	1,384
Stimulants ^{2,3}	1,087 ^a	840	234	222	853 ^a	618
Methamphetamine ³	415 ^a	253	41	56	373 ^a	197
Sedatives ²	260	226	34	27	226	199
ILLICIT DRUGS OTHER THAN MARIJUANA¹	6,952	6,754	1,503	1,433	5,449	5,320

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.15B Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Percentages, 2009 and 2010

Drug	Total (2009)	Total (2010)	Aged 12-20 (2009)	Aged 12-20 (2010)	Aged 21+ (2009)	Aged 21+ (2010)
ILLCIT DRUGS¹	13.2	13.5	37.6	39.0	11.1	11.4
Marijuana or Hashish	10.7	10.9	33.3	34.6	8.8	8.9
Cocaine (Including Crack)	1.2	1.0	1.9	2.0	1.1	0.9
Heroin	0.1	0.1	0.3	0.4	0.1	0.1
Hallucinogens	0.9	0.8	4.5	4.6	0.6	0.5
Inhalants	0.3	0.4	1.6	1.3	0.1	0.3
Pain Relievers ²	2.9	2.8	8.3	7.9	2.4	2.4
Tranquilizers ²	1.2	1.3	3.0	2.6	1.1	1.1
Stimulants ^{2,3}	0.8 ^a	0.6	2.2	2.2	0.7 ^a	0.5
Methamphetamine ³	0.3 ^a	0.2	0.4	0.6	0.3 ^a	0.2
Sedatives ²	0.2	0.2	0.3	0.3	0.2	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA¹	5.3	5.1	14.4	14.3	4.5	4.4

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.16A Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: Numbers in Thousands, 2009 and 2010

Drug¹	Total (2009)	Total (2010)	Aged 12-20 (2009)	Aged 12-20 (2010)	Aged 21+ (2009)	Aged 21+ (2010)
ILLCIT DRUGS²	7,782	8,091	1,790	1,870	5,992	6,221
Marijuana or Hashish	6,928	7,225	1,722	1,766	5,206	5,459
Cocaine (Including Crack)	388	303	30	46	357	257
Heroin	43	63	5	15	38	47
Hallucinogens	122	98	39	40	83	57
Inhalants	24	36	10	15	13	20
Pain Relievers ³	598	621	93	113	505	507
Tranquilizers ³	169	170	16	35	153	134
Stimulants ³	54	103	9	13	45	90
Methamphetamine ⁴	119	85	6	5	113	80
Sedatives ³	19	15	*	3	19	12
ILLCIT DRUGS OTHER THAN MARIJUANA²	1,376	1,258	171	226	1,205	1,032

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.16B Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: Percentages, 2009 and 2010

Drug¹	Total (2009)	Total (2010)	Aged 12-20 (2009)	Aged 12-20 (2010)	Aged 21+ (2009)	Aged 21+ (2010)
ILLCIT DRUGS²	6.0	6.2	17.5	19.0	5.0	5.1
Marijuana or Hashish	5.3	5.5	16.9	18.0	4.3	4.5
Cocaine (Including Crack)	0.3	0.2	0.3	0.5	0.3	0.2
Heroin	0.0	0.0	0.0	0.2	0.0	0.0
Hallucinogens	0.1	0.1	0.4	0.4	0.1	0.0
Inhalants	0.0	0.0	0.1	0.2	0.0	0.0
Pain Relievers ³	0.5	0.5	0.9	1.1	0.4	0.4
Tranquilizers ³	0.1	0.1	0.2	0.4	0.1	0.1
Stimulants ³	0.0	0.1	0.1	0.1	0.0	0.1
Methamphetamine ⁴	0.1	0.1	0.1	0.1	0.1	0.1
Sedatives ³	0.0	0.0	*	0.0	0.0	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA²	1.1	1.0	1.7 ^a	2.3	1.0	0.9

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.17A Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2009 and 2010

Drug	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
ILLCIT DRUGS¹	17,294	17,689	1,386	1,391	6,209	6,243	9,699	10,054
Marijuana or Hashish	13,992	14,302	1,175	1,205	5,428	5,450	7,389	7,646
Cocaine (Including Crack)	1,536	1,296	58	53	449	473	1,030	770
Heroin	134	169	8	6	46	81	81	82
Hallucinogens	1,137	1,076	169	165	571	629	397	282
Inhalants	341	475	106	94	119	119	115	262
Pain Relievers ²	3,776	3,730	325	304	1,337	1,260	2,114	2,166
Tranquilizers ²	1,581	1,643	114	82	542	513	924	1,048
Stimulants ^{2,3}	1,087 ^a	840	88	75	388	348	611	416
Methamphetamine ³	415 ^a	253	24	19	80	74	310 ^a	160
Sedatives ²	260	226	15	15	72	38	173	172
ILLCIT DRUGS OTHER THAN MARIJUANA¹	6,952	6,754	568	542	2,416	2,325	3,967	3,886

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.17B Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Percentages, 2009 and 2010

Drug	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
ILLCIT DRUGS¹	13.2	13.5	38.2 ^a	42.1	29.9	29.8	9.1	9.4
Marijuana or Hashish	10.7	10.9	32.4 ^a	36.4	26.1	26.0	7.0	7.1
Cocaine (Including Crack)	1.2	1.0	1.6	1.6	2.2	2.3	1.0	0.7
Heroin	0.1	0.1	0.2	0.2	0.2	0.4	0.1	0.1
Hallucinogens	0.9	0.8	4.7	5.0	2.7	3.0	0.4	0.3
Inhalants	0.3	0.4	2.9	2.8	0.6	0.6	0.1	0.2
Pain Relievers ²	2.9	2.8	9.0	9.2	6.4	6.0	2.0	2.0
Tranquilizers ²	1.2	1.3	3.2	2.5	2.6	2.4	0.9	1.0
Stimulants ^{2,3}	0.8 ^a	0.6	2.4	2.3	1.9	1.7	0.6	0.4
Methamphetamine ³	0.3 ^a	0.2	0.7	0.6	0.4	0.4	0.3 ^a	0.1
Sedatives ²	0.2	0.2	0.4	0.5	0.3 ^a	0.2	0.2	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA¹	5.3	5.1	15.7	16.4	11.6	11.1	3.7	3.6

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.18A Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2009 and 2010

Drug¹	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
ILLCIT DRUGS²	7,782	8,091	569	633	3,006	3,179	4,207	4,279
Marijuana or Hashish	6,928	7,225	535	608	2,834	2,962	3,559	3,656
Cocaine (Including Crack)	388	303	10	9	97	117	281	178
Heroin	43	63	3	*	6 ^a	32	33	31
Hallucinogens	122	98	9	14	63	55	49	29
Inhalants	24	36	8	12	9	15	8	9
Pain Relievers ³	598	621	24	37	199	199	375	384
Tranquilizers ³	169	170	3	10	58	55	109	105
Stimulants ³	54	103	2	3	20	50	32	50
Methamphetamine ⁴	119	85	2	2	19	13	98	70
Sedatives ³	19	15	0	3	8	2	11	10
ILLCIT DRUGS OTHER THAN MARIJUANA²	1,376	1,258	53	71	403	417	920	770

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.18B Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Percentages, 2009 and 2010

Drug¹	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
ILLCIT DRUGS²	6.0	6.2	16.1 ^b	19.6	14.6	15.3	4.0	4.0
Marijuana or Hashish	5.3	5.5	15.2 ^b	18.9	13.8	14.3	3.4	3.4
Cocaine (Including Crack)	0.3	0.2	0.3	0.3	0.5	0.6	0.3	0.2
Heroin	0.0	0.0	0.1	*	0.0 ^a	0.2	0.0	0.0
Hallucinogens	0.1	0.1	0.3	0.4	0.3	0.3	0.0	0.0
Inhalants	0.0	0.0	0.2	0.4	0.0	0.1	0.0	0.0
Pain Relievers ³	0.5	0.5	0.7	1.2	1.0	1.0	0.4	0.4
Tranquilizers ³	0.1	0.1	0.1	0.3	0.3	0.3	0.1	0.1
Stimulants ³	0.0	0.1	0.1	0.1	0.1	0.2	0.0	0.0
Methamphetamine ⁴	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1
Sedatives ³	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA²	1.1	1.0	1.5	2.2	1.9	2.0	0.9	0.7

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.