

**Table 6.24A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2009 and 2010**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	11,052	11,303
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	3,442	3,565	5,117	5,290
Cigars	7,496	7,154	5,773	6,023
Pipe Tobacco	1,105	1,176	982	974
<b>ALCOHOL</b>	39,375	38,538	91,246	92,804
Binge Alcohol Use <sup>2</sup>	26,252	25,476	33,309	33,074
Heavy Alcohol Use <sup>2</sup>	9,637	9,227	7,492	7,704

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.24B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Percentages, 2009 and 2010**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	5.7	5.8
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	5.9	6.1	2.6	2.7
Cigars	12.8	12.3	3.0	3.1
Pipe Tobacco	1.9	2.0	0.5	0.5
<b>ALCOHOL</b>	67.1	66.2	47.2	47.5
Binge Alcohol Use <sup>2</sup>	44.8	43.7	17.2	16.9
Heavy Alcohol Use <sup>2</sup>	16.4	15.8	3.9	3.9

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.25A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Numbers in Thousands, 2009 and 2010**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	670 <sup>a</sup>	566
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	303	282	274	280
Cigars	630 <sup>b</sup>	476	349	307
Pipe Tobacco	137	110	94 <sup>b</sup>	44
<b>ALCOHOL</b>	1,351	1,221	2,272	2,087
Binge Alcohol Use <sup>2</sup>	1,069	946	1,106 <sup>a</sup>	964
Heavy Alcohol Use <sup>2</sup>	379 <sup>b</sup>	285	138	124

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.25B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Percentages, 2009 and 2010**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	3.0 <sup>a</sup>	2.5
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	13.8	13.9	1.2	1.3
Cigars	28.8 <sup>b</sup>	23.4	1.6	1.4
Pipe Tobacco	6.2	5.4	0.4 <sup>b</sup>	0.2
<b>ALCOHOL</b>	61.7	60.1	10.1 <sup>a</sup>	9.4
Binge Alcohol Use <sup>2</sup>	48.8	46.6	4.9 <sup>a</sup>	4.3
Heavy Alcohol Use <sup>2</sup>	17.3 <sup>a</sup>	14.0	0.6	0.6

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.26A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Numbers in Thousands, 2009 and 2010**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	1,949 <sup>a</sup>	2,239
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1,269	1,290	784	894
Cigars	2,620	2,412	1,217 <sup>a</sup>	1,404
Pipe Tobacco	406	428	178	187
<b>ALCOHOL</b>	9,487	9,132	11,280	11,831
Binge Alcohol Use <sup>2</sup>	7,655	7,246	6,343	6,585
Heavy Alcohol Use <sup>2</sup>	3,090	2,956	1,522	1,683

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.26B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Percentages, 2009 and 2010**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	9.0 <sup>a</sup>	10.0
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	10.5	11.1	3.6	4.0
Cigars	21.8	20.7	5.6	6.3
Pipe Tobacco	3.4	3.7	0.8	0.8
<b>ALCOHOL</b>	78.9	78.3	52.3	52.8
Binge Alcohol Use <sup>2</sup>	63.6	62.1	29.4	29.4
Heavy Alcohol Use <sup>2</sup>	25.7	25.3	7.1	7.5

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.27A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2009 and 2010**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	8,433	8,498
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1,870	1,994	4,059	4,116
Cigars	4,245	4,267	4,207	4,313
Pipe Tobacco	562	637	710	743
<b>ALCOHOL</b>	28,536	28,185	77,694	78,886
Binge Alcohol Use <sup>2</sup>	17,528	17,284	25,860	25,525
Heavy Alcohol Use <sup>2</sup>	6,168	5,986	5,832	5,897

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.27B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Percentages, 2009 and 2010**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	5.7	5.6
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	4.2	4.5	2.7	2.7
Cigars	9.6	9.6	2.8	2.9
Pipe Tobacco	1.3	1.4	0.5	0.5
<b>ALCOHOL</b>	64.2	63.3	52.1	52.4
Binge Alcohol Use <sup>2</sup>	39.4	38.8	17.3	16.9
Heavy Alcohol Use <sup>2</sup>	13.9	13.4	3.9	3.9

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.