

Table 6.28A Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2009 and 2010

| Substance | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|-------------------------------------|------------------------------------|--------|--|--------|------------------------------------|--------|------------------|--------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2009 | 2010 | 2009 | 2010 | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS² | 11,463 | 10,927 | 20,159 | 19,848 | 16,165 | 15,839 | 21,926 | 22,944 |
| Cigarettes | 9,637 | 9,227 | 16,615 | 16,249 | 13,122 | 13,062 | 19,287 | 19,718 |
| Smokeless Tobacco | 2,149 | 1,992 | 2,761 | 2,779 | 1,571 | 1,604 | 2,079 | 2,480 |
| Cigars | 3,015 | 2,887 | 4,562 | 4,763 | 2,951 | 2,723 | 2,741 | 2,803 |
| Pipe Tobacco | 379 | 313 | 725 | 572 | 512 | 496 | 471 ^a | 768 |

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.28B Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2009 and 2010

| Substance | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|-------------------------------------|------------------------------------|------|--|------|------------------------------------|------|------------------|------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2009 | 2010 | 2009 | 2010 | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS² | 66.9 | 64.5 | 47.5 | 47.7 | 22.7 | 21.8 | 18.1 | 18.8 |
| Cigarettes | 56.3 | 54.5 | 39.2 | 39.0 | 18.5 | 17.9 | 15.9 | 16.1 |
| Smokeless Tobacco | 12.5 | 11.8 | 6.5 | 6.7 | 2.2 | 2.2 | 1.7 | 2.0 |
| Cigars | 17.6 | 17.1 | 10.8 | 11.4 | 4.2 | 3.7 | 2.3 | 2.3 |
| Pipe Tobacco | 2.2 | 1.9 | 1.7 | 1.4 | 0.7 | 0.7 | 0.4 ^a | 0.6 |

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.29A Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2009 and 2010

| Substance | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|-------------------------------------|------------------------------------|------|--|------|------------------------------------|------|--------|-------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2009 | 2010 | 2009 | 2010 | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS² | 431 ^b | 317 | 891 | 818 | 363 | 356 | 1,175 | 1,106 |
| Cigarettes | 379 ^b | 285 | 690 | 661 | 282 | 275 | 840 | 810 |
| Smokeless Tobacco | 87 | 63 | 204 | 199 | 53 | 64 | 232 | 235 |
| Cigars | 186 ^b | 121 | 312 | 286 | 147 | 104 | 335 | 272 |
| Pipe Tobacco | 56 ^a | 26 | 88 ^a | 51 | 27 | 21 | 60 | 56 |

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.29B Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2009 and 2010

| Substance | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|-------------------------------------|------------------------------------|------|--|------|------------------------------------|------|--------|------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2009 | 2010 | 2009 | 2010 | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS² | 83.3 | 77.6 | 53.8 | 54.5 | 25.0 | 25.5 | 5.6 | 5.3 |
| Cigarettes | 73.3 | 69.6 | 41.6 | 44.1 | 19.5 | 19.7 | 4.0 | 3.8 |
| Smokeless Tobacco | 16.8 | 15.5 | 12.3 | 13.3 | 3.7 | 4.6 | 1.1 | 1.1 |
| Cigars | 35.9 | 29.6 | 18.8 | 19.0 | 10.1 | 7.5 | 1.6 | 1.3 |
| Pipe Tobacco | 10.8 | 6.3 | 5.3 | 3.4 | 1.9 | 1.5 | 0.3 | 0.3 |

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.30A Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2009 and 2010

| Substance | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|-------------------------------------|------------------------------------|-------|--|-------|------------------------------------|-------|--------|-------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2009 | 2010 | 2009 | 2010 | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS² | 3,572 | 3,466 | 5,310 | 5,169 | 2,230 | 2,313 | 2,868 | 2,957 |
| Cigarettes | 3,090 | 2,956 | 4,565 | 4,290 | 1,832 | 1,886 | 2,544 | 2,534 |
| Smokeless Tobacco | 843 | 761 | 737 | 830 | 163 | 222 | 310 | 370 |
| Cigars | 1,155 | 1,144 | 1,452 | 1,431 | 604 | 565 | 626 | 676 |
| Pipe Tobacco | 184 | 146 | 220 | 258 | 91 | 96 | 89 | 115 |

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.30B Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2009 and 2010

| Substance | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|-------------------------------------|------------------------------------|------|--|------|------------------------------------|------|--------|------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2009 | 2010 | 2009 | 2010 | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS² | 77.4 | 74.7 | 56.6 | 56.2 | 32.9 | 32.4 | 22.4 | 22.6 |
| Cigarettes | 67.0 | 63.7 | 48.6 | 46.7 | 27.1 | 26.4 | 19.9 | 19.3 |
| Smokeless Tobacco | 18.3 | 16.4 | 7.9 | 9.0 | 2.4 | 3.1 | 2.4 | 2.8 |
| Cigars | 25.1 | 24.7 | 15.5 | 15.6 | 8.9 | 7.9 | 4.9 | 5.2 |
| Pipe Tobacco | 4.0 | 3.1 | 2.3 | 2.8 | 1.3 | 1.3 | 0.7 | 0.9 |

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.31A Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2009 and 2010

| Substance | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|-------------------------------------|------------------------------------|-------|--|--------|------------------------------------|--------|------------------|--------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2009 | 2010 | 2009 | 2010 | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS² | 7,461 | 7,144 | 13,957 | 13,861 | 13,572 | 13,170 | 17,883 | 18,881 |
| Cigarettes | 6,168 | 5,986 | 11,360 | 11,298 | 11,009 | 10,901 | 15,903 | 16,375 |
| Smokeless Tobacco | 1,219 | 1,168 | 1,819 | 1,750 | 1,355 | 1,317 | 1,536 | 1,875 |
| Cigars | 1,673 | 1,622 | 2,799 | 3,047 | 2,200 | 2,055 | 1,780 | 1,856 |
| Pipe Tobacco | 139 | 142 | 418 | 263 | 394 | 380 | 322 ^a | 596 |

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.31B Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2009 and 2010

| Substance | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|-------------------------------------|------------------------------------|------|--|------|------------------------------------|------|------------------|------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2009 | 2010 | 2009 | 2010 | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS² | 62.2 | 60.1 | 44.5 | 44.8 | 21.6 | 20.5 | 20.5 | 21.4 |
| Cigarettes | 51.4 | 50.4 | 36.2 | 36.5 | 17.5 | 17.0 | 18.2 | 18.6 |
| Smokeless Tobacco | 10.2 | 9.8 | 5.8 | 5.7 | 2.2 | 2.0 | 1.8 | 2.1 |
| Cigars | 13.9 | 13.6 | 8.9 | 9.9 | 3.5 | 3.2 | 2.0 | 2.1 |
| Pipe Tobacco | 1.2 | 1.2 | 1.3 | 0.9 | 0.6 | 0.6 | 0.4 ^a | 0.7 |

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.