

Table 6.47A Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	1,997	1,970	255	252	515	492	1,228	1,227
From More Than One Doctor	263	236	35	27	49	64	178	144
Wrote Fake Prescription	26	21	2	6	6	2	17	13
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	58 ^a	28	10	19	26 ^b	3	21	6
From Friend or Relative for Free	6,211	6,254	610	572	2,050	2,006	3,550	3,676
Bought from Friend or Relative	988 ^b	1,293	124	111	469 ^a	541	396 ^b	641
Took from Friend or Relative without Asking	603	549	168	140	189	170	245	239
Bought from Drug Dealer or Other Stranger	481	497	63	50	235	238	183	209
Bought on the Internet	43	44	3	0	7	3	33	41
Some Other Way	447	479	78	94	128	144	242	241

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.47B Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	18.0	17.3	18.9	19.8	14.0	13.4	20.1	19.1
From More Than One Doctor	2.4	2.1	2.6	2.1	1.3	1.8	2.9	2.2
Wrote Fake Prescription	0.2	0.2	0.2	0.5	0.2	0.1	0.3	0.2
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5 ^a	0.2	0.8	1.5	0.7 ^b	0.1	0.3	0.1
From Friend or Relative for Free	55.9	55.0	45.3	45.0	55.8	54.8	58.3	57.1
Bought from Friend or Relative	8.9 ^b	11.4	9.2	8.8	12.8 ^a	14.8	6.5 ^b	10.0
Took from Friend or Relative without Asking	5.4	4.8	12.5	11.0	5.2	4.7	4.0	3.7
Bought from Drug Dealer or Other Stranger	4.3	4.4	4.6	3.9	6.4	6.5	3.0	3.2
Bought on the Internet	0.4	0.4	0.2	0.0	0.2	0.1	0.5	0.6
Some Other Way	4.0	4.2	5.8	7.4	3.5	3.9	4.0	3.7

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.48A Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	4,489	4,453	310	287	1,339	1,281	2,841	2,886
From More Than One Doctor	188	200	27	28	58	69	102	102
Wrote Fake Prescription	3	13	0	1	3	1	*	11
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	31	14	5	5	8	8	18	2
From Friend or Relative for Free	343	356	66	47	148	148	129	161
Bought from Friend or Relative	220	255	31	30	119	127	70	98
Took from Friend or Relative without Asking	76	109	33	51	23	38	19	21
Bought from Drug Dealer or Other Stranger	89	129	20	17	47	68	23	44
Bought on the Internet	7	11	0	1	2	3	5	7
Some Other Way	52	70	18	21	23	22	10	27

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.48B Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	81.7	79.4	60.7	58.8	75.6	72.6	88.3	85.9
From More Than One Doctor	3.4	3.6	5.3	5.8	3.3	3.9	3.2	3.0
Wrote Fake Prescription	0.1	0.2	0.1	0.2	0.2	0.1	*	0.3
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.6	0.3	1.0	1.0	0.5	0.4	0.6	0.0
From Friend or Relative for Free	6.2	6.3	12.9	9.6	8.3	8.4	4.0	4.8
Bought from Friend or Relative	4.0	4.6	6.0	6.1	6.7	7.2	2.2	2.9
Took from Friend or Relative without Asking	1.4	2.0	6.5 ^a	10.5	1.3	2.1	0.6	0.6
Bought from Drug Dealer or Other Stranger	1.6	2.3	3.9	3.5	2.6	3.8	0.7	1.3
Bought on the Internet	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.2
Some Other Way	0.9	1.3	3.6	4.4	1.3	1.3	0.3	0.8

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.49A Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	1,227	1,276	129	123	277	289	821	863
From More Than One Doctor	273	231	21	25	65	63	186	144
Wrote Fake Prescription	20	27	3	5	9	2	9	19
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	50	30	12	12	18 ^a	6	20	11
From Friend or Relative for Free	2,626	2,835	247	244	858	936	1,521	1,655
Bought from Friend or Relative	1,008 ^b	1,260	101	88	437	496	470 ^a	676
Took from Friend or Relative without Asking	510	525	106	89	172	185	232	252
Bought from Drug Dealer or Other Stranger	536	540	65	55	245	269	226	216
Bought on the Internet	44	48	3	*	14	13	27	35
Some Other Way	225	301	48	41	72	96	105	164

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.49B Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	27.4	27.1	25.5	24.7	20.0	20.1	31.7	31.2
From More Than One Doctor	6.1	4.9	4.2	4.9	4.7	4.4	7.2	5.2
Wrote Fake Prescription	0.4	0.6	0.5	1.1	0.6	0.2	0.3	0.7
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.1	0.6	2.4	2.4	1.3 ^a	0.4	0.8	0.4
From Friend or Relative for Free	58.6	60.1	48.6	48.5	61.9	64.8	58.7	59.7
Bought from Friend or Relative	22.5 ^a	26.8	19.9	17.5	31.6	34.4	18.2 ^a	24.4
Took from Friend or Relative without Asking	11.4	11.2	20.9	17.8	12.4	12.8	9.0	9.1
Bought from Drug Dealer or Other Stranger	12.0	11.5	12.8	11.1	17.7	18.6	8.7	7.8
Bought on the Internet	1.0	1.0	0.6	*	1.0	0.9	1.0	1.3
Some Other Way	5.1	6.5	9.9	8.5	5.3	6.8	4.1	6.0

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.50A Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	1,832	1,967	127	116	563	626	1,142	1,224
From More Than One Doctor	238	222	29	24	81	80	128	118
Wrote Fake Prescription	24	33	3	1	11	9	10	23
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	52	23	9	3	16	16	28	4
From Friend or Relative for Free	370	373	53	39	141	157	176	177
Bought from Friend or Relative	322	342	34	38	144	157	144	147
Took from Friend or Relative without Asking	97	111	29	28	33	52	35	31
Bought from Drug Dealer or Other Stranger	176	219	29	23	90	115	58	81
Bought on the Internet	28 ^a	9	3	1	15 ^a	5	10	3
Some Other Way	68	74	14	18	18 ^a	36	36	20

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.50B Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	77.3	76.7	60.6	54.8	73.4	72.3	81.8	82.5
From More Than One Doctor	10.2	8.8	14.0	11.7	10.7	9.4	9.4	8.0
Wrote Fake Prescription	1.0	1.3	1.6	0.5	1.4	1.0	0.8	1.5
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2.2	0.9	4.1	1.4	2.1	1.9	2.0	0.2
From Friend or Relative for Free	15.8	14.7	25.6	18.9	18.6	18.3	12.8	12.1
Bought from Friend or Relative	13.8	13.5	16.4	18.1	18.9	18.3	10.5	10.0
Took from Friend or Relative without Asking	4.2	4.4	13.9	13.5	4.4	6.1	2.6	2.1
Bought from Drug Dealer or Other Stranger	7.5	8.6	13.9	10.8	11.8	13.5	4.2	5.5
Bought on the Internet	1.2 ^b	0.4	1.4	0.7	2.0 ^a	0.6	0.7	0.2
Some Other Way	3.0	3.0	7.1	9.0	2.5	4.3	2.7	1.4

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.51A Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	433 ^b	628	25	33	92 ^a	127	316 ^a	469
From More Than One Doctor	50	58	4	3	17	15	29	40
Wrote Fake Prescription	6	3	4	2	3	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	13	10	3	3	10	7	*	*
From Friend or Relative for Free	2,879	3,032	181	188	861	885	1,837	1,959
Bought from Friend or Relative	626	669	73	57	309	318	244	294
Took from Friend or Relative without Asking	274	190	43	37	87	75	143	78
Bought from Drug Dealer or Other Stranger	273	295	43	35	154	148	76	112
Bought on the Internet	91	21	3	1	4	3	84	17
Some Other Way	160	173	18	21	56	54	86	99

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.51B Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	9.0 ^a	12.4	6.3	8.6	5.8	7.8	11.2	15.3
From More Than One Doctor	1.0	1.1	1.1	0.9	1.1	0.9	1.0	1.3
Wrote Fake Prescription	0.1	0.1	0.9	0.5	0.2	0.1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3	0.2	0.8	0.8	0.6	0.5	*	*
From Friend or Relative for Free	59.9	59.7	45.7	49.7	54.1	54.2	65.2	63.9
Bought from Friend or Relative	13.0	13.2	18.5	15.0	19.4	19.5	8.7	9.6
Took from Friend or Relative without Asking	5.7 ^a	3.7	10.9	9.7	5.5	4.6	5.1	2.5
Bought from Drug Dealer or Other Stranger	5.7	5.8	10.7	9.3	9.7	9.0	2.7	3.7
Bought on the Internet	1.9	0.4	0.7	0.2	0.2	0.2	3.0	0.6
Some Other Way	3.3	3.4	4.5	5.4	3.5	3.3	3.0	3.2

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.52A Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	1,939	2,118	62 ^a	89	491	516	1,385	1,513
From More Than One Doctor	93	85	9	7	23	35	61	43
Wrote Fake Prescription	1	5	1	*	1	1	*	2
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	9	6	0	*	6	1	3	5
From Friend or Relative for Free	257	236	28	23	111	111	119	102
Bought from Friend or Relative	145 ^b	83	18	12	81	56	46 ^a	15
Took from Friend or Relative without Asking	33	25	17	12	9	11	*	2
Bought from Drug Dealer or Other Stranger	54	75	7	9	20 ^a	42	26	25
Bought on the Internet	5	8	0	*	4	1	0	7
Some Other Way	38	22	4	6	12	4	21	12

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.52B Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	75.3 ^a	79.6	42.1 ^a	55.6	64.7	66.3	83.0	87.7
From More Than One Doctor	3.6	3.2	6.1	4.6	3.0	4.5	3.7	2.5
Wrote Fake Prescription	0.1	0.2	0.3	*	0.1	0.1	*	0.1
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.4	0.2	0.3	*	0.8	0.1	0.2	0.3
From Friend or Relative for Free	10.0	8.9	18.7	14.3	14.6	14.3	7.1	5.9
Bought from Friend or Relative	5.6 ^b	3.1	12.4	7.4	10.7 ^a	7.2	2.8 ^a	0.9
Took from Friend or Relative without Asking	1.3	0.9	11.8	7.2	1.2	1.4	*	0.1
Bought from Drug Dealer or Other Stranger	2.1	2.8	5.0	5.3	2.7 ^a	5.3	1.6	1.4
Bought on the Internet	0.2	0.3	0.2	*	0.6	0.1	0.0	0.4
Some Other Way	1.5	0.8	3.0	3.9	1.6	0.6	1.3	0.7

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.53A Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	217 ^a	316	12	13	44	68	161	235
From More Than One Doctor	34	17	*	*	8	4	25	11
Wrote Fake Prescription	3	2	*	1	2	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	7	2	*	*	6	*	*	*
From Friend or Relative for Free	1,014	1,108	63 ^a	42	279	274	671	793
Bought from Friend or Relative	282 ^b	419	22	26	136	157	124 ^a	236
Took from Friend or Relative without Asking	125	96	16	13	45	43	64	39
Bought from Drug Dealer or Other Stranger	183	143	18	16	76	77	89	50
Bought on the Internet	46	13	3	1	3	4	*	8
Some Other Way	55	58	6	5	16	15	33	38

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.53B Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	13.0	16.8	9.1	12.2	8.5	12.5	15.9	19.1
From More Than One Doctor	2.0	0.9	*	*	1.5	0.8	2.5	0.9
Wrote Fake Prescription	0.2	0.1	*	1.2	0.4	0.1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.4	0.1	*	*	1.1	*	*	*
From Friend or Relative for Free	60.9	58.9	49.5 ^a	38.0	53.7	50.6	66.0	64.4
Bought from Friend or Relative	17.0 ^a	22.3	17.5	24.2	26.1	29.0	12.3	19.1
Took from Friend or Relative without Asking	7.5	5.1	12.6	12.4	8.6	8.0	6.3	3.2
Bought from Drug Dealer or Other Stranger	11.0	7.6	14.1	15.3	14.6	14.2	8.8	4.1
Bought on the Internet	2.8	0.7	2.1	1.0	0.7	0.8	*	0.6
Some Other Way	3.3	3.1	4.8	5.3	3.1	2.8	3.3	3.1

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.54A Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	708	781	*	*	164	165	518	600
From More Than One Doctor	85	52	*	*	21	21	62	30
Wrote Fake Prescription	0	2	*	*	*	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	3	*	*	*	1	*	*
From Friend or Relative for Free	120	130	*	*	63	48	45	76
Bought from Friend or Relative	72	67	*	*	37	36	24	25
Took from Friend or Relative without Asking	20	21	*	*	6	2	8	*
Bought from Drug Dealer or Other Stranger	47	55	*	*	16	19	27	33
Bought on the Internet	1	6	*	*	1	*	0	*
Some Other Way	23	8	*	*	6	4	16	2

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.54B Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	75.2	76.6	*	*	63.4	65.6	82.4	81.8
From More Than One Doctor	9.0	5.1	*	*	8.1	8.4	9.9	4.0
Wrote Fake Prescription	0.0	0.1	*	*	*	0.6	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2	0.3	*	*	*	0.6	*	*
From Friend or Relative for Free	12.8	12.8	*	*	24.3	19.1	7.2	10.4
Bought from Friend or Relative	7.7	6.6	*	*	14.4	14.5	3.8	3.4
Took from Friend or Relative without Asking	2.1	2.1	*	*	2.1	0.7	1.2	*
Bought from Drug Dealer or Other Stranger	5.0	5.4	*	*	6.3	7.5	4.3	4.5
Bought on the Internet	0.1	0.5	*	*	0.5	*	0.0	*
Some Other Way	2.4	0.9	*	*	2.3	1.5	2.6	0.3

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.55A Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	187	229	25	23	53 ^a	85	109	122
From More Than One Doctor	18	18	4	4	10	5	*	9
Wrote Fake Prescription	6	10	3	3	3	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	19	21	4	5	10 ^a	2	5	14
From Friend or Relative for Free	1,018	1,121	126 ^a	96	423 ^b	529	468	496
Bought from Friend or Relative	346	417	40	44	202 ^b	270	103	103
Took from Friend or Relative without Asking	108	106	30	21	34	34	44	50
Bought from Drug Dealer or Other Stranger	141	141	24	19	69	62	48	60
Bought on the Internet	58	34	1	3	14	9	44	23
Some Other Way	50	86	14	8	18	28	18	50

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.55B Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	9.6	10.5	9.3	10.0	6.3	8.3	12.9	13.0
From More Than One Doctor	0.9	0.8	1.5	1.7	1.1	0.5	*	1.0
Wrote Fake Prescription	0.3	0.4	1.1	1.2	0.3	0.0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.0	1.0	1.3	2.3	1.2 ^a	0.2	0.6	1.5
From Friend or Relative for Free	52.1	51.3	46.7	42.5	50.6	51.7	55.4	53.1
Bought from Friend or Relative	17.7	19.1	14.7	19.7	24.2	26.3	12.2	11.1
Took from Friend or Relative without Asking	5.5	4.9	11.1	9.5	4.1	3.3	5.2	5.4
Bought from Drug Dealer or Other Stranger	7.2	6.5	9.0	8.5	8.3	6.0	5.7	6.5
Bought on the Internet	3.0	1.6	0.3	1.1	1.7	0.9	5.2	2.4
Some Other Way	2.6	3.9	5.0	3.5	2.2	2.7	2.1	5.4

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.56A Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Stimulants ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	667	750	66	53	296	353	*	343
From More Than One Doctor	28	39	8	7	11	17	8	16
Wrote Fake Prescription	0	1	0	*	*	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	5	1	*	1	5	*	*
From Friend or Relative for Free	105	77	12	9	46	41	*	*
Bought from Friend or Relative	59	47	13	6	27	36	19	5
Took from Friend or Relative without Asking	9	7	6	3	3	3	*	1
Bought from Drug Dealer or Other Stranger	9 ^b	44	5	4	4 ^a	19	*	*
Bought on the Internet	6	8	*	*	3	*	*	*
Some Other Way	8	15	1	1	2	1	5	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.56B Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Stimulants ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	74.7	75.5	59.5	63.2	75.4	74.1	*	79.3
From More Than One Doctor	3.1	3.9	7.4	8.0	2.9	3.5	2.1	3.6
Wrote Fake Prescription	0.0	0.1	0.1	*	*	0.1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2	0.5	0.7	*	0.2	1.0	*	*
From Friend or Relative for Free	11.8	7.8	10.7	10.4	11.8	8.7	*	*
Bought from Friend or Relative	6.6	4.8	11.5	7.2	6.8	7.6	5.0	1.2
Took from Friend or Relative without Asking	1.0	0.7	5.3	3.7	0.7	0.6	*	0.2
Bought from Drug Dealer or Other Stranger	1.0 ^b	4.4	4.3	5.0	1.1 ^a	4.1	*	*
Bought on the Internet	0.7	0.8	*	*	0.7	*	*	*
Some Other Way	0.9	1.5	0.6	0.6	0.5	0.3	1.4	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.57A Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	89 ^b	173	14	14	27 ^b	57	*	102
From More Than One Doctor	21	19	4	2	13	5	*	12
Wrote Fake Prescription	5	3	3	2	2	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	12	9	2	*	5	1	*	*
From Friend or Relative for Free	343	405	35	34	128 ^a	181	*	190
Bought from Friend or Relative	166	208	19	15	83 ^a	122	64	71
Took from Friend or Relative without Asking	53	80	14	9	19	23	*	47
Bought from Drug Dealer or Other Stranger	72	89	10	16	41	38	21	35
Bought on the Internet	34	17	*	1	9	5	*	*
Some Other Way	15	32	7	4	6	7	*	21

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.57B Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	13.4 ^a	20.5	16.9	16.5	10.4	16.4	*	24.9
From More Than One Doctor	3.2	2.3	4.3	2.8	4.9	1.6	*	2.8
Wrote Fake Prescription	0.8	0.3	3.4	2.6	0.9	0.2	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.8	1.1	1.8	*	2.0	0.2	*	*
From Friend or Relative for Free	51.4	48.0	41.2	39.9	48.9	52.2	*	46.1
Bought from Friend or Relative	24.9	24.7	22.2	17.9	31.8	35.3	20.1	17.2
Took from Friend or Relative without Asking	8.0	9.5	16.0	11.2	7.2	6.7	*	11.5
Bought from Drug Dealer or Other Stranger	10.7	10.5	11.9	18.8	15.7	10.8	6.4	8.5
Bought on the Internet	5.1	2.0	*	1.2	3.5	1.5	*	*
Some Other Way	2.3	4.0	8.2	4.9	2.2	2.2	*	5.3

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.58A Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Stimulants ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	*	265	*	*	91	119	*	*
From More Than One Doctor	11	20	*	*	5	12	*	*
Wrote Fake Prescription	*	0	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	0	*	*	*	*	*	*
From Friend or Relative for Free	*	63	*	*	23	23	*	*
Bought from Friend or Relative	*	30	*	*	9	17	*	*
Took from Friend or Relative without Asking	9	11	*	*	4	*	*	*
Bought from Drug Dealer or Other Stranger	20	14	*	*	5	9	*	*
Bought on the Internet	7	*	*	*	*	*	*	*
Some Other Way	9	*	*	*	*	1	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.58B Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Stimulants ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	*	73.1	*	*	74.7	75.5	*	*
From More Than One Doctor	3.5	5.6	*	*	4.0	7.6	*	*
Wrote Fake Prescription	*	0.0	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	0.1	*	*	*	*	*	*
From Friend or Relative for Free	*	17.9	*	*	18.5	14.8	*	*
Bought from Friend or Relative	*	8.6	*	*	7.1	10.6	*	*
Took from Friend or Relative without Asking	2.9	3.2	*	*	3.0	*	*	*
Bought from Drug Dealer or Other Stranger	6.2	4.1	*	*	4.0	6.1	*	*
Bought on the Internet	2.3	*	*	*	*	*	*	*
Some Other Way	2.8	*	*	*	*	0.3	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.59A Source Where Methamphetamine was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From Friend or Relative for Free	419	389	46	*	158	128	214	223
Bought from Friend or Relative	294	288	15	9	68	55	211	224
Took from Friend or Relative without Asking	19	40	6	6	7	5	6	29
Bought from Drug Dealer or Other Stranger	212	175	9	10	57	58	146	107
Bought on the Internet	6	7	*	*	0	3	*	*
Some Other Way	27	40	2	*	12	6	14	28

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.59B Source Where Methamphetamine was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From Friend or Relative for Free	42.8	41.5	55.6	*	52.4	50.3	36.2	36.3
Bought from Friend or Relative	30.1	30.7	17.6	13.2	22.5	21.5	35.7	36.4
Took from Friend or Relative without Asking	1.9	4.2	6.7	8.9	2.4	2.0	1.0	4.6
Bought from Drug Dealer or Other Stranger	21.7	18.7	10.8	15.1	18.8	22.7	24.7	17.4
Bought on the Internet	0.7	0.7	*	*	0.1	1.0	*	*
Some Other Way	2.8	4.2	1.8	*	3.9	2.4	2.4	4.6

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.60A Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From Friend or Relative for Free	114	141	*	*	46	*	*	*
Bought from Friend or Relative	126	101	*	*	40	33	*	*
Took from Friend or Relative without Asking	4	11	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	90	51	*	*	32	19	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	14	9	*	*	8	7	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.60B Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From Friend or Relative for Free	32.9	44.7	*	*	35.6	*	*	*
Bought from Friend or Relative	36.3	32.0	*	*	31.1	30.7	*	*
Took from Friend or Relative without Asking	1.1	3.3	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	25.8	16.3	*	*	24.6	18.0	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	4.0	2.9	*	*	6.5	6.7	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.61A Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From Friend or Relative for Free	*	166	*	*	*	*	*	*
Bought from Friend or Relative	*	143	*	*	23	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	108	*	*	*	*	*	*
Bought on the Internet	0	4	*	*	*	*	*	*
Some Other Way	20	20	*	*	4	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.61B Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From Friend or Relative for Free	*	44.1	*	*	*	*	*	*
Bought from Friend or Relative	*	38.0	*	*	29.4	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	28.7	*	*	*	*	*	*
Bought on the Internet	0.0	1.0	*	*	*	*	*	*
Some Other Way	5.7	5.5	*	*	5.3	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.62A Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	5	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.62B Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	3.1	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.63A Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	102	112	9	16	16	28	*	69
From More Than One Doctor	20	20	2	1	*	4	17	*
Wrote Fake Prescription	8	1	*	0	*	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	8	5	1	*	3	*	*	*
From Friend or Relative for Free	419	465	41	42	100 ^a	70	*	353
Bought from Friend or Relative	40	72	9	15	19	19	12	38
Took from Friend or Relative without Asking	23	40	11	9	6	*	5	14
Bought from Drug Dealer or Other Stranger	30	33	6	2	6	5	18	26
Bought on the Internet	3	*	*	*	0	*	*	*
Some Other Way	24	33	4	3	4	4	17	25

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.63B Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	15.1	14.2	10.6	17.9	10.4 ^a	18.7	*	12.4
From More Than One Doctor	3.0	2.6	2.9	1.4	*	2.5	3.9	*
Wrote Fake Prescription	1.2	0.1	*	0.5	*	0.3	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.2	0.6	0.9	*	2.1	*	*	*
From Friend or Relative for Free	61.9	58.8	48.8	46.2	63.3 ^a	47.2	*	64.0
Bought from Friend or Relative	5.9	9.1	10.9	16.4	11.8	13.1	2.7	6.8
Took from Friend or Relative without Asking	3.3	5.1	12.8	10.3	3.9	*	1.3	2.5
Bought from Drug Dealer or Other Stranger	4.4	4.1	6.8	2.7	3.9	3.1	4.1	4.6
Bought on the Internet	0.4	*	*	*	0.1	*	*	*
Some Other Way	3.6	4.2	4.1	3.5	2.4	2.9	4.0	4.6

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.64A Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Sedatives ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	319	370	*	*	67	*	*	*
From More Than One Doctor	9	4	*	*	2	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	4	*	*	*	*	*	*
From Friend or Relative for Free	21	24	*	*	9	*	*	*
Bought from Friend or Relative	4	5	*	*	*	*	*	*
Took from Friend or Relative without Asking	1	5	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	1	8	*	*	*	*	*	*
Bought on the Internet	0	*	*	*	0	*	*	*
Some Other Way	3	3	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.64B Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Sedatives ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	88.2	86.1	*	*	79.7	*	*	*
From More Than One Doctor	2.4	1.0	*	*	2.6	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.2	1.0	*	*	*	*	*	*
From Friend or Relative for Free	5.7	5.5	*	*	10.3	*	*	*
Bought from Friend or Relative	1.1	1.2	*	*	*	*	*	*
Took from Friend or Relative without Asking	0.2	1.1	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	0.3	1.9	*	*	*	*	*	*
Bought on the Internet	0.0	*	*	*	0.1	*	*	*
Some Other Way	0.8	0.6	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.65A Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source for Past Month Nonmedical Use among Past Month Sedative Users ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	42	81	*	*	*	*	*	*
From More Than One Doctor	14	7	*	*	*	*	*	*
Wrote Fake Prescription	*	1	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	5	*	*	*	*	*	*
From Friend or Relative for Free	181	*	*	*	*	*	*	*
Bought from Friend or Relative	27	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	19	15	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	23	*	*	*	*	*	*	*
Bought on the Internet	1	0	*	*	*	*	*	*
Some Other Way	10	19	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.65B Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source for Past Month Nonmedical Use among Past Month Sedative Users ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	16.4	23.7	*	*	*	*	*	*
From More Than One Doctor	5.5	2.1	*	*	*	*	*	*
Wrote Fake Prescription	*	0.2	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	1.6	*	*	*	*	*	*
From Friend or Relative for Free	70.1	*	*	*	*	*	*	*
Bought from Friend or Relative	10.3	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	7.3	4.2	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	9.1	*	*	*	*	*	*	*
Bought on the Internet	0.4	0.0	*	*	*	*	*	*
Some Other Way	3.9	5.7	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.66A Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Sedatives ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.66B Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Source Where Friend or Relative Obtained Sedatives ¹	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.