

Table 6.71A Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2007-2008	2009-2010	2007-2008	2009-2010	2007-2008	2009-2010
ILLCIT DRUGS²	5,955 ^b	6,537	130	109	5,777 ^b	6,383
Marijuana and Hashish	4,316 ^b	4,904	97	90	4,181 ^b	4,785
Cocaine	501 ^a	385	9	3	487 ^a	378
Crack	94	99	7	2	87	96
Heroin	38	67	*	4	38	61
Hallucinogens	321 ^b	435	11	7	309 ^b	423
LSD	32	44	1	4	31	38
PCP	4	18	1	4	4	14
Ecstasy	206 ^a	291	*	7	196 ^b	280
Inhalants	142	144	6	6	136	137
Nonmedical Use of Psychotherapeutics ^{3,4}	2,222	2,263	27	27	2,172	2,212
Pain Relievers	1,608	1,653	17	21	1,571	1,609
OxyContin [®]	118 ^a	172	3	3	115 ^a	166
Tranquilizers	745	707	6	13	731	688
Stimulants ⁴	402	457	8	7	391	442
Methamphetamine ⁴	131	144	*	5	129	137
Sedatives	95	74	*	4	95	69
ILLCIT DRUGS OTHER THAN MARIJUANA²	2,784	2,850	51	37	2,708	2,786

*Low precision; no estimate reported.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.71B Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2007-2008	2009-2010	2007-2008	2009-2010	2007-2008	2009-2010
ILLCIT DRUGS²	9.6 ^b	10.6	5.1	4.4	9.8 ^b	10.9
Marijuana and Hashish	7.0 ^b	8.0	3.8	3.7	7.1 ^b	8.2
Cocaine	0.8 ^a	0.6	0.4	0.1	0.8 ^a	0.6
Crack	0.2	0.2	0.3	0.1	0.1	0.2
Heroin	0.1	0.1	*	0.2	0.1	0.1
Hallucinogens	0.5 ^b	0.7	0.4	0.3	0.5 ^b	0.7
LSD	0.1	0.1	0.0	0.2	0.1	0.1
PCP	0.0	0.0	0.0	0.2	0.0	0.0
Ecstasy	0.3 ^a	0.5	*	0.3	0.3 ^b	0.5
Inhalants	0.2	0.2	0.2	0.2	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{3,4}	3.6	3.7	1.0	1.1	3.7	3.8
Pain Relievers	2.6	2.7	0.7	0.9	2.7	2.7
OxyContin [®]	0.2 ^a	0.3	0.1	0.1	0.2 ^a	0.3
Tranquilizers	1.2	1.2	0.2	0.5	1.2	1.2
Stimulants ⁴	0.7	0.7	0.3	0.3	0.7	0.8
Methamphetamine ⁴	0.2	0.2	*	0.2	0.2	0.2
Sedatives	0.2	0.1	*	0.2	0.2	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	4.5	4.6	2.0	1.5	4.6	4.8

*Low precision; no estimate reported.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.72A Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2007-2008	2009-2010	2007-2008	2009-2010	2007-2008	2009-2010
TOTAL	5,955 ^b	6,537	130	109	5,777 ^b	6,383
AGE						
15-17	836	845	19	12	801	820
18-25	2,565 ^b	2,803	67	69	2,476 ^b	2,717
26-44	2,554 ^a	2,888	45	27	2,499 ^a	2,846
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5,308 ^a	5,658	117	88	5,154 ^a	5,531
White	4,167	4,402	76	59	4,071	4,321
Black or African American	874	976	36	25	825	935
American Indian or Alaska Native	35	45	*	*	34	42
Native Hawaiian or Other Pacific Islander	*	11	*	*	*	11
Asian	106	100	*	*	103	100
Two or More Races	100	125	*	*	96	122
Hispanic or Latino	647 ^b	879	13	21	623 ^b	853
TRIMESTER²						
First	N/A	N/A	56	59	N/A	N/A
Second	N/A	N/A	46	32	N/A	N/A
Third	N/A	N/A	24	16	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.72B Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2007-2008	2009-2010	2007-2008	2009-2010	2007-2008	2009-2010
TOTAL	9.6 ^b	10.6	5.1	4.4	9.8 ^b	10.9
AGE						
15-17	13.1	13.7	21.6	16.2	12.9	13.6
18-25	15.7 ^a	16.7	7.1	7.4	16.2 ^a	17.3
26-44	6.5 ^a	7.5	3.0	1.9	6.7 ^a	7.7
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	10.3 ^a	11.2	5.7	4.4	10.5 ^b	11.4
White	10.9 ^a	11.8	5.3	3.9	11.1 ^a	12.2
Black or African American	10.1	11.2	9.0	8.2	10.1	11.2
American Indian or Alaska Native	12.2	13.4	*	*	12.2	13.1
Native Hawaiian or Other Pacific Islander	*	4.3	*	*	*	4.2
Asian	3.2	3.1	*	*	3.3	3.2
Two or More Races	13.9	15.0	*	*	14.2	15.4
Hispanic or Latino	6.2 ^b	8.1	2.6	4.7	6.3 ^b	8.3
TRIMESTER²						
First	N/A	N/A	7.2	8.1	N/A	N/A
Second	N/A	N/A	5.0	3.5	N/A	N/A
Third	N/A	N/A	2.8	2.1	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.73A Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2007-2008	2009-2010	2007-2008	2009-2010	2007-2008	2009-2010
TOTAL	4,316 ^b	4,904	97	90	4,181 ^b	4,785
AGE						
15-17	624	646	15	11	597	626
18-25	2,025 ^b	2,294	51	61	1,956 ^b	2,220
26-44	1,667 ^a	1,964	31	18	1,628 ^a	1,939
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	3,875 ^b	4,312	91	72	3,755 ^b	4,214
White	3,005 ^a	3,326	55	48	2,935 ^a	3,262
Black or African American	674	779	33	21	629	750
American Indian or Alaska Native	28	32	*	*	27	30
Native Hawaiian or Other Pacific Islander	12	10	*	*	12	9
Asian	68	56	*	*	68	56
Two or More Races	88	109	*	*	84	107
Hispanic or Latino	441 ^b	592	6 ^a	18	426 ^b	571
TRIMESTER²						
First	N/A	N/A	51	49	N/A	N/A
Second	N/A	N/A	34	26	N/A	N/A
Third	N/A	N/A	11	14	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.73B Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2007-2008	2009-2010	2007-2008	2009-2010	2007-2008	2009-2010
TOTAL	7.0 ^b	8.0	3.8	3.7	7.1 ^b	8.2
AGE						
15-17	9.8	10.5	17.1	14.8	9.6	10.4
18-25	12.4 ^b	13.7	5.4	6.6	12.8 ^b	14.1
26-44	4.3 ^a	5.1	2.1	1.2	4.4 ^b	5.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	7.6 ^b	8.5	4.4	3.6	7.7 ^b	8.7
White	7.9 ^b	8.9	3.8	3.2	8.0 ^b	9.2
Black or African American	7.8	9.0	8.3	7.0	7.7	9.0
American Indian or Alaska Native	9.8	9.5	*	*	9.7	9.4
Native Hawaiian or Other Pacific Islander	4.3	3.7	*	*	4.4	3.6
Asian	2.1	1.7	*	*	2.2	1.8
Two or More Races	12.2	13.0	*	*	12.5	13.5
Hispanic or Latino	4.2 ^b	5.5	1.3 ^a	4.1	4.3 ^a	5.5
TRIMESTER²						
First	N/A	N/A	6.6	6.7	N/A	N/A
Second	N/A	N/A	3.7	2.9	N/A	N/A
Third	N/A	N/A	1.3	1.8	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.74A Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2007-2008	2009-2010	2007-2008	2009-2010	2007-2008	2009-2010
TOBACCO PRODUCTS²	17,381	16,959	427	408	16,836	16,468
Cigarettes	16,619	16,138	418	399	16,084	15,664
Smokeless Tobacco	235 ^a	319	3	8	226 ^a	308
Cigars	1,890 ^a	2,103	29	34	1,847 ^a	2,058
Pipe Tobacco	223 ^a	281	3	6	219	269
ALCOHOL	32,205	32,520	272	264	31,786	32,095
Binge Alcohol Use ³	14,444	14,566	116	90	14,256	14,409
Heavy Alcohol Use ³	3,262	3,228	21	23	3,220	3,193

*Low precision; no estimate reported.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.74B Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2007-2008	2009-2010	2007-2008	2009-2010	2007-2008	2009-2010
TOBACCO PRODUCTS²	28.1	27.6	16.7	16.6	28.6	28.1
Cigarettes	26.9	26.3	16.4	16.3	27.3	26.7
Smokeless Tobacco	0.4 ^a	0.5	0.1	0.3	0.4 ^a	0.5
Cigars	3.1 ^a	3.4	1.1	1.4	3.1 ^a	3.5
Pipe Tobacco	0.4 ^a	0.5	0.1	0.2	0.4	0.5
ALCOHOL	52.1	53.0	10.6	10.8	54.0	54.7
Binge Alcohol Use ³	23.4	23.7	4.5	3.7	24.2	24.6
Heavy Alcohol Use ³	5.3	5.3	0.8	1.0	5.5	5.4

*Low precision; no estimate reported.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.75A Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2007-2008	2009-2010	2007-2008	2009-2010	2007-2008	2009-2010
TOTAL	16,619	16,138	418	399	16,084	15,664
AGE						
15-17	948 ^b	837	18	17	914 ^b	805
18-25	5,186	5,146	208	212	4,935	4,903
26-44	10,485	10,155	192	170	10,236	9,956
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	14,896 ^a	14,166	390	373	14,415 ^a	13,725
White	12,287 ^a	11,605	307	314	11,932 ^a	11,234
Black or African American	1,814	1,799	69	41	1,720	1,749
American Indian or Alaska Native	143	133	*	*	139	129
Native Hawaiian or Other Pacific Islander	*	34	*	*	*	30
Asian	340	317	*	*	324	317
Two or More Races	219	278	*	*	207	265
Hispanic or Latino	1,723	1,971	28	26	1,669 ^a	1,940
TRIMESTER²						
First	N/A	N/A	164	177	N/A	N/A
Second	N/A	N/A	138	124	N/A	N/A
Third	N/A	N/A	107	96	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.75B Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2007-2008	2009-2010	2007-2008	2009-2010	2007-2008	2009-2010
TOTAL	26.9	26.3	16.4	16.3	27.3	26.7
AGE						
15-17	14.9 ^a	13.6	20.6	22.7	14.7 ^a	13.4
18-25	31.8	30.7	22.1	22.7	32.3	31.2
26-44	26.8	26.4	12.6	11.8	27.4	27.0
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	29.0	28.0	19.1	18.6	29.4	28.4
White	32.3	31.2	21.4	20.8	32.7	31.6
Black or African American	21.1	20.7	17.2	13.5	21.1	21.0
American Indian or Alaska Native	49.2	40.1	*	*	49.6	40.6
Native Hawaiian or Other Pacific Islander	*	13.2	*	*	*	11.9
Asian	10.3	9.7	*	*	10.4	10.2
Two or More Races	30.4	33.2	*	*	30.8	33.5
Hispanic or Latino	16.4	18.2	5.5	5.8	16.9	18.8
TRIMESTER²						
First	N/A	N/A	21.2	24.2	N/A	N/A
Second	N/A	N/A	15.2	13.5	N/A	N/A
Third	N/A	N/A	12.5	12.2	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.76A Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2007-2008	2009-2010	2007-2008	2009-2010	2007-2008	2009-2010
TOTAL	32,205	32,520	272	264	31,786	32,095
AGE						
15-17	1,516 ^b	1,350	14	10	1,487 ^b	1,328
18-25	9,383 ^a	9,603	102	89	9,229	9,468
26-44	21,306	21,567	155	166	21,070	21,299
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	28,091	28,088	239	220	27,735	27,734
White	22,482	22,152	158	168	22,257	21,885
Black or African American	3,849	4,048	77 ^a	38	3,748	3,977
American Indian or Alaska Native	150	140	*	*	149	140
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1,105	1,194	*	*	1,086	1,184
Two or More Races	383	447	*	*	373	443
Hispanic or Latino	4,114	4,432	33	44	4,051	4,361
TRIMESTER²						
First	N/A	N/A	160	146	N/A	N/A
Second	N/A	N/A	71	87	N/A	N/A
Third	N/A	N/A	30	25	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.76B Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2007-2008	2009-2010	2007-2008	2009-2010	2007-2008	2009-2010
TOTAL	52.1	53.0	10.6	10.8	54.0	54.7
AGE						
15-17	23.8 ^a	21.9	16.0	12.8	23.9 ^a	22.0
18-25	57.5	57.4	10.9	9.5	60.4	60.2
26-44	54.5	56.0	10.1	11.5	56.4	57.8
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	54.8	55.5	11.7	11.0	56.6	57.4
White	59.0	59.6	11.0	11.2	60.9	61.6
Black or African American	44.7	46.6	19.1	12.7	46.0	47.7
American Indian or Alaska Native	51.6	42.2	*	*	53.3	44.0
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	33.4	36.6	*	*	35.0	38.1
Two or More Races	53.2	53.5	*	*	55.4	56.0
Hispanic or Latino	39.3	40.9	6.4	9.9	41.0	42.3
TRIMESTER²						
First	N/A	N/A	20.7	19.9	N/A	N/A
Second	N/A	N/A	7.8	9.6	N/A	N/A
Third	N/A	N/A	3.5	3.2	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.77A Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2007-2008	2009-2010	2007-2008	2009-2010	2007-2008	2009-2010
TOTAL	14,444	14,566	116	90	14,256	14,409
AGE						
15-17	892 ^b	778	11	8	871 ^b	761
18-25	5,485	5,541	55	40	5,403	5,465
26-44	8,067	8,247	50	42	7,982	8,183
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	12,388	12,305	99	71	12,231	12,185
White	10,217	10,019	56	56	10,132	9,931
Black or African American	1,485	1,595	40	14	1,430	1,565
American Indian or Alaska Native	100	88	*	*	98	88
Native Hawaiian or Other Pacific Islander	*	44	*	*	*	44
Asian	335	352	*	*	323	352
Two or More Races	167	206	*	*	164	205
Hispanic or Latino	2,056	2,261	17	19	2,025	2,225
TRIMESTER²						
First	N/A	N/A	79	74	N/A	N/A
Second	N/A	N/A	17	12	N/A	N/A
Third	N/A	N/A	11 ^a	2	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.77B Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2007-2008	2009-2010	2007-2008	2009-2010	2007-2008	2009-2010
TOTAL	23.4	23.7	4.5	3.7	24.2	24.6
AGE						
15-17	14.0 ^a	12.6	12.3	10.3	14.0 ^a	12.6
18-25	33.6	33.1	5.9	4.3	35.4	34.8
26-44	20.6	21.4	3.3	2.9	21.4	22.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	24.1	24.3	4.8	3.6	25.0	25.2
White	26.8	26.9	3.9	3.7	27.7	28.0
Black or African American	17.2	18.3	10.0	4.7	17.6	18.8
American Indian or Alaska Native	34.2	26.3	*	*	35.2	27.6
Native Hawaiian or Other Pacific Islander	*	17.2	*	*	*	17.4
Asian	10.1	10.8	*	*	10.4	11.3
Two or More Races	23.3	24.7	*	*	24.3	25.9
Hispanic or Latino	19.6	20.8	3.3	4.3	20.5	21.6
TRIMESTER²						
First	N/A	N/A	10.3	10.1	N/A	N/A
Second	N/A	N/A	1.9	1.3	N/A	N/A
Third	N/A	N/A	1.3 ^a	0.2	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.78A Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2007-2008	2009-2010	2007-2008	2009-2010	2007-2008	2009-2010
TOTAL	3,262	3,228	21	23	3,220	3,193
AGE						
15-17	196	171	3	4	190	164
18-25	1,506	1,424	10	14	1,489	1,401
26-44	1,559	1,633	9	5	1,541	1,628
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2,922	2,871	20	17	2,884	2,845
White	2,563	2,465	16	13	2,538	2,445
Black or African American	249	290	3	4	237	285
American Indian or Alaska Native	21	23	*	*	21	23
Native Hawaiian or Other Pacific Islander	7	9	*	*	7	9
Asian	41	45	*	*	41	45
Two or More Races	40	39	*	*	39	38
Hispanic or Latino	340	357	1	6	336	348
TRIMESTER²						
First	N/A	N/A	18	20	N/A	N/A
Second	N/A	N/A	1	4	N/A	N/A
Third	N/A	N/A	3	*	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.78B Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2007-2008 and 2009-2010

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2007-2008	2009-2010	2007-2008	2009-2010	2007-2008	2009-2010
TOTAL	5.3	5.3	0.8	1.0	5.5	5.4
AGE						
15-17	3.1	2.8	3.0	5.5	3.1	2.7
18-25	9.2 ^a	8.5	1.1	1.5	9.8 ^a	8.9
26-44	4.0	4.2	0.6	0.4	4.1	4.4
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5.7	5.7	1.0	0.9	5.9	5.9
White	6.7	6.6	1.1	0.8	6.9	6.9
Black or African American	2.9	3.3	0.8	1.4	2.9	3.4
American Indian or Alaska Native	7.4	7.0	*	*	7.6	7.4
Native Hawaiian or Other Pacific Islander	2.8	3.6	*	*	2.8	3.6
Asian	1.2	1.4	*	*	1.3	1.4
Two or More Races	5.5	4.6	*	*	5.8	4.8
Hispanic or Latino	3.2	3.3	0.2	1.4	3.4	3.4
TRIMESTER²						
First	N/A	N/A	2.3	2.7	N/A	N/A
Second	N/A	N/A	0.1	0.4	N/A	N/A
Third	N/A	N/A	0.3	*	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

^a Difference between estimate and 2009-2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2009-2010 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.