

**Table 6.11C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2009 and 2010**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLICIT DRUGS<sup>2</sup></b>	230	218	269	246	196	247	191	227
Marijuana and Hashish	200	198	231	202	166	219	150	171
Cocaine	92	70	72	63	47	42	32	37
Crack	46	30	42	33	35	32	30	26
Heroin	21	17	14	13	19	27	20	26
Hallucinogens	63	62	51	40	34	22	25	26
LSD	17	14	11	11	4	7	6	4
PCP	15	3	9	4	2	*	*	20
Ecstasy	54	46	35	33	30	17	13	13
Inhalants	21	25	18	28	36	70	34	23
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	109	104	147	134	101	132	127	146
Pain Relievers	96	94	119	101	79	112	115	113
OxyContin <sup>®</sup>	26	28	38	33	9	24	38	48
Tranquilizers	65	59	81	75	63	69	59	72
Stimulants <sup>4</sup>	48	49	68	45	39	32	40	42
Methamphetamine <sup>4</sup>	36	21	42	24	28	26	29	31
Sedatives	28	23	29	54	19	17	32	40
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	149	139	167	150	121	141	133	152

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.11D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2009 and 2010**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>2</sup></b>	1.14	1.10	0.57	0.56	0.27	0.33	0.16	0.18
Marijuana and Hashish	1.03	1.01	0.50	0.47	0.23	0.29	0.12	0.14
Cocaine	0.52	0.40	0.17	0.15	0.07	0.06	0.03	0.03
Crack	0.27	0.17	0.10	0.08	0.05	0.04	0.03	0.02
Heroin	0.12	0.10	0.03	0.03	0.03	0.04	0.02	0.02
Hallucinogens	0.36	0.36	0.12	0.10	0.05	0.03	0.02	0.02
LSD	0.10	0.08	0.03	0.03	0.01	0.01	0.00	0.00
PCP	0.09	0.02	0.02	0.01	0.00	*	*	0.02
Ecstasy	0.31	0.27	0.08	0.08	0.04	0.02	0.01	0.01
Inhalants	0.12	0.14	0.04	0.07	0.05	0.10	0.03	0.02
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	0.61	0.60	0.33	0.32	0.14	0.18	0.10	0.12
Pain Relievers	0.54	0.54	0.27	0.24	0.11	0.15	0.09	0.09
OxyContin <sup>®</sup>	0.15	0.17	0.09	0.08	0.01	0.03	0.03	0.04
Tranquilizers	0.37	0.35	0.19	0.18	0.09	0.09	0.05	0.06
Stimulants <sup>4</sup>	0.28	0.28	0.16	0.11	0.05	0.04	0.03	0.03
Methamphetamine <sup>4</sup>	0.21	0.13	0.10	0.06	0.04	0.04	0.02	0.03
Sedatives	0.16	0.13	0.07	0.13	0.03	0.02	0.03	0.03
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	0.78	0.76	0.37	0.36	0.17	0.19	0.11	0.12

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.11P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLICIT DRUGS<sup>2</sup></b>	0.3640		0.7994		0.0067		0.2079	
Marijuana and Hashish	0.7053		0.5657		0.0107		0.1496	
Cocaine	0.1092		0.9569		0.5227		0.1650	
Crack	0.2742		0.6980		0.4675		0.9395	
Heroin	0.8004		0.7483		0.2803		0.7754	
Hallucinogens	0.9814		0.6424		0.6682		0.7760	
LSD	0.6816		0.6748		0.3747		0.3089	
PCP	0.0477		0.6352		*		*	
Ecstasy	0.6998		0.5400		0.7059		0.5506	
Inhalants	0.9486		0.0820		0.3224		0.8011	
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	0.2148		0.4560		0.0618		0.8534	
Pain Relievers	0.4517		0.2496		0.0652		0.4603	
OxyContin <sup>®</sup>	0.6831		0.8738		0.0152		0.9789	
Tranquilizers	0.0867		0.5709		0.1028		0.3679	
Stimulants <sup>4</sup>	0.2355		0.1310		0.4765		0.5756	
Methamphetamine <sup>4</sup>	0.0296		0.2623		0.6855		0.7715	
Sedatives	0.3981		0.6837		0.3106		0.4638	
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	0.1401		0.5313		0.1825		0.8506	

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.12C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2009 and 2010**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLICIT DRUGS<sup>2</sup></b>	25	24	37	41	25	25	47	48
Marijuana and Hashish	24	23	35	39	21	24	35	35
Cocaine	7	7	8	7	3	5	6	3
Crack	4	2	2	0	*	*	*	*
Heroin	3	1	1	2	*	1	3	2
Hallucinogens	11	10	12	13	5	8	10	10
LSD	5	7	2	5	2	3	3	3
PCP	4	1	2	1	1	*	1	2
Ecstasy	8	6	11	11	3	7	7	7
Inhalants	7	8	9	9	9	8	17	18
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	13	12	16	21	13	10	28	28
Pain Relievers	12	12	15	19	12	9	26	25
OxyContin <sup>®</sup>	7	6	5	6	4	2	5	6
Tranquilizers	9	8	8	8	6	4	6	10
Stimulants <sup>4</sup>	7	7	7	10	6	3	9	7
Methamphetamine <sup>4</sup>	4	3	4	6	3	1	5	3
Sedatives	3	3	3	4	2	*	8	6
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	17	16	22	25	15	14	35	34

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.12D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2009 and 2010**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>2</sup></b>	2.58	2.76	1.66	1.85	1.52	1.54	0.21	0.22
Marijuana and Hashish	2.65	3.05	1.64	1.80	1.31	1.49	0.16	0.17
Cocaine	1.40	1.69	0.47	0.45	0.21	0.34	0.03	0.01
Crack	0.71	0.45	0.10	0.02	*	*	*	*
Heroin	0.56	0.32	0.06	0.12	*	0.08	0.02	0.01
Hallucinogens	1.98	2.35	0.74	0.82	0.36	0.54	0.05	0.05
LSD	1.02	1.70	0.13	0.34	0.14	0.22	0.02	0.01
PCP	0.82	0.32	0.09	0.08	0.06	*	0.01	0.01
Ecstasy	1.46	1.39	0.65	0.73	0.24	0.48	0.03	0.03
Inhalants	1.25	1.78	0.54	0.58	0.60	0.59	0.08	0.09
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	2.26	2.72	0.92	1.28	0.87	0.69	0.13	0.13
Pain Relievers	2.10	2.66	0.87	1.21	0.78	0.62	0.12	0.12
OxyContin <sup>®</sup>	1.22	1.46	0.32	0.38	0.28	0.12	0.02	0.03
Tranquilizers	1.55	1.83	0.49	0.56	0.39	0.26	0.03	0.05
Stimulants <sup>4</sup>	1.41	1.72	0.41	0.67	0.39	0.22	0.04	0.03
Methamphetamine <sup>4</sup>	0.78	0.83	0.23	0.42	0.19	0.10	0.02	0.01
Sedatives	0.63	0.79	0.17	0.27	0.12	*	0.04	0.03
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	2.66	3.15	1.23	1.51	1.01	0.96	0.16	0.16

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.12P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLICIT DRUGS<sup>2</sup></b>	0.8435		0.0043		0.1244		0.8381	
Marijuana and Hashish	0.7737		0.0023		0.0465		0.4741	
Cocaine	0.6027		0.4275		0.2427		0.4579	
Crack	0.7407		0.1637		*		*	
Heroin	0.2178		0.4293		*		0.4999	
Hallucinogens	0.6811		0.5351		0.1192		0.8675	
LSD	0.7043		0.0701		0.2060		0.6141	
PCP	0.2351		0.7668		*		0.5400	
Ecstasy	0.5581		0.4797		0.0674		0.9303	
Inhalants	0.8314		0.8262		0.9361		0.2112	
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	0.5537		0.0559		0.0916		0.7371	
Pain Relievers	0.9961		0.0775		0.0937		0.5410	
OxyContin <sup>®</sup>	0.2058		0.9788		0.2984		0.9559	
Tranquilizers	0.4424		0.9629		0.1350		0.0762	
Stimulants <sup>4</sup>	0.5253		0.4589		0.5401		0.4426	
Methamphetamine <sup>4</sup>	0.5748		0.7715		0.6684		0.2672	
Sedatives	0.9790		0.3513		*		0.6133	
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	0.5206		0.0550		0.8220		0.9886	

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.13C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2009 and 2010**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>2</sup></b>	99	94	97	94	55	58	51	58
Marijuana and Hashish	92	88	90	85	53	54	42	54
Cocaine	29	30	19	23	8	10	5	7
Crack	5	8	8	12	5	5	2	2
Heroin	6	10	7	12	4	6	7	3
Hallucinogens	31	35	25	24	14	16	12	10
LSD	12	11	9	10	3	6	5	3
PCP	2	2	*	2	*	*	1	*
Ecstasy	20	26	21	19	12	15	10	8
Inhalants	9	10	13	16	8	7	10	6
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	46	40	45	45	21	28	31	28
Pain Relievers	38	34	37	39	18	25	28	24
OxyContin <sup>®</sup>	10	11	14	18	4	6	6	12
Tranquilizers	26	25	23	23	12	16	13	10
Stimulants <sup>4</sup>	23	23	18	20	10	9	11	13
Methamphetamine <sup>4</sup>	10	5	10	14	6	6	2	4
Sedatives	8	7	9	5	7	3	9	8
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	60	57	54	50	26	32	33	30

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.13D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2009 and 2010**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>2</sup></b>	1.22	1.23	0.82	0.80	0.72	0.73	0.38	0.41
Marijuana and Hashish	1.24	1.24	0.78	0.77	0.70	0.70	0.32	0.39
Cocaine	0.60	0.61	0.20	0.25	0.12	0.14	0.04	0.06
Crack	0.11	0.18	0.09	0.13	0.07	0.07	0.02	0.02
Heroin	0.12	0.21	0.07	0.13	0.05	0.09	0.05	0.03
Hallucinogens	0.62	0.70	0.26	0.26	0.20	0.23	0.10	0.07
LSD	0.26	0.24	0.09	0.11	0.05	0.08	0.04	0.02
PCP	0.05	0.05	*	0.02	*	*	0.01	*
Ecstasy	0.43	0.53	0.22	0.21	0.18	0.21	0.08	0.06
Inhalants	0.19	0.22	0.14	0.17	0.11	0.10	0.08	0.04
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>								
Pain Relievers	0.83	0.76	0.46	0.46	0.30	0.39	0.24	0.21
OxyContin <sup>®</sup>	0.75	0.67	0.39	0.40	0.26	0.35	0.22	0.18
Tranquilizers	0.22	0.23	0.15	0.20	0.05	0.09	0.05	0.09
Stimulants <sup>4</sup>	0.53	0.51	0.24	0.25	0.17	0.22	0.10	0.08
Methamphetamine <sup>4</sup>	0.47	0.45	0.19	0.22	0.14	0.13	0.09	0.10
Sedatives	0.22	0.11	0.10	0.15	0.09	0.08	0.01	0.03
Sedatives	0.18	0.15	0.10	0.05	0.10	0.04	0.07	0.06
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	0.98	0.99	0.53	0.50	0.36	0.44	0.25	0.22

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.13P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>2</sup></b>	0.5553		0.2790		0.0090		0.0372	
Marijuana and Hashish	0.8106		0.1338		0.0067		0.0073	
Cocaine	0.7113		0.2601		0.6258		0.2884	
Crack	0.1191		0.2600		0.9016		0.8611	
Heroin	0.3770		0.1434		0.3955		0.1107	
Hallucinogens	0.4739		0.9140		0.2726		0.8194	
LSD	0.8917		0.5775		0.5499		0.3657	
PCP	0.8080		*		*		*	
Ecstasy	0.6057		0.6090		0.2591		0.7416	
Inhalants	0.6535		0.8126		0.9266		0.2472	
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	0.0048		0.5070		0.2115		0.8107	
Pain Relievers	0.0114		0.7628		0.1194		0.7676	
OxyContin <sup>®</sup>	0.7156		0.2885		0.0731		0.0195	
Tranquilizers	0.0600		0.7902		0.2311		0.1779	
Stimulants <sup>4</sup>	0.0614		0.3050		0.4573		0.0920	
Methamphetamine <sup>4</sup>	0.0538		0.3916		0.9904		0.1421	
Sedatives	0.1833		0.4910		0.1166		0.5970	
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	0.1889		0.3226		0.1675		0.9818	

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.14C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2009 and 2010**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>2</sup></b>	202	189	246	222	186	234	175	213
Marijuana and Hashish	174	173	208	180	159	205	135	156
Cocaine	85	61	68	59	46	40	31	36
Crack	46	28	41	31	35	32	30	26
Heroin	20	14	12	5	19	26	18	26
Hallucinogens	53	48	43	29	31	11	20	22
LSD	11	3	7	*	*	*	*	*
PCP	14	*	9	3	2	*	*	20
Ecstasy	48	37	25	24	27	*	*	7
Inhalants	17	22	7	22	34	69	28	13
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	95	93	141	124	97	127	120	140
Pain Relievers	86	84	114	91	75	108	109	106
OxyContin <sup>®</sup>	23	25	34	27	7	24	37	46
Tranquilizers	57	50	76	71	61	67	58	71
Stimulants <sup>4</sup>	41	41	64	39	37	30	36	39
Methamphetamine <sup>4</sup>	34	20	40	18	27	26	28	30
Sedatives	27	21	28	54	18	16	29	39
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	131	119	155	139	115	135	124	144

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.14D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2009 and 2010**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>2</sup></b>	1.48	1.39	0.70	0.68	0.29	0.35	0.20	0.24
Marijuana and Hashish	1.32	1.29	0.61	0.56	0.25	0.31	0.15	0.17
Cocaine	0.69	0.50	0.21	0.19	0.07	0.06	0.04	0.04
Crack	0.38	0.24	0.13	0.10	0.06	0.05	0.03	0.03
Heroin	0.17	0.12	0.04	0.02	0.03	0.04	0.02	0.03
Hallucinogens	0.43	0.40	0.14	0.09	0.05	0.02	0.02	0.02
LSD	0.09	0.02	0.02	*	*	*	*	*
PCP	0.12	*	0.03	0.01	0.00	*	*	0.02
Ecstasy	0.39	0.31	0.08	0.08	0.04	*	*	0.01
Inhalants	0.14	0.18	0.02	0.07	0.05	0.11	0.03	0.01
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	0.76	0.77	0.43	0.40	0.15	0.20	0.14	0.16
Pain Relievers	0.70	0.70	0.35	0.29	0.12	0.17	0.12	0.12
OxyContin <sup>®</sup>	0.19	0.21	0.11	0.09	0.01	0.04	0.04	0.05
Tranquilizers	0.47	0.42	0.24	0.23	0.10	0.10	0.07	0.08
Stimulants <sup>4</sup>	0.34	0.35	0.20	0.13	0.06	0.05	0.04	0.04
Methamphetamine <sup>4</sup>	0.28	0.17	0.13	0.06	0.04	0.04	0.03	0.03
Sedatives	0.22	0.18	0.09	0.17	0.03	0.03	0.03	0.04
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	1.01	0.95	0.47	0.44	0.18	0.21	0.14	0.16

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.14P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLICIT DRUGS<sup>2</sup></b>	0.4849		0.8955		0.0451		0.3987	
Marijuana and Hashish	0.8459		0.7288		0.0983		0.3776	
Cocaine	0.0962		0.7448		0.3876		0.1849	
Crack	0.1719		0.4925		0.4512		0.9279	
Heroin	0.6091		0.2662		0.3852		0.4518	
Hallucinogens	0.7848		0.4538		0.0929		0.7072	
LSD	0.2930		*		*		*	
PCP	*		0.5221		*		*	
Ecstasy	0.5748		0.5186		*		*	
Inhalants	0.8083		0.0087		0.2898		0.3942	
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	0.8687		0.4398		0.0678		0.8409	
Pain Relievers	0.8541		0.1786		0.0840		0.5564	
OxyContin <sup>®</sup>	0.9133		0.7244		0.0308		0.6297	
Tranquilizers	0.3375		0.5914		0.1266		0.3557	
Stimulants <sup>4</sup>	0.8127		0.0353		0.6495		0.8237	
Methamphetamine <sup>4</sup>	0.1034		0.1055		0.7009		0.7848	
Sedatives	0.6459		0.6440		0.6475		0.4563	
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	0.3263		0.5773		0.2609		0.8463	

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.15C Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Numbers in Thousands, 2009 and 2010**

<b>Drug</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-20 (2009)</b>	<b>Aged 12-20 (2010)</b>	<b>Aged 21+ (2009)</b>	<b>Aged 21+ (2010)</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	418	442	109	114	394	421
Marijuana or Hashish	360	377	104	108	338	356
Cocaine (Including Crack)	134	104	23	26	131	101
Heroin	32	34	8	12	31	32
Hallucinogens	88	79	36	35	80	68
Inhalants	45	80	20	17	40	78
Pain Relievers <sup>2</sup>	174	178	42	48	167	171
Tranquilizers <sup>2</sup>	121	118	26	24	117	115
Stimulants <sup>2,3</sup>	91	75	23	24	88	69
Methamphetamine <sup>3</sup>	61	42	9	15	61	39
Sedatives <sup>2</sup>	46	61	9	8	44	60
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	261	259	62	65	247	250

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Refers to nonmedical use; does not include over-the-counter drugs.

<sup>3</sup> Estimates in the designated rows include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.15D Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Percentages, 2009 and 2010**

<b>Drug</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-20 (2009)</b>	<b>Aged 12-20 (2010)</b>	<b>Aged 21+ (2009)</b>	<b>Aged 21+ (2010)</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	0.30	0.31	0.72	0.83	0.31	0.32
Marijuana or Hashish	0.27	0.27	0.72	0.82	0.27	0.28
Cocaine (Including Crack)	0.10	0.08	0.22	0.25	0.11	0.08
Heroin	0.02	0.03	0.08	0.12	0.03	0.03
Hallucinogens	0.07	0.06	0.33	0.34	0.07	0.06
Inhalants	0.03	0.06	0.19	0.17	0.03	0.06
Pain Relievers <sup>2</sup>	0.13	0.14	0.38	0.45	0.14	0.14
Tranquilizers <sup>2</sup>	0.09	0.09	0.24	0.24	0.10	0.09
Stimulants <sup>2,3</sup>	0.07	0.06	0.21	0.23	0.07	0.06
Methamphetamine <sup>3</sup>	0.05	0.03	0.09	0.14	0.05	0.03
Sedatives <sup>2</sup>	0.04	0.05	0.08	0.08	0.04	0.05
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.19	0.19	0.52	0.59	0.20	0.20

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Refers to nonmedical use; does not include over-the-counter drugs.

<sup>3</sup> Estimates in the designated rows include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.15P Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

<b>Drug</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-20 (2009)</b>	<b>Aged 12-20 (2010)</b>	<b>Aged 21+ (2009)</b>	<b>Aged 21+ (2010)</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	0.5911		0.1974		0.5979	
Marijuana or Hashish	0.6319		0.2131		0.6464	
Cocaine (Including Crack)	0.1366		0.7940		0.1193	
Heroin	0.4773		0.3340		0.6495	
Hallucinogens	0.5529		0.8642		0.5767	
Inhalants	0.1557		0.2572		0.0514	
Pain Relievers <sup>2</sup>	0.7842		0.4076		0.9694	
Tranquilizers <sup>2</sup>	0.7500		0.2316		0.5258	
Stimulants <sup>2,3</sup>	0.0292		0.9282		0.0298	
Methamphetamine <sup>3</sup>	0.0281		0.3125		0.0139	
Sedatives <sup>2</sup>	0.6482		0.6281		0.7017	
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.4977		0.8865		0.5950	

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Refers to nonmedical use; does not include over-the-counter drugs.

<sup>3</sup> Estimates in the designated rows include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.16C Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: Standard Errors of Numbers in Thousands, 2009 and 2010**

<b>Drug<sup>1</sup></b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-20 (2009)</b>	<b>Aged 12-20 (2010)</b>	<b>Aged 21+ (2009)</b>	<b>Aged 21+ (2010)</b>
<b>ILLCIT DRUGS<sup>2</sup></b>	271	269	72	71	253	256
Marijuana or Hashish	253	247	72	69	234	235
Cocaine (Including Crack)	54	49	7	11	53	47
Heroin	20	19	2	8	20	17
Hallucinogens	34	26	11	10	31	24
Inhalants	9	14	4	8	8	12
Pain Relievers <sup>3</sup>	73	65	13	14	72	63
Tranquilizers <sup>3</sup>	38	32	5	10	37	31
Stimulants <sup>3</sup>	15	27	4	4	14	27
Methamphetamine <sup>4</sup>	34	26	5	2	34	26
Sedatives <sup>3</sup>	10	11	*	3	10	10
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	112	98	18	24	110	95

\*Low precision; no estimate reported.

<sup>1</sup> Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Refers to nonmedical use; does not include over-the-counter drugs.

<sup>4</sup> Estimates in the designated row include data from new methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.16D Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: Standard Errors of Percentages, 2009 and 2010**

<b>Drug<sup>1</sup></b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-20 (2009)</b>	<b>Aged 12-20 (2010)</b>	<b>Aged 21+ (2009)</b>	<b>Aged 21+ (2010)</b>
<b>ILLCIT DRUGS<sup>2</sup></b>	0.20	0.20	0.62	0.62	0.21	0.20
Marijuana or Hashish	0.19	0.18	0.62	0.60	0.19	0.19
Cocaine (Including Crack)	0.04	0.04	0.07	0.11	0.04	0.04
Heroin	0.02	0.01	0.02	0.08	0.02	0.01
Hallucinogens	0.03	0.02	0.10	0.10	0.03	0.02
Inhalants	0.01	0.01	0.04	0.08	0.01	0.01
Pain Relievers <sup>3</sup>	0.06	0.05	0.13	0.15	0.06	0.05
Tranquilizers <sup>3</sup>	0.03	0.02	0.05	0.10	0.03	0.03
Stimulants <sup>3</sup>	0.01	0.02	0.04	0.04	0.01	0.02
Methamphetamine <sup>4</sup>	0.03	0.02	0.04	0.02	0.03	0.02
Sedatives <sup>3</sup>	0.01	0.01	*	0.03	0.01	0.01
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	0.09	0.07	0.17	0.24	0.09	0.08

\*Low precision; no estimate reported.

<sup>1</sup> Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Refers to nonmedical use; does not include over-the-counter drugs.

<sup>4</sup> Estimates in the designated row include data from new methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.16P Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

<b>Drug<sup>1</sup></b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-20 (2009)</b>	<b>Aged 12-20 (2010)</b>	<b>Aged 21+ (2009)</b>	<b>Aged 21+ (2010)</b>
<b>ILLICIT DRUGS<sup>2</sup></b>	0.4558		0.0871		0.6065	
Marijuana or Hashish	0.4450		0.2108		0.5169	
Cocaine (Including Crack)	0.2431		0.1735		0.1455	
Heroin	0.4632		0.1633		0.7321	
Hallucinogens	0.5797		0.8328		0.5137	
Inhalants	0.4794		0.5444		0.6257	
Pain Relievers <sup>3</sup>	0.8427		0.2213		0.9746	
Tranquilizers <sup>3</sup>	0.9850		0.0724		0.6714	
Stimulants <sup>3</sup>	0.1214		0.4571		0.1485	
Methamphetamine <sup>4</sup>	0.4257		0.9118		0.4288	
Sedatives <sup>3</sup>	0.7414		*		0.5847	
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	0.3841		0.0327		0.1858	

\*Low precision; no estimate reported.

<sup>1</sup> Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Refers to nonmedical use; does not include over-the-counter drugs.

<sup>4</sup> Estimates in the designated row include data from new methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.17C Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2009 and 2010**

<b>Drug</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	418	442	53	57	162	160	371	395
Marijuana or Hashish	360	377	49	54	149	148	318	333
Cocaine (Including Crack)	134	104	11	11	36	39	124	94
Heroin	32	34	3	2	9	16	31	30
Hallucinogens	88	79	17	19	45	48	74	57
Inhalants	45	80	14	14	18	20	39	76
Pain Relievers <sup>2</sup>	174	178	22	25	58	61	161	163
Tranquilizers <sup>2</sup>	121	118	13	11	39	38	111	109
Stimulants <sup>2,3</sup>	91	75	12	13	33	34	84	65
Methamphetamine <sup>3</sup>	61	42	6	7	15	16	59	38
Sedatives <sup>2</sup>	46	61	5	5	14	9	42	60
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	261	259	31	33	91	88	233	236

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Refers to nonmedical use; does not include over-the-counter drugs.

<sup>3</sup> Estimates in the designated rows include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.17D Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2009 and 2010**

<b>Drug</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
<b>ILLICIT DRUGS<sup>1</sup></b>	0.30	0.31	1.09	1.21	0.56	0.56	0.33	0.35
Marijuana or Hashish	0.27	0.27	1.06	1.17	0.54	0.55	0.29	0.30
Cocaine (Including Crack)	0.10	0.08	0.31	0.33	0.17	0.18	0.12	0.09
Heroin	0.02	0.03	0.08	0.07	0.05	0.08	0.03	0.03
Hallucinogens	0.07	0.06	0.46	0.54	0.21	0.22	0.07	0.05
Inhalants	0.03	0.06	0.39	0.42	0.09	0.10	0.04	0.07
Pain Relievers <sup>2</sup>	0.13	0.14	0.59	0.71	0.26	0.28	0.15	0.15
Tranquilizers <sup>2</sup>	0.09	0.09	0.37	0.33	0.18	0.18	0.10	0.10
Stimulants <sup>2,3</sup>	0.07	0.06	0.32	0.38	0.15	0.15	0.08	0.06
Methamphetamine <sup>3</sup>	0.05	0.03	0.17	0.22	0.07	0.08	0.06	0.04
Sedatives <sup>2</sup>	0.04	0.05	0.13	0.16	0.07	0.04	0.04	0.06
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.19	0.19	0.77	0.91	0.38	0.36	0.21	0.21

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Refers to nonmedical use; does not include over-the-counter drugs.

<sup>3</sup> Estimates in the designated rows include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.17P Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

<b>Drug</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	0.5911		0.0221		0.8802		0.5882	
Marijuana or Hashish	0.6319		0.0120		0.8481		0.6528	
Cocaine (Including Crack)	0.1366		0.9620		0.7065		0.0804	
Heroin	0.4773		0.6872		0.0664		0.9961	
Hallucinogens	0.5529		0.6314		0.4053		0.1974	
Inhalants	0.1557		0.8567		0.9698		0.0841	
Pain Relievers <sup>2</sup>	0.7842		0.8247		0.2475		0.8733	
Tranquilizers <sup>2</sup>	0.7500		0.1829		0.5039		0.4536	
Stimulants <sup>2,3</sup>	0.0292		0.7447		0.3218		0.0606	
Methamphetamine <sup>3</sup>	0.0281		0.6991		0.7339		0.0299	
Sedatives <sup>2</sup>	0.6482		0.8606		0.0498		0.9843	
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.4977		0.5629		0.2939		0.7270	

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Refers to nonmedical use; does not include over-the-counter drugs.

<sup>3</sup> Estimates in the designated rows include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.18C Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2009 and 2010**

<b>Drug<sup>1</sup></b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
<b>ILLCIT DRUGS<sup>2</sup></b>	271	269	34	38	101	104	238	237
Marijuana or Hashish	253	247	33	38	99	99	221	218
Cocaine (Including Crack)	54	49	4	4	16	20	51	43
Heroin	20	19	2	*	3	10	20	16
Hallucinogens	34	26	4	5	15	13	29	22
Inhalants	9	14	4	7	3	10	7	8
Pain Relievers <sup>3</sup>	73	65	6	8	19	21	70	61
Tranquilizers <sup>3</sup>	38	32	2	4	11	12	36	30
Stimulants <sup>3</sup>	15	27	1	1	6	15	14	23
Methamphetamine <sup>4</sup>	34	26	1	2	8	6	33	26
Sedatives <sup>3</sup>	10	11	0	3	5	2	8	10
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	112	98	9	12	32	36	106	90

\*Low precision; no estimate reported.

<sup>1</sup> Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Refers to nonmedical use; does not include over-the-counter drugs.

<sup>4</sup> Estimates in the designated row include data from new methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.18D Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2009 and 2010**

<b>Drug<sup>1</sup></b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
<b>ILLCIT DRUGS<sup>2</sup></b>	0.20	0.20	0.85	1.00	0.43	0.43	0.22	0.22
Marijuana or Hashish	0.19	0.18	0.84	0.98	0.43	0.41	0.20	0.20
Cocaine (Including Crack)	0.04	0.04	0.10	0.13	0.08	0.10	0.05	0.04
Heroin	0.02	0.01	0.05	*	0.01	0.05	0.02	0.01
Hallucinogens	0.03	0.02	0.11	0.14	0.07	0.06	0.03	0.02
Inhalants	0.01	0.01	0.11	0.21	0.02	0.05	0.01	0.01
Pain Relievers <sup>3</sup>	0.06	0.05	0.16	0.25	0.09	0.10	0.07	0.06
Tranquilizers <sup>3</sup>	0.03	0.02	0.04	0.12	0.05	0.06	0.03	0.03
Stimulants <sup>3</sup>	0.01	0.02	0.04	0.04	0.03	0.07	0.01	0.02
Methamphetamine <sup>4</sup>	0.03	0.02	0.04	0.05	0.04	0.03	0.03	0.02
Sedatives <sup>3</sup>	0.01	0.01	0.01	0.09	0.02	0.01	0.01	0.01
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	0.09	0.07	0.25	0.36	0.15	0.17	0.10	0.08

\*Low precision; no estimate reported.

<sup>1</sup> Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Refers to nonmedical use; does not include over-the-counter drugs.

<sup>4</sup> Estimates in the designated row include data from new methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.18P Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009**

<b>Drug<sup>1</sup></b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
<b>ILLCIT DRUGS<sup>2</sup></b>	0.4558		0.0084		0.2575		0.9027	
Marijuana or Hashish	0.4450		0.0051		0.3988		0.8198	
Cocaine (Including Crack)	0.2431		0.9347		0.4577		0.1182	
Heroin	0.4632		*		0.0148		0.9291	
Hallucinogens	0.5797		0.4272		0.6447		0.5946	
Inhalants	0.4794		0.5274		0.5231		0.9277	
Pain Relievers <sup>3</sup>	0.8427		0.1068		0.9556		0.9517	
Tranquilizers <sup>3</sup>	0.9850		0.0920		0.8661		0.9052	
Stimulants <sup>3</sup>	0.1214		0.5164		0.0678		0.5136	
Methamphetamine <sup>4</sup>	0.4257		0.7258		0.5278		0.4917	
Sedatives <sup>3</sup>	0.7414		0.3573		0.2296		0.9161	
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	0.3841		0.0981		0.8311		0.2374	

\*Low precision; no estimate reported.

<sup>1</sup> Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Refers to nonmedical use; does not include over-the-counter drugs.

<sup>4</sup> Estimates in the designated row include data from new methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.