

**Table 6.19C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2009 and 2010**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	311	303	138	168	260	292	133	146
Marijuana and Hashish	270	273	126	143	218	241	83	94
Cocaine	115	95	25	30	63	40	19	22
Crack	70	53	24	22	11	16	19	14
Heroin	30	34	20	17	12	*	*	20
Hallucinogens	74	71	14	13	49	31	21	23
LSD	21	17	5	4	4	9	4	*
PCP	17	3	1	1	3	*	*	20
Ecstasy	57	54	11	10	42	26	6	8
Inhalants	26	39	12	5	36	70	32	23
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	150	150	79	88	146	160	101	110
Pain Relievers	120	130	73	74	123	122	89	84
OxyContin <sup>®</sup>	42	43	38	48	17	22	6	5
Tranquilizers	89	88	48	52	84	76	43	50
Stimulants <sup>3</sup>	76	61	35	26	48	40	18	32
Methamphetamine <sup>3</sup>	52	33	28	20	30	26	5	23
Sedatives	38	32	25	14	25	52	20	38
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	195	184	79	93	170	170	109	114

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.19D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2009 and 2010**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	0.69	0.67	0.72	0.81	0.28	0.31	0.13	0.14
Marijuana and Hashish	0.62	0.62	0.65	0.70	0.24	0.26	0.08	0.09
Cocaine	0.29	0.24	0.13	0.15	0.07	0.04	0.02	0.02
Crack	0.18	0.13	0.12	0.11	0.01	0.02	0.02	0.01
Heroin	0.08	0.09	0.10	0.08	0.01	*	*	0.02
Hallucinogens	0.18	0.18	0.07	0.06	0.05	0.03	0.02	0.02
LSD	0.05	0.04	0.02	0.02	0.00	0.01	0.00	*
PCP	0.04	0.01	0.01	0.00	0.00	*	*	0.02
Ecstasy	0.14	0.14	0.06	0.05	0.05	0.03	0.01	0.01
Inhalants	0.07	0.10	0.06	0.02	0.04	0.08	0.03	0.02
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.37	0.37	0.41	0.44	0.16	0.17	0.10	0.11
Pain Relievers	0.29	0.33	0.38	0.37	0.13	0.13	0.09	0.08
OxyContin <sup>®</sup>	0.11	0.11	0.19	0.24	0.02	0.02	0.01	0.00
Tranquilizers	0.22	0.22	0.25	0.26	0.09	0.08	0.04	0.05
Stimulants <sup>3</sup>	0.19	0.16	0.18	0.13	0.05	0.04	0.02	0.03
Methamphetamine <sup>3</sup>	0.13	0.09	0.15	0.10	0.03	0.03	0.00	0.02
Sedatives	0.10	0.08	0.13	0.07	0.03	0.06	0.02	0.04
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.46	0.44	0.42	0.46	0.18	0.18	0.11	0.11

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.19P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	0.9433		0.5080		0.2138		0.2985	
Marijuana and Hashish	0.7563		0.5373		0.3422		0.1443	
Cocaine	0.3587		0.2420		0.2476		0.4772	
Crack	0.1516		0.8843		0.6208		0.9220	
Heroin	0.2491		0.6428		*		*	
Hallucinogens	0.7877		0.7430		0.1883		0.9113	
LSD	0.6254		0.5695		0.0533		*	
PCP	0.0359		0.9921		*		*	
Ecstasy	0.9458		0.5401		0.2027		0.9022	
Inhalants	0.4821		0.0519		0.2021		0.7086	
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.4506		0.9651		0.4853		0.8155	
Pain Relievers	0.6534		0.4183		0.7247		0.7303	
OxyContin <sup>®</sup>	0.4008		0.9918		0.8194		0.9318	
Tranquilizers	0.3718		0.8775		0.7638		0.2994	
Stimulants <sup>3</sup>	0.0139		0.6242		1.0000		0.1519	
Methamphetamine <sup>3</sup>	0.0112		0.6509		0.9613		0.2464	
Sedatives	0.2640		0.3543		0.6778		0.1491	
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.4032		0.9583		0.7963		0.8799	

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.20C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2009 and 2010**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	41	44	23	21	34	35	40	42
Marijuana and Hashish	39	41	21	18	30	32	27	29
Cocaine	9	11	4	2	7	2	3	2
Crack	4	2	*	*	*	0	*	*
Heroin	3	2	3	*	1	1	1	2
Hallucinogens	15	17	6	5	8	8	7	9
LSD	6	9	2	3	1	3	2	2
PCP	5	2	*	0	1	1	1	2
Ecstasy	13	13	2	3	6	7	6	6
Inhalants	8	11	5	4	12	9	16	18
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	21	22	10	12	12	16	25	25
Pain Relievers	19	20	9	10	12	15	24	22
OxyContin <sup>®</sup>	9	8	4	4	4	2	3	4
Tranquilizers	12	11	2	8	5	5	6	7
Stimulants <sup>3</sup>	11	11	5	2	4	6	7	7
Methamphetamine <sup>3</sup>	6	6	3	2	1	5	3	2
Sedatives	4	5	5	2	2	2	7	5
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	24	28	14	13	19	19	31	31

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.20D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2009 and 2010**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	1.77	2.02	2.17	2.19	1.26	1.38	0.19	0.20
Marijuana and Hashish	1.77	2.03	2.11	1.95	1.17	1.30	0.13	0.14
Cocaine	0.64	0.85	0.53	0.29	0.31	0.11	0.02	0.01
Crack	0.30	0.15	*	*	*	0.01	*	*
Heroin	0.22	0.17	0.38	*	0.03	0.06	0.01	0.01
Hallucinogens	1.03	1.30	0.74	0.55	0.34	0.37	0.04	0.04
LSD	0.44	0.70	0.24	0.31	0.05	0.14	0.01	0.01
PCP	0.33	0.12	*	0.05	0.04	0.05	0.01	0.01
Ecstasy	0.94	1.03	0.26	0.39	0.25	0.31	0.03	0.03
Inhalants	0.60	0.90	0.56	0.48	0.50	0.42	0.08	0.09
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	1.42	1.53	1.15	1.36	0.50	0.74	0.12	0.12
Pain Relievers	1.31	1.45	1.07	1.22	0.52	0.69	0.12	0.11
OxyContin <sup>®</sup>	0.64	0.66	0.48	0.54	0.16	0.10	0.02	0.02
Tranquilizers	0.89	0.86	0.25	0.93	0.23	0.25	0.03	0.03
Stimulants <sup>3</sup>	0.81	0.88	0.61	0.23	0.17	0.30	0.03	0.03
Methamphetamine <sup>3</sup>	0.46	0.46	0.39	0.21	0.03	0.23	0.02	0.01
Sedatives	0.32	0.40	0.55	0.29	0.08	0.09	0.03	0.03
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	1.53	1.89	1.54	1.50	0.80	0.86	0.15	0.15

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.20P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	0.5475		0.7007		0.0049		0.9025	
Marijuana and Hashish	0.5700		0.1643		0.0014		0.8686	
Cocaine	0.5276		0.6878		0.3070		0.5146	
Crack	0.2853		*		*		*	
Heroin	0.4414		*		0.4190		0.5225	
Hallucinogens	0.6578		1.0000		0.7871		0.7975	
LSD	0.2788		0.7879		0.1459		0.6867	
PCP	0.1262		*		0.8533		0.6433	
Ecstasy	0.6183		0.3202		0.3720		0.7496	
Inhalants	0.6745		0.9398		0.5137		0.1918	
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.3205		0.0995		0.1609		0.3098	
Pain Relievers	0.5944		0.1308		0.2344		0.2121	
OxyContin <sup>®</sup>	0.1957		0.7517		0.5189		0.7201	
Tranquilizers	0.1827		0.0442		0.8966		0.4561	
Stimulants <sup>3</sup>	0.4251		0.0407		0.3234		0.7890	
Methamphetamine <sup>3</sup>	0.3522		0.3192		0.3359		0.5725	
Sedatives	0.8077		0.3226		0.9240		0.9764	
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.7439		0.3186		0.5204		0.7237	

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.21C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2009 and 2010**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	129	124	39	45	78	85	32	38
Marijuana and Hashish	120	116	35	42	71	77	23	33
Cocaine	33	36	4	7	13	15	3	2
Crack	10	16	1	2	4	2	2	*
Heroin	9	16	6	3	1	0	2	1
Hallucinogens	36	42	12	8	24	20	4	5
LSD	15	14	4	3	4	8	3	1
PCP	*	3	1	*	2	*	*	*
Ecstasy	26	32	10	7	18	16	2	5
Inhalants	15	17	10	3	9	11	4	5
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	62	57	21	23	33	37	24	18
Pain Relievers	50	51	17	19	27	31	23	15
OxyContin <sup>®</sup>	17	21	4	11	6	5	4	3
Tranquilizers	33	35	10	9	18	16	8	4
Stimulants <sup>3</sup>	29	28	11	11	14	16	3	7
Methamphetamine <sup>3</sup>	15	16	0	4	2	4	1	1
Sedatives	13	8	*	3	6	5	2	7
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	76	73	23	24	42	43	24	19

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.21D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2009 and 2010**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	0.85	0.90	1.34	1.48	0.61	0.61	0.31	0.34
Marijuana and Hashish	0.85	0.90	1.25	1.42	0.56	0.57	0.22	0.31
Cocaine	0.33	0.37	0.15	0.27	0.12	0.13	0.03	0.02
Crack	0.10	0.17	0.04	0.09	0.03	0.02	0.02	*
Heroin	0.10	0.18	0.25	0.13	0.01	0.00	0.02	0.01
Hallucinogens	0.36	0.43	0.45	0.32	0.21	0.17	0.04	0.05
LSD	0.16	0.15	0.16	0.11	0.03	0.07	0.03	0.01
PCP	*	0.03	0.04	*	0.02	*	*	*
Ecstasy	0.27	0.34	0.38	0.27	0.16	0.14	0.02	0.04
Inhalants	0.15	0.18	0.38	0.12	0.08	0.10	0.03	0.04
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.54	0.57	0.78	0.85	0.29	0.29	0.23	0.17
Pain Relievers	0.47	0.52	0.64	0.74	0.24	0.25	0.22	0.14
OxyContin <sup>®</sup>	0.17	0.23	0.15	0.43	0.05	0.04	0.04	0.03
Tranquilizers	0.33	0.37	0.38	0.36	0.15	0.13	0.08	0.04
Stimulants <sup>3</sup>	0.29	0.30	0.42	0.42	0.13	0.13	0.03	0.07
Methamphetamine <sup>3</sup>	0.16	0.17	0.01	0.17	0.01	0.03	0.01	0.01
Sedatives	0.14	0.08	*	0.12	0.05	0.04	0.02	0.07
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.64	0.66	0.84	0.89	0.36	0.34	0.23	0.18

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.21P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	0.7053		0.0619		0.5806		0.1474	
Marijuana and Hashish	0.6324		0.0816		0.7734		0.0098	
Cocaine	0.4077		0.1703		0.8489		0.7474	
Crack	0.0626		0.5442		0.9227		*	
Heroin	0.0456		0.1323		0.5149		0.6494	
Hallucinogens	0.1238		0.8306		0.7321		0.9680	
LSD	0.9696		0.6663		0.0800		0.3364	
PCP	*		*		*		*	
Ecstasy	0.3103		0.9412		0.6965		0.4213	
Inhalants	0.9870		0.1114		0.8879		0.6649	
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.1675		0.0951		0.8679		0.2233	
Pain Relievers	0.4702		0.1153		0.8283		0.0750	
OxyContin <sup>®</sup>	0.1515		0.0060		0.6935		0.8131	
Tranquilizers	0.6602		0.7190		0.0975		0.0658	
Stimulants <sup>3</sup>	0.1384		0.2610		0.2158		0.1056	
Methamphetamine <sup>3</sup>	0.6152		0.0458		0.1062		0.3749	
Sedatives	0.0422		*		0.8662		0.1509	
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.6241		0.1559		0.8493		0.1800	

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.22C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2009 and 2010**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	272	268	127	160	243	273	123	134
Marijuana and Hashish	234	239	115	135	203	224	74	82
Cocaine	109	86	25	30	59	37	19	21
Crack	70	50	24	21	11	16	18	14
Heroin	28	30	18	16	12	*	*	20
Hallucinogens	61	52	4	8	42	23	20	20
LSD	13	3	*	*	*	*	*	*
PCP	17	*	*	*	*	*	*	20
Ecstasy	48	40	4	7	37	19	*	*
Inhalants	20	32	5	*	33	68	28	13
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	133	134	75	83	142	155	96	105
Pain Relievers	108	115	70	69	119	118	84	80
OxyContin <sup>®</sup>	37	35	37	46	15	22	*	*
Tranquilizers	80	78	46	51	81	74	41	49
Stimulants <sup>3</sup>	69	52	33	24	46	37	16	30
Methamphetamine <sup>3</sup>	49	28	28	20	30	25	3	23
Sedatives	36	30	23	13	23	52	19	37
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	174	163	75	88	159	164	103	107

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.22D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2009 and 2010**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	0.86	0.82	0.79	0.93	0.30	0.34	0.17	0.19
Marijuana and Hashish	0.76	0.76	0.71	0.79	0.26	0.28	0.10	0.11
Cocaine	0.37	0.30	0.16	0.18	0.08	0.05	0.03	0.03
Crack	0.24	0.18	0.15	0.13	0.01	0.02	0.03	0.02
Heroin	0.10	0.11	0.11	0.10	0.01	*	*	0.03
Hallucinogens	0.21	0.19	0.03	0.05	0.05	0.03	0.03	0.03
LSD	0.04	0.01	*	*	*	*	*	*
PCP	0.06	*	*	*	*	*	*	0.03
Ecstasy	0.17	0.14	0.03	0.04	0.05	0.02	*	*
Inhalants	0.07	0.12	0.03	*	0.04	0.09	0.04	0.02
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.45	0.45	0.47	0.50	0.18	0.19	0.13	0.15
Pain Relievers	0.37	0.39	0.44	0.42	0.15	0.15	0.12	0.11
OxyContin <sup>®</sup>	0.13	0.13	0.23	0.28	0.02	0.03	*	*
Tranquilizers	0.28	0.27	0.29	0.31	0.10	0.09	0.06	0.07
Stimulants <sup>3</sup>	0.24	0.19	0.21	0.15	0.06	0.05	0.02	0.04
Methamphetamine <sup>3</sup>	0.17	0.10	0.18	0.12	0.04	0.03	0.00	0.03
Sedatives	0.13	0.11	0.14	0.08	0.03	0.07	0.03	0.05
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.58	0.54	0.48	0.53	0.20	0.20	0.14	0.15

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.22P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	0.8913		0.6807		0.3841		0.4991	
Marijuana and Hashish	0.7127		0.6176		0.6058		0.5217	
Cocaine	0.1850		0.3262		0.2807		0.3936	
Crack	0.0660		0.9285		0.6146		0.9929	
Heroin	0.6728		0.9022		*		*	
Hallucinogens	0.5656		0.2341		0.1581		1.0000	
LSD	0.0951		*		*		*	
PCP	*		*		*		*	
Ecstasy	0.4644		0.4930		0.1430		*	
Inhalants	0.4153		*		0.1306		0.5576	
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	1.0000		0.5206		0.5318		0.4661	
Pain Relievers	0.9727		0.1636		0.7677		0.7076	
OxyContin <sup>®</sup>	0.5441		0.6211		0.6403		*	
Tranquilizers	0.2756		0.9971		0.9643		0.2283	
Stimulants <sup>3</sup>	0.0640		0.5139		0.5774		0.2818	
Methamphetamine <sup>3</sup>	0.0157		0.5481		0.7396		0.1814	
Sedatives	0.5907		0.5247		0.6538		0.2171	
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.5423		0.6817		0.7901		0.6024	

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.23C Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: Standard Errors of Numbers in Thousands, 2009 and 2010**

<b>Cigarette Use/Level of Alcohol Use</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
<b>TOTAL</b>	457	488	66	70	134	135	401	441
Heavy Alcohol Use <sup>1</sup>	230	218	25	24	99	94	202	189
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	269	246	37	41	97	94	246	222
Alcohol Use But Not Binge Use <sup>1</sup>	196	247	25	25	55	58	186	234
No Alcohol Use	191	227	47	48	51	58	175	213
<b>PAST MONTH CIGARETTE USE</b>	345	355	48	48	132	131	302	318
Heavy Alcohol Use <sup>1</sup>	190	174	22	22	85	76	166	156
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	191	172	29	31	76	75	173	149
Alcohol Use But Not Binge Use <sup>1</sup>	132	150	18	16	39	41	127	140
No Alcohol Use	138	168	23	21	39	45	127	160
<b>NO PAST MONTH CIGARETTE USE</b>	299	330	53	57	86	95	275	308
Heavy Alcohol Use <sup>1</sup>	116	109	12	11	39	44	106	98
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	177	169	25	25	55	55	165	159
Alcohol Use But Not Binge Use <sup>1</sup>	149	195	18	20	37	41	141	188
No Alcohol Use	133	146	40	42	32	38	123	134

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.23D Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: Standard Errors of Percentages, 2009 and 2010**

<b>Cigarette Use/Level of Alcohol Use</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
<b>TOTAL</b>	0.18	0.19	0.27	0.29	0.40	0.40	0.21	0.23
Heavy Alcohol Use <sup>1</sup>	1.14	1.10	2.58	2.76	1.22	1.23	1.48	1.39
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.57	0.56	1.66	1.85	0.82	0.80	0.70	0.68
Alcohol Use But Not Binge Use <sup>1</sup>	0.27	0.33	1.52	1.54	0.72	0.73	0.29	0.35
No Alcohol Use	0.16	0.18	0.21	0.22	0.38	0.41	0.20	0.24
<b>PAST MONTH CIGARETTE USE</b>	0.53	0.54	1.46	1.58	0.73	0.77	0.62	0.64
Heavy Alcohol Use <sup>1</sup>	1.49	1.53	2.92	3.11	1.48	1.50	2.13	2.15
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	1.00	0.95	2.49	2.69	1.17	1.29	1.33	1.19
Alcohol Use But Not Binge Use <sup>1</sup>	0.95	1.08	3.99	4.24	1.76	1.75	1.08	1.20
No Alcohol Use	0.72	0.81	2.17	2.19	1.34	1.48	0.79	0.93
<b>NO PAST MONTH CIGARETTE USE</b>	0.15	0.17	0.22	0.24	0.36	0.37	0.18	0.20
Heavy Alcohol Use <sup>1</sup>	1.46	1.32	5.68	5.65	1.84	1.79	1.73	1.58
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.64	0.65	2.12	2.31	1.00	0.97	0.77	0.78
Alcohol Use But Not Binge Use <sup>1</sup>	0.25	0.32	1.40	1.57	0.70	0.72	0.27	0.34
No Alcohol Use	0.13	0.14	0.19	0.20	0.31	0.34	0.17	0.19

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.23P Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

<b>Cigarette Use/Level of Alcohol Use</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
<b>TOTAL</b>	0.3272		0.8213		0.5823		0.3940	
Heavy Alcohol Use <sup>1</sup>	0.3640		0.8435		0.5553		0.4849	
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.7994		0.0043		0.2790		0.8955	
Alcohol Use But Not Binge Use <sup>1</sup>	0.0067		0.1244		0.0090		0.0451	
No Alcohol Use	0.2079		0.8381		0.0372		0.3987	
<b>PAST MONTH CIGARETTE USE</b>	0.8527		0.9708		0.3415		0.8351	
Heavy Alcohol Use <sup>1</sup>	0.3770		0.8527		0.5541		0.5595	
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.7025		0.2718		0.8034		0.6625	
Alcohol Use But Not Binge Use <sup>1</sup>	0.0629		0.9549		0.1122		0.1348	
No Alcohol Use	0.5080		0.7007		0.0619		0.6807	
<b>NO PAST MONTH CIGARETTE USE</b>	0.1124		0.2371		0.2252		0.2750	
Heavy Alcohol Use <sup>1</sup>	0.9410		0.6469		0.5422		0.7968	
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.9737		0.0100		0.1733		0.8723	
Alcohol Use But Not Binge Use <sup>1</sup>	0.0303		0.0538		0.0176		0.1243	
No Alcohol Use	0.2985		0.9025		0.1474		0.4991	

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.