

Table 6.24C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2009 and 2010

| Substance | CIGARETTE USE IN PAST MONTH | | | |
|-------------------------------------|-----------------------------|------|--------|-------|
| | Any Use | | No Use | |
| | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS¹ | N/A | N/A | 361 | 370 |
| Cigarettes | N/A | N/A | N/A | N/A |
| Smokeless Tobacco | 162 | 175 | 241 | 255 |
| Cigars | 261 | 262 | 271 | 256 |
| Pipe Tobacco | 100 | 90 | 112 | 123 |
| ALCOHOL | 719 | 706 | 1,437 | 1,445 |
| Binge Alcohol Use ² | 546 | 516 | 711 | 687 |
| Heavy Alcohol Use ² | 313 | 294 | 289 | 326 |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.24D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2009 and 2010

| Substance | CIGARETTE USE IN PAST MONTH | | | |
|-------------------------------------|-----------------------------|------|--------|------|
| | Any Use | | No Use | |
| | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS¹ | N/A | N/A | 0.18 | 0.18 |
| Cigarettes | N/A | N/A | N/A | N/A |
| Smokeless Tobacco | 0.27 | 0.30 | 0.12 | 0.13 |
| Cigars | 0.42 | 0.42 | 0.14 | 0.13 |
| Pipe Tobacco | 0.17 | 0.15 | 0.06 | 0.06 |
| ALCOHOL | 0.68 | 0.70 | 0.45 | 0.47 |
| Binge Alcohol Use ² | 0.68 | 0.70 | 0.32 | 0.30 |
| Heavy Alcohol Use ² | 0.49 | 0.47 | 0.15 | 0.16 |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.24P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2010 Versus 2009

| Substance | CIGARETTE USE IN PAST MONTH | | | |
|-------------------------------------|-----------------------------|------|--------|------|
| | Any Use | | No Use | |
| | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS¹ | N/A | | 0.7932 | |
| Cigarettes | N/A | | N/A | |
| Smokeless Tobacco | 0.5169 | | 0.7392 | |
| Cigars | 0.4073 | | 0.6051 | |
| Pipe Tobacco | 0.5543 | | 0.9069 | |
| ALCOHOL | 0.3273 | | 0.6662 | |
| Binge Alcohol Use ² | 0.2849 | | 0.4694 | |
| Heavy Alcohol Use ² | 0.3734 | | 0.7643 | |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

110407

Table 6.25C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2009 and 2010

| Substance | CIGARETTE USE IN PAST MONTH | | | |
|-------------------------------------|-----------------------------|------|--------|------|
| | Any Use | | No Use | |
| | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS¹ | N/A | N/A | 36 | 31 |
| Cigarettes | N/A | N/A | N/A | N/A |
| Smokeless Tobacco | 23 | 22 | 22 | 22 |
| Cigars | 35 | 28 | 25 | 23 |
| Pipe Tobacco | 16 | 15 | 15 | 9 |
| ALCOHOL | 52 | 55 | 71 | 66 |
| Binge Alcohol Use ² | 46 | 46 | 48 | 44 |
| Heavy Alcohol Use ² | 26 | 24 | 16 | 14 |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.25D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Percentages, 2009 and 2010

| Substance | CIGARETTE USE IN PAST MONTH | | | |
|-------------------------------------|-----------------------------|------|--------|------|
| | Any Use | | No Use | |
| | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS¹ | N/A | N/A | 0.15 | 0.14 |
| Cigarettes | N/A | N/A | N/A | N/A |
| Smokeless Tobacco | 0.94 | 0.98 | 0.10 | 0.10 |
| Cigars | 1.33 | 1.27 | 0.11 | 0.10 |
| Pipe Tobacco | 0.70 | 0.71 | 0.06 | 0.04 |
| ALCOHOL | 1.40 | 1.54 | 0.29 | 0.27 |
| Binge Alcohol Use ² | 1.46 | 1.50 | 0.20 | 0.19 |
| Heavy Alcohol Use ² | 1.09 | 1.05 | 0.07 | 0.06 |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.25P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2010 Versus 2009

| Substance | CIGARETTE USE IN PAST MONTH | | | |
|-------------------------------------|-----------------------------|------|--------|------|
| | Any Use | | No Use | |
| | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS¹ | N/A | | 0.0248 | |
| Cigarettes | N/A | | N/A | |
| Smokeless Tobacco | 0.9650 | | 0.8099 | |
| Cigars | 0.0048 | | 0.2254 | |
| Pipe Tobacco | 0.4067 | | 0.0034 | |
| ALCOHOL | 0.4580 | | 0.0472 | |
| Binge Alcohol Use ² | 0.2931 | | 0.0280 | |
| Heavy Alcohol Use ² | 0.0306 | | 0.5364 | |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

110407

Table 6.26C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2009 and 2010

| Substance | CIGARETTE USE IN PAST MONTH | | | |
|-------------------------------------|-----------------------------|------|--------|------|
| | Any Use | | No Use | |
| | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS¹ | N/A | N/A | 78 | 89 |
| Cigarettes | N/A | N/A | N/A | N/A |
| Smokeless Tobacco | 66 | 63 | 46 | 52 |
| Cigars | 90 | 82 | 59 | 71 |
| Pipe Tobacco | 33 | 34 | 23 | 23 |
| ALCOHOL | 211 | 189 | 244 | 259 |
| Binge Alcohol Use ² | 191 | 164 | 168 | 179 |
| Heavy Alcohol Use ² | 119 | 105 | 70 | 86 |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.26D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Percentages, 2009 and 2010

| Substance | CIGARETTE USE IN PAST MONTH | | | |
|-------------------------------------|-----------------------------|------|--------|------|
| | Any Use | | No Use | |
| | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS¹ | N/A | N/A | 0.33 | 0.35 |
| Cigarettes | N/A | N/A | N/A | N/A |
| Smokeless Tobacco | 0.48 | 0.50 | 0.21 | 0.22 |
| Cigars | 0.59 | 0.61 | 0.25 | 0.29 |
| Pipe Tobacco | 0.26 | 0.29 | 0.11 | 0.10 |
| ALCOHOL | 0.65 | 0.66 | 0.64 | 0.61 |
| Binge Alcohol Use ² | 0.76 | 0.73 | 0.57 | 0.58 |
| Heavy Alcohol Use ² | 0.76 | 0.71 | 0.30 | 0.35 |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.26P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2010 Versus 2009

| Substance | CIGARETTE USE IN PAST MONTH | | | |
|-------------------------------------|-----------------------------|------|--------|------|
| | Any Use | | No Use | |
| | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS¹ | N/A | | 0.0465 | |
| Cigarettes | N/A | | N/A | |
| Smokeless Tobacco | 0.4531 | | 0.2468 | |
| Cigars | 0.1905 | | 0.1079 | |
| Pipe Tobacco | 0.4361 | | 0.9671 | |
| ALCOHOL | 0.5257 | | 0.5920 | |
| Binge Alcohol Use ² | 0.1314 | | 0.9535 | |
| Heavy Alcohol Use ² | 0.7213 | | 0.3063 | |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

110407

Table 6.27C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2009 and 2010

| Substance | CIGARETTE USE IN PAST MONTH | | | |
|-------------------------------------|-----------------------------|------|--------|-------|
| | Any Use | | No Use | |
| | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS¹ | N/A | N/A | 351 | 354 |
| Cigarettes | N/A | N/A | N/A | N/A |
| Smokeless Tobacco | 145 | 161 | 235 | 244 |
| Cigars | 241 | 244 | 261 | 245 |
| Pipe Tobacco | 93 | 82 | 108 | 121 |
| ALCOHOL | 659 | 667 | 1,398 | 1,405 |
| Binge Alcohol Use ² | 495 | 479 | 681 | 657 |
| Heavy Alcohol Use ² | 280 | 277 | 281 | 310 |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.27D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2009 and 2010

| Substance | CIGARETTE USE IN PAST MONTH | | | |
|-------------------------------------|-----------------------------|------|--------|------|
| | Any Use | | No Use | |
| | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS¹ | N/A | N/A | 0.22 | 0.23 |
| Cigarettes | N/A | N/A | N/A | N/A |
| Smokeless Tobacco | 0.32 | 0.36 | 0.15 | 0.16 |
| Cigars | 0.51 | 0.52 | 0.17 | 0.16 |
| Pipe Tobacco | 0.21 | 0.18 | 0.07 | 0.08 |
| ALCOHOL | 0.86 | 0.89 | 0.56 | 0.59 |
| Binge Alcohol Use ² | 0.85 | 0.87 | 0.40 | 0.37 |
| Heavy Alcohol Use ² | 0.59 | 0.59 | 0.19 | 0.20 |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 6.27P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2010 Versus 2009

| Substance | CIGARETTE USE IN PAST MONTH | | | |
|-------------------------------------|-----------------------------|------|--------|------|
| | Any Use | | No Use | |
| | 2009 | 2010 | 2009 | 2010 |
| TOBACCO PRODUCTS¹ | N/A | | 0.9700 | |
| Cigarettes | N/A | | N/A | |
| Smokeless Tobacco | 0.5709 | | 0.9585 | |
| Cigars | 0.9766 | | 0.8518 | |
| Pipe Tobacco | 0.5489 | | 0.8694 | |
| ALCOHOL | 0.4420 | | 0.7089 | |
| Binge Alcohol Use ² | 0.5880 | | 0.4806 | |
| Heavy Alcohol Use ² | 0.5875 | | 0.9834 | |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.