

**Table 6.28C Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2009 and 2010**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>2</sup></b>	345	325	485	465	468	463	556	572
Cigarettes	313	294	434	408	421	433	514	528
Smokeless Tobacco	134	141	154	157	140	132	155	172
Cigars	168	164	204	212	182	165	163	176
Pipe Tobacco	47	44	92	62	80	79	77	103

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.28D Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2009 and 2010**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>2</sup></b>	1.22	1.24	0.81	0.81	0.60	0.58	0.40	0.42
Cigarettes	1.26	1.26	0.79	0.77	0.56	0.55	0.38	0.40
Smokeless Tobacco	0.73	0.77	0.35	0.37	0.19	0.18	0.13	0.14
Cigars	0.89	0.88	0.46	0.49	0.26	0.23	0.13	0.14
Pipe Tobacco	0.27	0.26	0.21	0.15	0.11	0.11	0.06	0.08

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.28P Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>2</sup></b>	0.1752		0.8732		0.2367		0.2419	
Cigarettes	0.3260		0.9156		0.5054		0.6962	
Smokeless Tobacco	0.4430		0.7430		0.9777		0.0879	
Cigars	0.6670		0.2854		0.2260		0.8747	
Pipe Tobacco	0.3441		0.2052		0.8038		0.0230	

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.29C Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2009 and 2010**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>2</sup></b>	28	25	42	42	26	25	48	45
Cigarettes	26	24	38	38	24	23	40	37
Smokeless Tobacco	10	10	19	19	9	12	21	19
Cigars	18	13	23	23	18	13	26	22
Pipe Tobacco	10	7	14	11	9	6	11	10

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.29D Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2009 and 2010**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>2</sup></b>	2.13	2.77	1.76	1.85	1.52	1.55	0.22	0.21
Cigarettes	2.66	2.97	1.77	1.81	1.45	1.45	0.19	0.18
Smokeless Tobacco	1.86	2.31	1.09	1.16	0.62	0.81	0.10	0.09
Cigars	2.73	2.92	1.27	1.39	1.14	0.90	0.12	0.10
Pipe Tobacco	1.88	1.64	0.81	0.68	0.60	0.46	0.05	0.05

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.29P Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>2</sup></b>	0.1076		0.7930		0.8416		0.2495	
Cigarettes	0.3670		0.3374		0.9070		0.5399	
Smokeless Tobacco	0.6650		0.5496		0.3670		0.9384	
Cigars	0.1115		0.9153		0.0683		0.0588	
Pipe Tobacco	0.0628		0.0790		0.6152		0.7925	

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.30C Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2009 and 2010**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>2</sup></b>	135	124	138	127	79	79	89	93
Cigarettes	119	105	126	112	68	69	82	86
Smokeless Tobacco	62	51	42	48	19	24	27	31
Cigars	64	66	64	62	39	38	43	43
Pipe Tobacco	22	19	24	27	16	16	15	17

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.30D Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2009 and 2010**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>2</sup></b>	1.03	1.08	0.89	0.83	0.94	0.93	0.65	0.60
Cigarettes	1.14	1.29	0.89	0.86	0.86	0.83	0.61	0.58
Smokeless Tobacco	1.03	0.92	0.43	0.49	0.28	0.33	0.21	0.24
Cigars	0.98	1.10	0.58	0.61	0.56	0.51	0.32	0.32
Pipe Tobacco	0.45	0.39	0.25	0.29	0.23	0.23	0.12	0.13

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.30P Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>2</sup></b>	0.0558		0.7804		0.6970		0.8453	
Cigarettes	0.0575		0.1185		0.6053		0.5302	
Smokeless Tobacco	0.1556		0.0627		0.1031		0.1968	
Cigars	0.7808		0.9110		0.1764		0.5454	
Pipe Tobacco	0.1490		0.2143		0.9964		0.3023	

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.31C Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2009 and 2010**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>2</sup></b>	307	305	466	443	458	454	535	551
Cigarettes	280	277	417	390	414	426	499	512
Smokeless Tobacco	113	132	146	146	138	128	150	167
Cigars	152	151	194	204	176	161	151	167
Pipe Tobacco	40	41	85	56	77	77	76	101

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.31D Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2009 and 2010**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>2</sup></b>	1.68	1.70	1.05	1.05	0.67	0.64	0.53	0.57
Cigarettes	1.71	1.71	1.03	0.99	0.62	0.61	0.51	0.54
Smokeless Tobacco	0.91	1.05	0.45	0.47	0.22	0.20	0.17	0.19
Cigars	1.17	1.16	0.58	0.63	0.28	0.25	0.17	0.19
Pipe Tobacco	0.34	0.35	0.27	0.18	0.12	0.12	0.09	0.11

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 6.31P Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2009	2010	2009	2010	2009	2010	2009	2010
<b>TOBACCO PRODUCTS<sup>2</sup></b>	0.3963		0.8091		0.2367		0.2130	
Cigarettes	0.6741		0.8071		0.5263		0.6013	
Smokeless Tobacco	0.8072		0.8374		0.7199		0.1353	
Cigars	0.8622		0.2595		0.4100		0.7880	
Pipe Tobacco	0.9388		0.1418		0.8321		0.0309	

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.