

**Table 6.47C Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

<b>Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users<sup>1</sup></b>	<b>Total (2007-2008)</b>	<b>Total (2009-2010)</b>	<b>Aged 12-17 (2007-2008)</b>	<b>Aged 12-17 (2009-2010)</b>	<b>Aged 18-25 (2007-2008)</b>	<b>Aged 18-25 (2009-2010)</b>	<b>Aged 26+ (2007-2008)</b>	<b>Aged 26+ (2009-2010)</b>
From One Doctor	93	105	15	16	26	26	89	101
From More Than One Doctor	36	35	5	5	9	9	35	34
Wrote Fake Prescription	11	9	1	2	3	1	10	9
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	11	6	3	5	7	2	8	4
From Friend or Relative for Free	159	162	25	24	58	59	142	150
Bought from Friend or Relative	50	72	11	9	25	27	42	66
Took from Friend or Relative without Asking	41	40	12	11	16	15	36	35
Bought from Drug Dealer or Other Stranger	42	45	8	7	15	19	38	41
Bought on the Internet	14	19	1	0	3	1	14	19
Some Other Way	43	47	9	10	13	14	38	42

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.47D Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

<b>Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users<sup>1</sup></b>	<b>Total (2007-2008)</b>	<b>Total (2009-2010)</b>	<b>Aged 12-17 (2007-2008)</b>	<b>Aged 12-17 (2009-2010)</b>	<b>Aged 18-25 (2007-2008)</b>	<b>Aged 18-25 (2009-2010)</b>	<b>Aged 26+ (2007-2008)</b>	<b>Aged 26+ (2009-2010)</b>
From One Doctor	0.76	0.82	1.02	1.08	0.65	0.66	1.31	1.39
From More Than One Doctor	0.32	0.30	0.40	0.38	0.23	0.23	0.55	0.52
Wrote Fake Prescription	0.10	0.08	0.08	0.16	0.08	0.04	0.17	0.14
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.10	0.06	0.23	0.36	0.18	0.05	0.13	0.06
From Friend or Relative for Free	0.94	1.00	1.28	1.33	0.95	0.95	1.56	1.67
Bought from Friend or Relative	0.44	0.58	0.75	0.70	0.64	0.67	0.68	0.97
Took from Friend or Relative without Asking	0.36	0.34	0.81	0.83	0.40	0.40	0.58	0.54
Bought from Drug Dealer or Other Stranger	0.37	0.38	0.54	0.53	0.41	0.49	0.62	0.62
Bought on the Internet	0.13	0.17	0.10	0.02	0.07	0.04	0.23	0.29
Some Other Way	0.37	0.40	0.61	0.78	0.36	0.37	0.61	0.65

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.47P Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:  
P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	0.5666		0.5459		0.5195		0.5753	
From More Than One Doctor	0.5333		0.4017		0.1989		0.3956	
Wrote Fake Prescription	0.6983		0.1174		0.2209		0.7017	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.0154		0.0972		0.0009		0.0788	
From Friend or Relative for Free	0.5401		0.8832		0.4270		0.6350	
Bought from Friend or Relative	0.0004		0.6671		0.0277		0.0025	
Took from Friend or Relative without Asking	0.2216		0.2008		0.3799		0.6929	
Bought from Drug Dealer or Other Stranger	0.9394		0.3307		0.8751		0.7888	
Bought on the Internet	0.9898		0.0911		0.1528		0.8041	
Some Other Way	0.7248		0.0994		0.3755		0.7980	

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.48C Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Pain Relievers <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	141	142	17	16	46	46	130	134
From More Than One Doctor	22	27	4	6	8	10	20	23
Wrote Fake Prescription	2	11	0	1	2	1	*	11
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	14	5	2	3	3	3	13	2
From Friend or Relative for Free	30	40	8	7	13	15	26	37
Bought from Friend or Relative	21	24	5	5	11	13	16	20
Took from Friend or Relative without Asking	9	15	5	8	5	8	5	10
Bought from Drug Dealer or Other Stranger	14	18	4	3	8	9	11	14
Bought on the Internet	4	5	0	1	2	2	3	4
Some Other Way	8	12	4	5	5	5	5	10

\*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.48D Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Pain Relievers <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	0.89	1.01	2.03	2.18	1.12	1.29	1.25	1.48
From More Than One Doctor	0.40	0.47	0.82	1.15	0.44	0.54	0.62	0.69
Wrote Fake Prescription	0.04	0.19	0.07	0.12	0.11	0.06	*	0.31
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.25	0.08	0.47	0.69	0.16	0.16	0.41	0.05
From Friend or Relative for Free	0.53	0.69	1.42	1.31	0.70	0.79	0.78	1.07
Bought from Friend or Relative	0.38	0.43	0.89	1.08	0.62	0.72	0.51	0.59
Took from Friend or Relative without Asking	0.16	0.26	0.96	1.44	0.28	0.42	0.15	0.31
Bought from Drug Dealer or Other Stranger	0.26	0.32	0.77	0.71	0.43	0.52	0.34	0.42
Bought on the Internet	0.07	0.09	0.07	0.11	0.09	0.10	0.10	0.13
Some Other Way	0.15	0.21	0.79	0.91	0.29	0.30	0.16	0.29

\*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.48P Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source Where Friend or Relative Obtained Pain Relievers <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	0.0881		0.5294		0.0722		0.2175	
From More Than One Doctor	0.8093		0.7023		0.3791		0.8792	
Wrote Fake Prescription	0.3825		0.3119		0.4369		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3036		0.9690		0.8870		0.2087	
From Friend or Relative for Free	0.8993		0.0846		0.9509		0.5517	
Bought from Friend or Relative	0.3433		0.9587		0.6060		0.3421	
Took from Friend or Relative without Asking	0.0631		0.0205		0.1073		0.9503	
Bought from Drug Dealer or Other Stranger	0.0982		0.7144		0.0764		0.2665	
Bought on the Internet	0.6047		0.8247		0.7781		0.6789	
Some Other Way	0.2614		0.5160		0.9021		0.1942	

\*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.49C Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	79	87	11	11	19	20	75	85
From More Than One Doctor	38	36	4	5	9	9	36	33
Wrote Fake Prescription	6	11	1	2	3	1	5	10
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	10	7	3	4	4	3	8	6
From Friend or Relative for Free	100	109	15	16	34	38	91	101
Bought from Friend or Relative	56	70	9	8	24	25	50	63
Took from Friend or Relative without Asking	39	47	9	9	15	16	36	43
Bought from Drug Dealer or Other Stranger	40	37	7	7	17	19	36	31
Bought on the Internet	14	19	2	*	4	5	13	18
Some Other Way	26	39	6	6	10	11	23	37

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.49D Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

<b>Source for Past Month Nonmedical Use among Past Month Pain Reliever Users<sup>1</sup></b>	<b>Total (2007-2008)</b>	<b>Total (2009-2010)</b>	<b>Aged 12-17 (2007-2008)</b>	<b>Aged 12-17 (2009-2010)</b>	<b>Aged 18-25 (2007-2008)</b>	<b>Aged 18-25 (2009-2010)</b>	<b>Aged 26+ (2007-2008)</b>	<b>Aged 26+ (2009-2010)</b>
From One Doctor	1.42	1.51	1.88	1.85	1.22	1.21	2.35	2.42
From More Than One Doctor	0.82	0.75	0.82	0.93	0.66	0.62	1.35	1.16
Wrote Fake Prescription	0.14	0.22	0.19	0.39	0.23	0.09	0.20	0.37
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.21	0.16	0.63	0.75	0.31	0.21	0.31	0.20
From Friend or Relative for Free	1.53	1.52	2.11	2.17	1.45	1.35	2.49	2.46
Bought from Friend or Relative	1.16	1.28	1.60	1.47	1.47	1.44	1.78	2.03
Took from Friend or Relative without Asking	0.85	0.93	1.68	1.56	0.98	1.01	1.34	1.47
Bought from Drug Dealer or Other Stranger	0.86	0.77	1.32	1.28	1.10	1.15	1.34	1.10
Bought on the Internet	0.31	0.40	0.38	*	0.29	0.34	0.51	0.66
Some Other Way	0.58	0.80	1.24	1.18	0.75	0.77	0.88	1.30

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.49P Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	0.8754		0.7432		1.0000		0.8781	
From More Than One Doctor	0.2847		0.5576		0.6795		0.2656	
Wrote Fake Prescription	0.6386		0.1943		0.0622		0.4000	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.0686		0.9333		0.0307		0.2874	
From Friend or Relative for Free	0.4955		0.9737		0.1454		0.7853	
Bought from Friend or Relative	0.0126		0.2910		0.1684		0.0201	
Took from Friend or Relative without Asking	0.8550		0.1904		0.7906		0.9498	
Bought from Drug Dealer or Other Stranger	0.6689		0.3288		0.5822		0.5979	
Bought on the Internet	0.9523		*		0.8242		0.8021	
Some Other Way	0.1620		0.3976		0.1732		0.2218	

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.50C Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Pain Relievers <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	84	93	10	10	27	30	79	89
From More Than One Doctor	27	31	5	5	10	11	25	28
Wrote Fake Prescription	9	13	2	1	3	4	8	13
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	16	5	3	1	4	5	15	2
From Friend or Relative for Free	34	37	6	6	13	14	31	33
Bought from Friend or Relative	27	31	5	6	14	14	23	27
Took from Friend or Relative without Asking	15	17	5	5	6	9	14	13
Bought from Drug Dealer or Other Stranger	18	27	5	5	10	13	15	23
Bought on the Internet	7	3	1	1	4	2	5	2
Some Other Way	14	11	3	4	4	7	12	8

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.50D Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

<b>Source Where Friend or Relative Obtained Pain Relievers<sup>1</sup></b>	<b>Total (2007-2008)</b>	<b>Total (2009-2010)</b>	<b>Aged 12-17 (2007-2008)</b>	<b>Aged 12-17 (2009-2010)</b>	<b>Aged 18-25 (2007-2008)</b>	<b>Aged 18-25 (2009-2010)</b>	<b>Aged 26+ (2007-2008)</b>	<b>Aged 26+ (2009-2010)</b>
From One Doctor	1.58	1.49	2.93	3.33	1.72	1.78	2.39	2.30
From More Than One Doctor	1.10	1.15	2.19	2.34	1.21	1.17	1.72	1.80
Wrote Fake Prescription	0.37	0.52	0.79	0.27	0.44	0.47	0.56	0.86
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.66	0.21	1.33	0.60	0.55	0.55	1.07	0.15
From Friend or Relative for Free	1.39	1.35	2.67	2.53	1.56	1.46	2.13	2.14
Bought from Friend or Relative	1.14	1.17	2.24	2.48	1.62	1.54	1.67	1.74
Took from Friend or Relative without Asking	0.64	0.65	2.10	2.29	0.72	1.01	0.98	0.89
Bought from Drug Dealer or Other Stranger	0.75	1.02	2.19	2.06	1.26	1.41	1.05	1.50
Bought on the Internet	0.29	0.12	0.68	0.43	0.56	0.29	0.36	0.12
Some Other Way	0.58	0.46	1.42	1.94	0.56	0.85	0.90	0.53

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.50P Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source Where Friend or Relative Obtained Pain Relievers <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	0.8154		0.1929		0.6379		0.8468	
From More Than One Doctor	0.3790		0.4944		0.4503		0.5887	
Wrote Fake Prescription	0.6957		0.2185		0.5718		0.4453	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.0580		0.0656		0.8061		0.1030	
From Friend or Relative for Free	0.5898		0.0695		0.8747		0.8191	
Bought from Friend or Relative	0.8520		0.6157		0.7889		0.8252	
Took from Friend or Relative without Asking	0.8020		0.9053		0.1703		0.7249	
Bought from Drug Dealer or Other Stranger	0.3860		0.2650		0.3821		0.4732	
Bought on the Internet	0.0090		0.4105		0.0235		0.1716	
Some Other Way	0.9898		0.4406		0.0766		0.2161	

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.51C Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	43	61	4	6	10	13	41	59
From More Than One Doctor	14	19	2	2	5	5	12	19
Wrote Fake Prescription	3	1	2	1	2	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	5	2	2	4	5	*	*
From Friend or Relative for Free	111	120	13	13	35	36	105	114
Bought from Friend or Relative	44	42	9	7	21	22	37	35
Took from Friend or Relative without Asking	45	20	6	5	10	9	43	18
Bought from Drug Dealer or Other Stranger	23	30	6	5	13	13	18	24
Bought on the Internet	43	15	1	1	2	2	43	15
Some Other Way	32	26	4	4	10	9	30	24

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.51D Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

<b>Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users<sup>1</sup></b>	<b>Total (2007-2008)</b>	<b>Total (2009-2010)</b>	<b>Aged 12-17 (2007-2008)</b>	<b>Aged 12-17 (2009-2010)</b>	<b>Aged 18-25 (2007-2008)</b>	<b>Aged 18-25 (2009-2010)</b>	<b>Aged 26+ (2007-2008)</b>	<b>Aged 26+ (2009-2010)</b>
From One Doctor	0.87	1.12	1.11	1.47	0.64	0.78	1.42	1.79
From More Than One Doctor	0.28	0.38	0.54	0.51	0.33	0.30	0.44	0.60
Wrote Fake Prescription	0.06	0.02	0.45	0.21	0.13	0.06	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.09	0.10	0.42	0.47	0.26	0.29	*	*
From Friend or Relative for Free	1.54	1.46	2.36	2.38	1.46	1.41	2.45	2.24
Bought from Friend or Relative	0.88	0.80	1.90	1.66	1.13	1.18	1.28	1.10
Took from Friend or Relative without Asking	0.91	0.40	1.43	1.28	0.63	0.53	1.48	0.58
Bought from Drug Dealer or Other Stranger	0.48	0.56	1.55	1.24	0.77	0.78	0.63	0.77
Bought on the Internet	0.87	0.29	0.36	0.14	0.12	0.12	1.48	0.47
Some Other Way	0.64	0.50	0.89	1.10	0.63	0.54	1.03	0.76

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.51P Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:  
P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	0.0171		0.1831		0.0538		0.0713	
From More Than One Doctor	0.8165		0.7861		0.7087		0.6903	
Wrote Fake Prescription	0.3098		0.4707		0.5464		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5875		0.9646		0.6356		*	
From Friend or Relative for Free	0.9084		0.2289		0.9466		0.6761	
Bought from Friend or Relative	0.9081		0.1534		0.9562		0.5853	
Took from Friend or Relative without Asking	0.0493		0.5445		0.3005		0.1069	
Bought from Drug Dealer or Other Stranger	0.8656		0.4717		0.5670		0.3561	
Bought on the Internet	0.1105		0.2179		0.7895		0.1179	
Some Other Way	0.9205		0.5041		0.7696		0.8931	

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.52C Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Tranquilizers <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	91	104	7	9	25	26	88	101
From More Than One Doctor	17	15	3	3	5	7	16	13
Wrote Fake Prescription	1	2	0	*	1	1	*	2
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	4	0	*	3	1	3	3
From Friend or Relative for Free	32	25	5	5	13	13	29	21
Bought from Friend or Relative	19	11	4	3	11	9	15	5
Took from Friend or Relative without Asking	9	5	4	3	3	4	*	2
Bought from Drug Dealer or Other Stranger	12	13	2	3	5	8	11	10
Bought on the Internet	3	4	0	*	3	1	0	4
Some Other Way	14	7	2	2	5	2	13	6

\*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.52D Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Tranquilizers <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	1.62	1.31	3.80	3.80	1.99	1.92	2.22	1.60
From More Than One Doctor	0.67	0.57	2.26	1.70	0.64	0.89	0.96	0.76
Wrote Fake Prescription	0.04	0.09	0.26	*	0.11	0.09	*	0.11
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.17	0.13	0.29	*	0.43	0.07	0.17	0.19
From Friend or Relative for Free	1.18	0.93	3.06	2.65	1.53	1.60	1.66	1.16
Bought from Friend or Relative	0.73	0.42	2.49	1.68	1.31	1.11	0.88	0.31
Took from Friend or Relative without Asking	0.33	0.20	2.74	1.83	0.46	0.51	*	0.10
Bought from Drug Dealer or Other Stranger	0.47	0.50	1.61	1.69	0.60	0.96	0.67	0.60
Bought on the Internet	0.10	0.16	0.20	*	0.34	0.10	0.03	0.24
Some Other Way	0.55	0.26	1.58	1.53	0.59	0.24	0.77	0.35

\*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.52P Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: *P* Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source Where Friend or Relative Obtained Tranquilizers <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	0.0445		0.0124		0.5571		0.0934	
From More Than One Doctor	0.6489		0.5849		0.1852		0.3871	
Wrote Fake Prescription	0.2249		*		0.9157		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5507		*		0.1030		0.7267	
From Friend or Relative for Free	0.4549		0.2855		0.8893		0.5472	
Bought from Friend or Relative	0.0027		0.0955		0.0462		0.0428	
Took from Friend or Relative without Asking	0.3638		0.1724		0.7606		*	
Bought from Drug Dealer or Other Stranger	0.2869		0.8986		0.0184		0.8981	
Bought on the Internet	0.6020		*		0.2105		0.1280	
Some Other Way	0.2905		0.7013		0.1018		0.4749	

\*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.53C Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	31	33	3	4	7	10	29	32
From More Than One Doctor	12	6	*	*	3	2	11	5
Wrote Fake Prescription	2	1	*	1	2	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	3	2	*	*	3	*	*	*
From Friend or Relative for Free	68	82	8	5	19	19	65	78
Bought from Friend or Relative	30	39	4	4	14	14	27	36
Took from Friend or Relative without Asking	24	15	4	4	8	7	22	13
Bought from Drug Dealer or Other Stranger	27	18	4	4	10	9	25	14
Bought on the Internet	24	7	1	1	2	3	*	7
Some Other Way	11	16	2	2	5	4	10	15

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.53D Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	1.74	1.70	2.19	3.06	1.30	1.66	2.66	2.51
From More Than One Doctor	0.70	0.31	*	*	0.60	0.31	1.10	0.43
Wrote Fake Prescription	0.12	0.05	*	0.63	0.39	0.12	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.20	0.09	*	*	0.58	*	*	*
From Friend or Relative for Free	2.37	2.41	4.31	4.06	2.48	2.36	3.68	3.35
Bought from Friend or Relative	1.74	1.91	3.12	3.53	2.26	2.12	2.53	2.73
Took from Friend or Relative without Asking	1.35	0.79	2.72	3.19	1.41	1.33	2.08	1.03
Bought from Drug Dealer or Other Stranger	1.55	0.94	2.71	2.97	1.73	1.64	2.35	1.13
Bought on the Internet	1.43	0.38	1.10	0.65	0.37	0.48	*	0.54
Some Other Way	0.69	0.82	1.56	1.89	0.87	0.70	1.01	1.21

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.53P Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	0.1186		0.3762		0.0621		0.3733	
From More Than One Doctor	0.1432		*		0.3013		0.1849	
Wrote Fake Prescription	0.6944		*		0.4689		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.1598		*		*		*	
From Friend or Relative for Free	0.5541		0.0465		0.3655		0.7531	
Bought from Friend or Relative	0.0439		0.1608		0.3316		0.0674	
Took from Friend or Relative without Asking	0.1277		0.9659		0.7700		0.1777	
Bought from Drug Dealer or Other Stranger	0.0620		0.7755		0.8518		0.0701	
Bought on the Internet	0.1554		0.3767		0.8374		*	
Some Other Way	0.8465		0.8249		0.7827		0.9052	

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.54C Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Tranquilizers <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	55	68	*	*	14	14	54	67
From More Than One Doctor	24	13	*	*	5	6	23	12
Wrote Fake Prescription	0	1	*	*	*	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1	2	*	*	*	1	*	*
From Friend or Relative for Free	17	24	*	*	10	8	13	22
Bought from Friend or Relative	11	12	*	*	7	6	8	9
Took from Friend or Relative without Asking	7	10	*	*	3	1	6	*
Bought from Drug Dealer or Other Stranger	13	15	*	*	4	5	13	14
Bought on the Internet	1	4	*	*	1	*	0	*
Some Other Way	10	3	*	*	3	2	10	2

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.54D Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Tranquilizers <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	2.77	2.89	*	*	3.44	3.26	3.76	3.79
From More Than One Doctor	2.35	1.28	*	*	1.80	2.08	3.41	1.57
Wrote Fake Prescription	0.02	0.08	*	*	*	0.32	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.15	0.16	*	*	*	0.50	*	*
From Friend or Relative for Free	1.77	2.21	*	*	3.24	2.75	2.04	2.84
Bought from Friend or Relative	1.18	1.21	*	*	2.57	2.36	1.27	1.28
Took from Friend or Relative without Asking	0.73	0.93	*	*	1.03	0.42	0.88	*
Bought from Drug Dealer or Other Stranger	1.34	1.47	*	*	1.55	1.83	1.91	1.92
Bought on the Internet	0.08	0.37	*	*	0.30	*	0.01	*
Some Other Way	1.10	0.33	*	*	1.05	0.83	1.57	0.31

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.54P Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source Where Friend or Relative Obtained Tranquilizers <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	0.7382		*		0.6328		0.9112	
From More Than One Doctor	0.1405		*		0.8920		0.1171	
Wrote Fake Prescription	0.1127		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7890		*		*		*	
From Friend or Relative for Free	0.9922		*		0.2166		0.3758	
Bought from Friend or Relative	0.5060		*		0.9940		0.8493	
Took from Friend or Relative without Asking	0.9627		*		0.1866		*	
Bought from Drug Dealer or Other Stranger	0.8434		*		0.6231		0.9212	
Bought on the Internet	0.2824		*		*		*	
Some Other Way	0.1688		*		0.5452		0.1599	

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.55C Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

Source for Most Recent Nonmedical Use among Past Year Stimulant Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	27	26	4	5	8	10	25	24
From More Than One Doctor	6	6	2	2	4	3	*	5
Wrote Fake Prescription	3	7	2	1	2	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	6	10	2	2	4	2	4	10
From Friend or Relative for Free	58	62	11	9	25	32	51	53
Bought from Friend or Relative	28	29	6	6	18	18	20	20
Took from Friend or Relative without Asking	16	19	5	4	7	7	14	17
Bought from Drug Dealer or Other Stranger	16	31	4	4	9	9	13	29
Bought on the Internet	15	11	1	2	4	3	14	10
Some Other Way	11	17	3	3	5	5	9	16

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

<sup>1</sup> Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.55D Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

<b>Source for Most Recent Nonmedical Use among Past Year Stimulant Users<sup>1</sup></b>	<b>Total (2007-2008)</b>	<b>Total (2009-2010)</b>	<b>Aged 12-17 (2007-2008)</b>	<b>Aged 12-17 (2009-2010)</b>	<b>Aged 18-25 (2007-2008)</b>	<b>Aged 18-25 (2009-2010)</b>	<b>Aged 26+ (2007-2008)</b>	<b>Aged 26+ (2009-2010)</b>
From One Doctor	1.32	1.13	1.53	2.02	0.94	0.90	2.78	2.43
From More Than One Doctor	0.32	0.29	0.60	0.89	0.44	0.25	*	0.58
Wrote Fake Prescription	0.13	0.32	0.65	0.56	0.22	0.01	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.30	0.46	0.72	1.04	0.44	0.16	0.48	1.03
From Friend or Relative for Free	1.92	1.87	2.83	3.32	1.90	1.74	3.87	4.00
Bought from Friend or Relative	1.30	1.27	1.98	2.60	1.75	1.55	2.30	2.08
Took from Friend or Relative without Asking	0.79	0.83	1.60	1.66	0.77	0.67	1.60	1.76
Bought from Drug Dealer or Other Stranger	0.80	1.31	1.51	1.69	1.10	0.80	1.47	2.94
Bought on the Internet	0.77	0.49	0.19	0.75	0.50	0.33	1.64	1.07
Some Other Way	0.55	0.78	1.20	1.12	0.59	0.53	1.06	1.71

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

<sup>1</sup> Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.55P Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:  
P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source for Most Recent Nonmedical Use among Past Year Stimulant Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	0.6106		0.8042		0.1269		0.9801	
From More Than One Doctor	0.7868		0.8816		0.2289		*	
Wrote Fake Prescription	0.6517		0.8710		0.1492		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.9726		0.4104		0.0312		0.4770	
From Friend or Relative for Free	0.7686		0.3432		0.6677		0.6838	
Bought from Friend or Relative	0.4456		0.1445		0.3516		0.7016	
Took from Friend or Relative without Asking	0.5489		0.4946		0.4548		0.9420	
Bought from Drug Dealer or Other Stranger	0.6134		0.8098		0.1045		0.8074	
Bought on the Internet	0.1217		0.2754		0.1840		0.1652	
Some Other Way	0.1432		0.3570		0.4811		0.1071	

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

<sup>1</sup> Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.56C Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Stimulants <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	43	50	9	7	22	24	*	43
From More Than One Doctor	7	10	3	3	3	5	5	8
Wrote Fake Prescription	0	1	0	*	*	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1	3	0	*	1	3	*	*
From Friend or Relative for Free	23	17	3	3	7	8	*	*
Bought from Friend or Relative	11	9	4	2	5	8	9	3
Took from Friend or Relative without Asking	3	3	3	1	1	2	*	1
Bought from Drug Dealer or Other Stranger	3	13	2	2	2	6	*	*
Bought on the Internet	4	6	*	*	2	*	*	*
Some Other Way	4	10	0	0	1	1	3	*

\*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.56D Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Stimulants <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	2.61	2.32	4.72	4.78	2.40	2.43	*	4.75
From More Than One Doctor	0.76	0.99	2.59	2.88	0.86	1.04	1.26	1.88
Wrote Fake Prescription	0.01	0.07	0.07	*	*	0.09	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.11	0.26	0.43	*	0.21	0.54	*	*
From Friend or Relative for Free	2.35	1.64	2.68	3.07	1.75	1.64	*	*
Bought from Friend or Relative	1.19	0.88	3.11	2.26	1.37	1.55	2.18	0.73
Took from Friend or Relative without Asking	0.35	0.28	2.42	1.66	0.35	0.37	*	0.19
Bought from Drug Dealer or Other Stranger	0.30	1.28	1.62	1.99	0.48	1.15	*	*
Bought on the Internet	0.46	0.62	*	*	0.48	*	*	*
Some Other Way	0.41	1.00	0.32	0.47	0.34	0.18	0.87	*

\*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.56P Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: *P* Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source Where Friend or Relative Obtained Stimulants <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	0.8414		0.5883		0.7127		*	
From More Than One Doctor	0.5051		0.8835		0.6621		0.5032	
Wrote Fake Prescription	0.1725		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2538		*		0.1818		*	
From Friend or Relative for Free	0.1830		0.9427		0.2059		*	
Bought from Friend or Relative	0.2269		0.2684		0.6909		0.0944	
Took from Friend or Relative without Asking	0.5681		0.6021		0.9124		*	
Bought from Drug Dealer or Other Stranger	0.0094		0.7889		0.0167		*	
Bought on the Internet	0.8961		*		*		*	
Some Other Way	0.5518		0.9159		0.6608		*	

\*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.57C Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

Source for Past Month Nonmedical Use among Past Month Stimulant Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	17	25	3	4	6	9	*	22
From More Than One Doctor	7	7	1	1	5	2	*	6
Wrote Fake Prescription	2	1	2	1	2	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	6	1	*	3	1	*	*
From Friend or Relative for Free	37	37	5	5	12	18	*	32
Bought from Friend or Relative	20	24	4	4	12	13	15	19
Took from Friend or Relative without Asking	13	18	3	2	6	6	*	16
Bought from Drug Dealer or Other Stranger	12	15	3	4	7	7	9	13
Bought on the Internet	13	8	*	1	4	3	*	*
Some Other Way	4	9	2	2	2	2	*	8

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

<sup>1</sup> Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.57D Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

Source for Past Month Nonmedical Use among Past Month Stimulant Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	2.42	2.58	3.38	4.18	2.30	2.34	*	4.81
From More Than One Doctor	1.05	0.80	1.69	1.12	1.84	0.70	*	1.51
Wrote Fake Prescription	0.36	0.15	1.98	1.31	0.62	0.16	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.74	0.70	1.16	*	1.04	0.17	*	*
From Friend or Relative for Free	3.56	3.06	4.47	5.19	3.41	3.04	*	5.54
Bought from Friend or Relative	2.79	2.55	3.76	4.04	3.61	2.90	4.63	4.25
Took from Friend or Relative without Asking	1.79	1.97	3.38	2.83	2.01	1.68	*	3.71
Bought from Drug Dealer or Other Stranger	1.74	1.69	3.02	3.94	2.53	1.83	2.77	3.02
Bought on the Internet	1.90	0.93	*	0.80	1.63	0.76	*	*
Some Other Way	0.60	1.09	2.38	2.28	0.85	0.73	*	2.13

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

<sup>1</sup> Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.57P Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source for Past Month Nonmedical Use among Past Month Stimulant Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	0.0447		0.9417		0.0602		*	
From More Than One Doctor	0.4954		0.4444		0.0923		*	
Wrote Fake Prescription	0.2346		0.7357		0.2440		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5005		*		0.0998		*	
From Friend or Relative for Free	0.4467		0.8419		0.4722		*	
Bought from Friend or Relative	0.9554		0.4486		0.4492		0.6500	
Took from Friend or Relative without Asking	0.5808		0.2294		0.8430		*	
Bought from Drug Dealer or Other Stranger	0.9211		0.1592		0.1249		0.6137	
Bought on the Internet	0.1493		*		0.2577		*	
Some Other Way	0.1743		0.3255		0.9634		*	

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

<sup>1</sup> Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.58C Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Stimulants <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	*	28	*	*	11	12	*	*
From More Than One Doctor	4	7	*	*	2	5	*	*
Wrote Fake Prescription	*	0	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	0	*	*	*	*	*	*
From Friend or Relative for Free	*	16	*	*	5	6	*	*
Bought from Friend or Relative	*	8	*	*	3	6	*	*
Took from Friend or Relative without Asking	5	5	*	*	2	*	*	*
Bought from Drug Dealer or Other Stranger	7	5	*	*	2	3	*	*
Bought on the Internet	4	*	*	*	*	*	*	*
Some Other Way	5	*	*	*	*	0	*	*

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

<sup>1</sup> Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.58D Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Stimulants <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	*	4.56	*	*	4.03	4.39	*	*
From More Than One Doctor	1.15	2.03	*	*	1.90	2.95	*	*
Wrote Fake Prescription	*	0.03	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	0.09	*	*	*	*	*	*
From Friend or Relative for Free	*	4.10	*	*	3.66	3.57	*	*
Bought from Friend or Relative	*	2.33	*	*	2.27	3.42	*	*
Took from Friend or Relative without Asking	1.49	1.29	*	*	1.51	*	*	*
Bought from Drug Dealer or Other Stranger	2.12	1.30	*	*	1.71	2.01	*	*
Bought on the Internet	1.38	*	*	*	*	*	*	*
Some Other Way	1.66	*	*	*	*	0.25	*	*

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

<sup>1</sup> Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.58P Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source Where Friend or Relative Obtained Stimulants <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	*		*		0.8906		*	
From More Than One Doctor	0.3905		*		0.3059		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	*		*		0.4651		*	
Bought from Friend or Relative	*		*		0.3874		*	
Took from Friend or Relative without Asking	0.8772		*		*		*	
Bought from Drug Dealer or Other Stranger	0.4056		*		0.4339		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine use items added in 2005 and 2006.

<sup>1</sup> Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.59C Source Where Methamphetamine was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

<b>Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users<sup>1</sup></b>	<b>Total (2007-2008)</b>	<b>Total (2009-2010)</b>	<b>Aged 12-17 (2007-2008)</b>	<b>Aged 12-17 (2009-2010)</b>	<b>Aged 18-25 (2007-2008)</b>	<b>Aged 18-25 (2009-2010)</b>	<b>Aged 26+ (2007-2008)</b>	<b>Aged 26+ (2009-2010)</b>
From Friend or Relative for Free	35	33	6	*	14	14	31	30
Bought from Friend or Relative	37	52	4	3	9	9	35	49
Took from Friend or Relative without Asking	6	14	2	2	3	3	4	14
Bought from Drug Dealer or Other Stranger	32	25	3	3	9	9	30	23
Bought on the Internet	4	5	*	*	0	2	*	*
Some Other Way	9	12	1	*	3	3	9	11

\*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.59D Source Where Methamphetamine was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

<b>Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users<sup>1</sup></b>	<b>Total (2007-2008)</b>	<b>Total (2009-2010)</b>	<b>Aged 12-17 (2007-2008)</b>	<b>Aged 12-17 (2009-2010)</b>	<b>Aged 18-25 (2007-2008)</b>	<b>Aged 18-25 (2009-2010)</b>	<b>Aged 26+ (2007-2008)</b>	<b>Aged 26+ (2009-2010)</b>
From Friend or Relative for Free	2.88	3.42	5.51	*	3.05	3.99	4.37	4.56
Bought from Friend or Relative	2.90	4.18	4.11	3.96	2.52	3.31	4.56	5.61
Took from Friend or Relative without Asking	0.57	1.45	2.78	3.01	1.09	0.99	0.64	2.18
Bought from Drug Dealer or Other Stranger	2.88	2.50	3.38	3.72	2.64	3.18	4.46	3.49
Bought on the Internet	0.43	0.49	*	*	0.07	0.66	*	*
Some Other Way	0.96	1.29	1.04	*	1.13	1.03	1.47	1.79

\*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.59P Source Where Methamphetamine was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From Friend or Relative for Free	0.7529		*		0.6871		0.9912	
Bought from Friend or Relative	0.9073		0.4873		0.8161		0.9262	
Took from Friend or Relative without Asking	0.1381		0.6004		0.7925		0.1112	
Bought from Drug Dealer or Other Stranger	0.4359		0.3853		0.3548		0.2006	
Bought on the Internet	0.9010		*		0.1588		*	
Some Other Way	0.3747		*		0.3498		0.3372	

\*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.60C Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Methamphetamine <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From Friend or Relative for Free	17	22	*	*	8	*	*	*
Bought from Friend or Relative	20	16	*	*	7	6	*	*
Took from Friend or Relative without Asking	2	6	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	17	10	*	*	7	4	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	5	3	*	*	3	3	*	*

\*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.60D Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Methamphetamine <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From Friend or Relative for Free	4.32	4.85	*	*	4.85	*	*	*
Bought from Friend or Relative	4.26	4.43	*	*	4.35	4.99	*	*
Took from Friend or Relative without Asking	0.70	1.79	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	4.27	3.04	*	*	4.56	3.87	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	1.33	1.01	*	*	2.38	2.66	*	*

\*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.60P Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source Where Friend or Relative Obtained Methamphetamine <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From Friend or Relative for Free	0.0803		*		*		*	
Bought from Friend or Relative	0.4871		*		0.9466		*	
Took from Friend or Relative without Asking	0.2402		*		*		*	
Bought from Drug Dealer or Other Stranger	0.0816		*		0.2786		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.5112		*		0.9508		*	

\*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.61C Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

<b>Source for Past Month Nonmedical Use among Past Month Methamphetamine Users<sup>1</sup></b>	<b>Total (2007-2008)</b>	<b>Total (2009-2010)</b>	<b>Aged 12-17 (2007-2008)</b>	<b>Aged 12-17 (2009-2010)</b>	<b>Aged 18-25 (2007-2008)</b>	<b>Aged 18-25 (2009-2010)</b>	<b>Aged 26+ (2007-2008)</b>	<b>Aged 26+ (2009-2010)</b>
From Friend or Relative for Free	*	23	*	*	*	*	*	*
Bought from Friend or Relative	*	26	*	*	5	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	21	*	*	*	*	*	*
Bought on the Internet	0	3	*	*	*	*	*	*
Some Other Way	7	9	*	*	2	*	*	*

\*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.61D Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

<b>Source for Past Month Nonmedical Use among Past Month Methamphetamine Users<sup>1</sup></b>	<b>Total (2007-2008)</b>	<b>Total (2009-2010)</b>	<b>Aged 12-17 (2007-2008)</b>	<b>Aged 12-17 (2009-2010)</b>	<b>Aged 18-25 (2007-2008)</b>	<b>Aged 18-25 (2009-2010)</b>	<b>Aged 26+ (2007-2008)</b>	<b>Aged 26+ (2009-2010)</b>
From Friend or Relative for Free	*	5.03	*	*	*	*	*	*
Bought from Friend or Relative	*	5.20	*	*	5.26	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	4.63	*	*	*	*	*	*
Bought on the Internet	0.03	0.77	*	*	*	*	*	*
Some Other Way	1.99	2.52	*	*	2.12	*	*	*

\*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.61P Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

<b>Source for Past Month Nonmedical Use among Past Month Methamphetamine Users<sup>1</sup></b>	<b>Total (2007-2008)</b>	<b>Total (2009-2010)</b>	<b>Aged 12-17 (2007-2008)</b>	<b>Aged 12-17 (2009-2010)</b>	<b>Aged 18-25 (2007-2008)</b>	<b>Aged 18-25 (2009-2010)</b>	<b>Aged 26+ (2007-2008)</b>	<b>Aged 26+ (2009-2010)</b>
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	0.1933		*		*		*	
Some Other Way	0.9353		*		*		*	

\*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.62C Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

<b>Source Where Friend or Relative Obtained Methamphetamine<sup>1</sup></b>	<b>Total (2007-2008)</b>	<b>Total (2009-2010)</b>	<b>Aged 12-17 (2007-2008)</b>	<b>Aged 12-17 (2009-2010)</b>	<b>Aged 18-25 (2007-2008)</b>	<b>Aged 18-25 (2009-2010)</b>	<b>Aged 26+ (2007-2008)</b>	<b>Aged 26+ (2009-2010)</b>
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	3	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.62D Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Methamphetamine <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	1.74	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.62P Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source Where Friend or Relative Obtained Methamphetamine <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

\*Low precision; no estimate reported.

NOTE: Estimates in this table include data from new methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.63C Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:  
Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

Source for Most Recent Nonmedical Use among Past Year Sedative Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	21	21	3	4	4	6	*	20
From More Than One Doctor	9	11	1	1	*	2	9	*
Wrote Fake Prescription	5	1	*	0	*	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	4	1	*	2	*	*	*
From Friend or Relative for Free	49	53	6	7	12	9	*	53
Bought from Friend or Relative	9	17	3	4	5	5	7	16
Took from Friend or Relative without Asking	6	11	3	3	3	*	3	9
Bought from Drug Dealer or Other Stranger	10	11	3	1	2	2	9	11
Bought on the Internet	2	*	*	*	0	*	*	*
Some Other Way	8	13	2	2	2	2	8	13

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.63D Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

Source for Most Recent Nonmedical Use among Past Year Sedative Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	3.04	2.51	3.05	4.07	2.21	3.46	*	3.36
From More Than One Doctor	1.27	1.39	1.73	1.05	*	1.59	1.94	*
Wrote Fake Prescription	0.73	0.07	*	0.44	*	0.29	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.63	0.47	0.62	*	1.41	*	*	*
From Friend or Relative for Free	3.98	3.83	5.09	5.37	4.01	5.03	*	5.14
Bought from Friend or Relative	1.26	2.03	2.91	3.70	2.70	3.19	1.52	2.71
Took from Friend or Relative without Asking	0.83	1.42	3.60	3.23	1.98	*	0.70	1.54
Bought from Drug Dealer or Other Stranger	1.47	1.38	2.83	1.14	1.34	1.16	2.12	1.93
Bought on the Internet	0.34	*	*	*	0.06	*	*	*
Some Other Way	1.22	1.58	1.91	1.99	1.08	1.16	1.84	2.22

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.63P Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:  
P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source for Most Recent Nonmedical Use among Past Year Sedative Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	0.8235		0.1570		0.0437		*	
From More Than One Doctor	0.8123		0.4771		*		*	
Wrote Fake Prescription	0.1331		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.4796		*		*		*	
From Friend or Relative for Free	0.5744		0.7283		0.0151		*	
Bought from Friend or Relative	0.1737		0.2390		0.7527		0.1836	
Took from Friend or Relative without Asking	0.2872		0.5987		*		0.4751	
Bought from Drug Dealer or Other Stranger	0.8975		0.1798		0.6618		0.8567	
Bought on the Internet	*		*		*		*	
Some Other Way	0.7867		0.8334		0.7553		0.8262	

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.64C Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Sedatives <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	43	51	*	*	9	*	*	*
From More Than One Doctor	4	3	*	*	1	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	3	2	*	*	*	*	*	*
From Friend or Relative for Free	6	11	*	*	3	*	*	*
Bought from Friend or Relative	2	2	*	*	*	*	*	*
Took from Friend or Relative without Asking	0	3	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	1	4	*	*	*	*	*	*
Bought on the Internet	0	*	*	*	0	*	*	*
Some Other Way	2	2	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.64D Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Sedatives <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	2.46	3.29	*	*	4.40	*	*	*
From More Than One Doctor	1.05	0.64	*	*	1.65	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.80	0.54	*	*	*	*	*	*
From Friend or Relative for Free	1.70	2.43	*	*	3.14	*	*	*
Bought from Friend or Relative	0.63	0.56	*	*	*	*	*	*
Took from Friend or Relative without Asking	0.13	0.66	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	0.31	1.01	*	*	*	*	*	*
Bought on the Internet	0.03	*	*	*	0.14	*	*	*
Some Other Way	0.49	0.43	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.64P Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: *P* Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source Where Friend or Relative Obtained Sedatives <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	0.6169		*		*		*	
From More Than One Doctor	0.2418		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7876		*		*		*	
From Friend or Relative for Free	0.9451		*		*		*	
Bought from Friend or Relative	0.9832		*		*		*	
Took from Friend or Relative without Asking	0.1730		*		*		*	
Bought from Drug Dealer or Other Stranger	0.1367		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.7538		*		*		*	

\*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>1</sup> Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.65C Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

Source for Past Month Nonmedical Use among Past Month Sedative Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	11	19	*	*	*	*	*	*
From More Than One Doctor	7	5	*	*	*	*	*	*
Wrote Fake Prescription	*	0	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	4	*	*	*	*	*	*
From Friend or Relative for Free	36	*	*	*	*	*	*	*
Bought from Friend or Relative	9	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	5	6	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	8	*	*	*	*	*	*	*
Bought on the Internet	1	0	*	*	*	*	*	*
Some Other Way	4	9	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.65D Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

Source for Past Month Nonmedical Use among Past Month Sedative Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	4.12	5.10	*	*	*	*	*	*
From More Than One Doctor	2.67	1.31	*	*	*	*	*	*
Wrote Fake Prescription	*	0.13	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	1.08	*	*	*	*	*	*
From Friend or Relative for Free	5.35	*	*	*	*	*	*	*
Bought from Friend or Relative	3.39	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	2.07	1.71	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	3.00	*	*	*	*	*	*	*
Bought on the Internet	0.24	0.03	*	*	*	*	*	*
Some Other Way	1.79	2.45	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.65P Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source for Past Month Nonmedical Use among Past Month Sedative Users <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	0.2742		*		*		*	
From More Than One Doctor	0.2509		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	0.2543		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	0.1620		*		*		*	
Some Other Way	0.5543		*		*		*	

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.66C Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Sedatives <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.66D Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010**

Source Where Friend or Relative Obtained Sedatives <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

**Table 6.66P Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008**

Source Where Friend or Relative Obtained Sedatives <sup>1</sup>	Total (2007-2008)	Total (2009-2010)	Aged 12-17 (2007-2008)	Aged 12-17 (2009-2010)	Aged 18-25 (2007-2008)	Aged 18-25 (2009-2010)	Aged 26+ (2007-2008)	Aged 26+ (2009-2010)
From One Doctor	*		*		*		*	
From More Than One Doctor	*		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

\*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

<sup>1</sup> Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.