

Table 6.71C Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

| Drug | Total ¹ | | PREGNANCY STATUS | | | |
|--|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| ILLCIT DRUGS² | 135 | 140 | 19 | 11 | 132 | 139 |
| Marijuana and Hashish | 108 | 117 | 17 | 10 | 106 | 117 |
| Cocaine | 37 | 37 | 6 | 2 | 36 | 37 |
| Crack | 16 | 23 | 6 | 1 | 15 | 23 |
| Heroin | 9 | 14 | * | 3 | 9 | 13 |
| Hallucinogens | 28 | 33 | 10 | 4 | 26 | 33 |
| LSD | 6 | 8 | 1 | 4 | 6 | 6 |
| PCP | 2 | 7 | 1 | 4 | 2 | 6 |
| Ecstasy | 23 | 27 | * | 4 | 21 | 26 |
| Inhalants | 17 | 19 | 3 | 4 | 16 | 19 |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 80 | 82 | 8 | 6 | 80 | 82 |
| Pain Relievers | 66 | 69 | 4 | 6 | 65 | 68 |
| OxyContin [®] | 15 | 20 | 1 | 2 | 15 | 20 |
| Tranquilizers | 47 | 43 | 2 | 5 | 47 | 42 |
| Stimulants ⁴ | 32 | 36 | 7 | 4 | 31 | 35 |
| Methamphetamine ⁴ | 21 | 22 | * | 4 | 20 | 22 |
| Sedatives | 14 | 12 | * | 4 | 14 | 11 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 93 | 94 | 14 | 7 | 91 | 93 |

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.71D Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010

| Drug | Total ¹ | | PREGNANCY STATUS | | | |
|--|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| ILLCIT DRUGS² | 0.21 | 0.21 | 0.73 | 0.47 | 0.21 | 0.22 |
| Marijuana and Hashish | 0.17 | 0.18 | 0.63 | 0.42 | 0.17 | 0.19 |
| Cocaine | 0.06 | 0.06 | 0.24 | 0.08 | 0.06 | 0.06 |
| Crack | 0.03 | 0.04 | 0.23 | 0.04 | 0.03 | 0.04 |
| Heroin | 0.01 | 0.02 | * | 0.10 | 0.01 | 0.02 |
| Hallucinogens | 0.05 | 0.05 | 0.39 | 0.17 | 0.04 | 0.06 |
| LSD | 0.01 | 0.01 | 0.02 | 0.16 | 0.01 | 0.01 |
| PCP | 0.00 | 0.01 | 0.02 | 0.16 | 0.00 | 0.01 |
| Ecstasy | 0.04 | 0.04 | * | 0.17 | 0.04 | 0.04 |
| Inhalants | 0.03 | 0.03 | 0.13 | 0.16 | 0.03 | 0.03 |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 0.13 | 0.13 | 0.31 | 0.25 | 0.13 | 0.14 |
| Pain Relievers | 0.10 | 0.11 | 0.16 | 0.23 | 0.11 | 0.11 |
| OxyContin [®] | 0.02 | 0.03 | 0.06 | 0.06 | 0.02 | 0.03 |
| Tranquilizers | 0.08 | 0.07 | 0.09 | 0.22 | 0.08 | 0.07 |
| Stimulants ⁴ | 0.05 | 0.06 | 0.26 | 0.17 | 0.05 | 0.06 |
| Methamphetamine ⁴ | 0.03 | 0.04 | * | 0.16 | 0.03 | 0.04 |
| Sedatives | 0.02 | 0.02 | * | 0.16 | 0.02 | 0.02 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 0.14 | 0.15 | 0.55 | 0.28 | 0.15 | 0.15 |

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.71P Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008

| Drug | Total ¹ | | PREGNANCY STATUS | | | |
|--|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| ILLCIT DRUGS² | 0.0006 | | 0.4355 | | 0.0004 | |
| Marijuana and Hashish | 0.0000 | | 0.8655 | | 0.0000 | |
| Cocaine | 0.0206 | | 0.3906 | | 0.0267 | |
| Crack | 0.8200 | | 0.3644 | | 0.6946 | |
| Heroin | 0.0643 | | * | | 0.1342 | |
| Hallucinogens | 0.0066 | | 0.7377 | | 0.0052 | |
| LSD | 0.2280 | | 0.3939 | | 0.4090 | |
| PCP | 0.0652 | | 0.3939 | | 0.0995 | |
| Ecstasy | 0.0112 | | * | | 0.0089 | |
| Inhalants | 0.8920 | | 0.9344 | | 0.9501 | |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 0.6316 | | 0.8815 | | 0.6588 | |
| Pain Relievers | 0.5607 | | 0.4543 | | 0.6263 | |
| OxyContin [®] | 0.0244 | | 0.7426 | | 0.0330 | |
| Tranquilizers | 0.5925 | | 0.2337 | | 0.5148 | |
| Stimulants ⁴ | 0.2243 | | 0.8884 | | 0.2533 | |
| Methamphetamine ⁴ | 0.6484 | | * | | 0.7709 | |
| Sedatives | 0.2682 | | * | | 0.1513 | |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 0.5117 | | 0.4169 | | 0.4777 | |

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.72C Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 135 | 140 | 19 | 11 | 132 | 139 |
| AGE | | | | | | |
| 15-17 | 29 | 30 | 4 | 3 | 28 | 29 |
| 18-25 | 58 | 56 | 9 | 8 | 65 | 65 |
| 26-44 | 111 | 118 | 17 | 8 | 109 | 118 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 127 | 130 | 19 | 10 | 124 | 129 |
| White | 112 | 112 | 14 | 8 | 110 | 112 |
| Black or African American | 54 | 56 | 12 | 5 | 51 | 55 |
| American Indian or Alaska Native | 6 | 6 | * | * | 6 | 6 |
| Native Hawaiian or Other Pacific Islander | * | 3 | * | * | * | 3 |
| Asian | 19 | 17 | * | * | 19 | 17 |
| Two or More Races | 14 | 14 | * | * | 14 | 14 |
| Hispanic or Latino | 40 | 53 | 4 | 6 | 40 | 52 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 12 | 9 | N/A | N/A |
| Second | N/A | N/A | 13 | 6 | N/A | N/A |
| Third | N/A | N/A | 8 | 4 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.72D Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 0.21 | 0.21 | 0.73 | 0.47 | 0.21 | 0.22 |
| AGE | | | | | | |
| 15-17 | 0.41 | 0.43 | 4.25 | 3.50 | 0.42 | 0.43 |
| 18-25 | 0.36 | 0.34 | 0.88 | 0.85 | 0.37 | 0.35 |
| 26-44 | 0.27 | 0.29 | 1.09 | 0.52 | 0.28 | 0.30 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.23 | 0.24 | 0.88 | 0.50 | 0.24 | 0.25 |
| White | 0.27 | 0.29 | 0.99 | 0.56 | 0.28 | 0.30 |
| Black or African American | 0.57 | 0.60 | 2.78 | 1.73 | 0.57 | 0.61 |
| American Indian or Alaska Native | 1.88 | 1.88 | * | * | 1.95 | 1.94 |
| Native Hawaiian or Other Pacific Islander | * | 1.28 | * | * | * | 1.30 |
| Asian | 0.56 | 0.53 | * | * | 0.59 | 0.56 |
| Two or More Races | 1.89 | 1.62 | * | * | 1.99 | 1.70 |
| Hispanic or Latino | 0.38 | 0.46 | 0.86 | 1.24 | 0.40 | 0.48 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 1.51 | 1.26 | N/A | N/A |
| Second | N/A | N/A | 1.34 | 0.64 | N/A | N/A |
| Third | N/A | N/A | 0.93 | 0.47 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.72P Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 0.0006 | | 0.4355 | | 0.0004 | |
| AGE | | | | | | |
| 15-17 | 0.3246 | | 0.3267 | | 0.2224 | |
| 18-25 | 0.0335 | | 0.7992 | | 0.0347 | |
| 26-44 | 0.0161 | | 0.3750 | | 0.0129 | |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.0101 | | 0.1806 | | 0.0060 | |
| White | 0.0229 | | 0.2159 | | 0.0125 | |
| Black or African American | 0.1752 | | 0.8086 | | 0.1711 | |
| American Indian or Alaska Native | 0.6275 | | * | | 0.7221 | |
| Native Hawaiian or Other Pacific Islander | * | | * | | * | |
| Asian | 0.8581 | | * | | 0.8959 | |
| Two or More Races | 0.6591 | | * | | 0.6386 | |
| Hispanic or Latino | 0.0013 | | 0.1725 | | 0.0018 | |
| TRIMESTER² | | | | | | |
| First | N/A | | 0.6376 | | N/A | |
| Second | N/A | | 0.3128 | | N/A | |
| Third | N/A | | 0.4644 | | N/A | |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.73C Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 108 | 117 | 17 | 10 | 106 | 117 |
| AGE | | | | | | |
| 15-17 | 25 | 26 | 4 | 3 | 24 | 25 |
| 18-25 | 52 | 52 | 7 | 8 | 57 | 59 |
| 26-44 | 86 | 98 | 15 | 6 | 84 | 98 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 103 | 110 | 17 | 9 | 100 | 110 |
| White | 91 | 95 | 12 | 8 | 90 | 95 |
| Black or African American | 47 | 49 | 12 | 5 | 44 | 48 |
| American Indian or Alaska Native | 5 | 5 | * | * | 5 | 5 |
| Native Hawaiian or Other Pacific Islander | 5 | 3 | * | * | 5 | 3 |
| Asian | 15 | 12 | * | * | 15 | 12 |
| Two or More Races | 14 | 13 | * | * | 13 | 13 |
| Hispanic or Latino | 31 | 40 | 3 | 5 | 31 | 40 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 12 | 8 | N/A | N/A |
| Second | N/A | N/A | 11 | 5 | N/A | N/A |
| Third | N/A | N/A | 3 | 3 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.73D Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 0.17 | 0.18 | 0.63 | 0.42 | 0.17 | 0.19 |
| AGE | | | | | | |
| 15-17 | 0.36 | 0.38 | 3.99 | 3.39 | 0.37 | 0.38 |
| 18-25 | 0.32 | 0.31 | 0.73 | 0.82 | 0.33 | 0.33 |
| 26-44 | 0.22 | 0.25 | 0.94 | 0.43 | 0.22 | 0.26 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.19 | 0.21 | 0.78 | 0.44 | 0.20 | 0.22 |
| White | 0.23 | 0.25 | 0.80 | 0.51 | 0.24 | 0.26 |
| Black or African American | 0.51 | 0.53 | 2.76 | 1.53 | 0.50 | 0.54 |
| American Indian or Alaska Native | 1.72 | 1.54 | * | * | 1.78 | 1.60 |
| Native Hawaiian or Other Pacific Islander | 1.75 | 1.21 | * | * | 1.78 | 1.23 |
| Asian | 0.45 | 0.37 | * | * | 0.48 | 0.39 |
| Two or More Races | 1.86 | 1.56 | * | * | 1.96 | 1.63 |
| Hispanic or Latino | 0.29 | 0.37 | 0.50 | 1.21 | 0.31 | 0.38 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 1.50 | 1.15 | N/A | N/A |
| Second | N/A | N/A | 1.18 | 0.58 | N/A | N/A |
| Third | N/A | N/A | 0.38 | 0.45 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.73P Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 0.0000 | | 0.8655 | | 0.0000 | |
| AGE | | | | | | |
| 15-17 | 0.2054 | | 0.6594 | | 0.1388 | |
| 18-25 | 0.0032 | | 0.3142 | | 0.0044 | |
| 26-44 | 0.0118 | | 0.4179 | | 0.0085 | |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.0005 | | 0.3395 | | 0.0002 | |
| White | 0.0017 | | 0.4986 | | 0.0010 | |
| Black or African American | 0.1127 | | 0.6851 | | 0.0778 | |
| American Indian or Alaska Native | 0.9187 | | * | | 0.9252 | |
| Native Hawaiian or Other Pacific Islander | 0.7801 | | * | | 0.7389 | |
| Asian | 0.5538 | | * | | 0.5278 | |
| Two or More Races | 0.7411 | | * | | 0.6926 | |
| Hispanic or Latino | 0.0086 | | 0.0312 | | 0.0137 | |
| TRIMESTER² | | | | | | |
| First | N/A | | 0.9777 | | N/A | |
| Second | N/A | | 0.5200 | | N/A | |
| Third | N/A | | 0.3941 | | N/A | |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.74C Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

| Substance | Total ¹ | | PREGNANCY STATUS | | | |
|-------------------------------------|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOBACCO PRODUCTS² | 266 | 249 | 33 | 32 | 260 | 245 |
| Cigarettes | 258 | 245 | 33 | 31 | 253 | 241 |
| Smokeless Tobacco | 28 | 31 | 2 | 4 | 28 | 31 |
| Cigars | 69 | 78 | 6 | 6 | 69 | 78 |
| Pipe Tobacco | 21 | 22 | 2 | 4 | 21 | 21 |
| ALCOHOL | 421 | 405 | 29 | 27 | 419 | 400 |
| Binge Alcohol Use ³ | 243 | 237 | 18 | 14 | 241 | 236 |
| Heavy Alcohol Use ³ | 100 | 103 | 6 | 6 | 100 | 102 |

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.74D Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010

| Substance | Total ¹ | | PREGNANCY STATUS | | | |
|-------------------------------------|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOBACCO PRODUCTS² | 0.36 | 0.35 | 1.20 | 1.18 | 0.37 | 0.36 |
| Cigarettes | 0.36 | 0.35 | 1.20 | 1.18 | 0.37 | 0.36 |
| Smokeless Tobacco | 0.05 | 0.05 | 0.07 | 0.17 | 0.05 | 0.05 |
| Cigars | 0.11 | 0.13 | 0.25 | 0.24 | 0.11 | 0.13 |
| Pipe Tobacco | 0.03 | 0.04 | 0.06 | 0.16 | 0.04 | 0.04 |
| ALCOHOL | 0.40 | 0.39 | 1.06 | 1.02 | 0.42 | 0.40 |
| Binge Alcohol Use ³ | 0.32 | 0.32 | 0.70 | 0.55 | 0.33 | 0.34 |
| Heavy Alcohol Use ³ | 0.16 | 0.16 | 0.23 | 0.25 | 0.16 | 0.17 |

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.74P Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: *P* Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008

| Substance | Total ¹ | | PREGNANCY STATUS | | | |
|-------------------------------------|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOBACCO PRODUCTS² | 0.3056 | | 0.9792 | | 0.3306 | |
| Cigarettes | 0.2145 | | 0.9686 | | 0.2433 | |
| Smokeless Tobacco | 0.0415 | | 0.2991 | | 0.0454 | |
| Cigars | 0.0338 | | 0.4416 | | 0.0378 | |
| Pipe Tobacco | 0.0436 | | 0.4531 | | 0.0764 | |
| ALCOHOL | 0.1449 | | 0.9073 | | 0.1943 | |
| Binge Alcohol Use ³ | 0.4455 | | 0.3335 | | 0.4354 | |
| Heavy Alcohol Use ³ | 0.9178 | | 0.7414 | | 0.9184 | |

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.75C Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 258 | 245 | 33 | 31 | 253 | 241 |
| AGE | | | | | | |
| 15-17 | 30 | 28 | 4 | 4 | 29 | 28 |
| 18-25 | 72 | 68 | 16 | 16 | 89 | 88 |
| 26-44 | 222 | 217 | 28 | 27 | 219 | 215 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 239 | 228 | 31 | 31 | 233 | 224 |
| White | 216 | 209 | 26 | 29 | 213 | 205 |
| Black or African American | 85 | 88 | 15 | 8 | 81 | 87 |
| American Indian or Alaska Native | 19 | 16 | * | * | 19 | 16 |
| Native Hawaiian or Other Pacific Islander | * | 7 | * | * | * | 6 |
| Asian | 38 | 39 | * | * | 37 | 39 |
| Two or More Races | 20 | 23 | * | * | 20 | 23 |
| Hispanic or Latino | 84 | 95 | 10 | 6 | 83 | 94 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 20 | 21 | N/A | N/A |
| Second | N/A | N/A | 20 | 17 | N/A | N/A |
| Third | N/A | N/A | 14 | 15 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.75D Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 0.36 | 0.35 | 1.20 | 1.18 | 0.37 | 0.36 |
| AGE | | | | | | |
| 15-17 | 0.43 | 0.41 | 3.84 | 4.28 | 0.43 | 0.42 |
| 18-25 | 0.44 | 0.41 | 1.51 | 1.47 | 0.46 | 0.43 |
| 26-44 | 0.50 | 0.51 | 1.72 | 1.75 | 0.52 | 0.52 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.39 | 0.40 | 1.37 | 1.39 | 0.40 | 0.41 |
| White | 0.45 | 0.47 | 1.66 | 1.67 | 0.46 | 0.49 |
| Black or African American | 0.82 | 0.86 | 3.36 | 2.57 | 0.83 | 0.88 |
| American Indian or Alaska Native | 3.86 | 3.74 | * | * | 3.98 | 3.92 |
| Native Hawaiian or Other Pacific Islander | * | 2.72 | * | * | * | 2.46 |
| Asian | 1.09 | 1.15 | * | * | 1.12 | 1.19 |
| Two or More Races | 2.49 | 2.38 | * | * | 2.58 | 2.48 |
| Hispanic or Latino | 0.72 | 0.77 | 1.88 | 1.35 | 0.76 | 0.81 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 2.34 | 2.47 | N/A | N/A |
| Second | N/A | N/A | 1.92 | 1.80 | N/A | N/A |
| Third | N/A | N/A | 1.66 | 1.88 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.75P Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 0.2145 | | 0.9686 | | 0.2433 | |
| AGE | | | | | | |
| 15-17 | 0.0275 | | 0.7145 | | 0.0285 | |
| 18-25 | 0.0839 | | 0.8012 | | 0.0720 | |
| 26-44 | 0.5310 | | 0.7550 | | 0.5964 | |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.0625 | | 0.8173 | | 0.0742 | |
| White | 0.1080 | | 0.7950 | | 0.1283 | |
| Black or African American | 0.7629 | | 0.3859 | | 0.9017 | |
| American Indian or Alaska Native | 0.0923 | | * | | 0.1097 | |
| Native Hawaiian or Other Pacific Islander | * | | * | | * | |
| Asian | 0.7319 | | * | | 0.9003 | |
| Two or More Races | 0.4125 | | * | | 0.4520 | |
| Hispanic or Latino | 0.1150 | | 0.8706 | | 0.0996 | |
| TRIMESTER² | | | | | | |
| First | N/A | | 0.3695 | | N/A | |
| Second | N/A | | 0.5328 | | N/A | |
| Third | N/A | | 0.8968 | | N/A | |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.76C Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 421 | 405 | 29 | 27 | 419 | 400 |
| AGE | | | | | | |
| 15-17 | 42 | 38 | 4 | 2 | 41 | 38 |
| 18-25 | 81 | 81 | 11 | 11 | 156 | 156 |
| 26-44 | 356 | 351 | 26 | 24 | 354 | 346 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 389 | 373 | 28 | 23 | 386 | 368 |
| White | 340 | 329 | 20 | 21 | 338 | 323 |
| Black or African American | 140 | 136 | 17 | 8 | 139 | 136 |
| American Indian or Alaska Native | 18 | 16 | * | * | 18 | 16 |
| Native Hawaiian or Other Pacific Islander | * | * | * | * | * | * |
| Asian | 77 | 90 | * | * | 78 | 89 |
| Two or More Races | 31 | 32 | * | * | 30 | 32 |
| Hispanic or Latino | 147 | 145 | 9 | 14 | 146 | 143 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 22 | 19 | N/A | N/A |
| Second | N/A | N/A | 15 | 17 | N/A | N/A |
| Third | N/A | N/A | 8 | 8 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.76D Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 0.40 | 0.39 | 1.06 | 1.02 | 0.42 | 0.40 |
| AGE | | | | | | |
| 15-17 | 0.57 | 0.51 | 3.78 | 2.98 | 0.57 | 0.52 |
| 18-25 | 0.50 | 0.48 | 1.09 | 1.13 | 0.50 | 0.49 |
| 26-44 | 0.56 | 0.56 | 1.59 | 1.60 | 0.58 | 0.58 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.43 | 0.40 | 1.24 | 1.08 | 0.44 | 0.41 |
| White | 0.47 | 0.46 | 1.34 | 1.28 | 0.48 | 0.47 |
| Black or African American | 1.02 | 0.97 | 3.79 | 2.36 | 1.07 | 1.00 |
| American Indian or Alaska Native | 3.84 | 3.61 | * | * | 3.94 | 3.81 |
| Native Hawaiian or Other Pacific Islander | * | * | * | * | * | * |
| Asian | 1.92 | 1.91 | * | * | 2.02 | 2.03 |
| Two or More Races | 2.83 | 2.60 | * | * | 2.80 | 2.66 |
| Hispanic or Latino | 0.94 | 0.98 | 1.79 | 2.90 | 0.97 | 1.02 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 2.51 | 2.30 | N/A | N/A |
| Second | N/A | N/A | 1.54 | 1.76 | N/A | N/A |
| Third | N/A | N/A | 0.92 | 0.99 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.76P Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 0.1449 | | 0.9073 | | 0.1943 | |
| AGE | | | | | | |
| 15-17 | 0.0143 | | 0.5100 | | 0.0191 | |
| 18-25 | 0.8082 | | 0.3574 | | 0.7542 | |
| 26-44 | 0.0635 | | 0.5320 | | 0.1000 | |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.1816 | | 0.6701 | | 0.1816 | |
| White | 0.4005 | | 0.9387 | | 0.2860 | |
| Black or African American | 0.1553 | | 0.1533 | | 0.2273 | |
| American Indian or Alaska Native | 0.0525 | | * | | 0.0696 | |
| Native Hawaiian or Other Pacific Islander | * | | * | | * | |
| Asian | 0.2049 | | * | | 0.2486 | |
| Two or More Races | 0.9240 | | * | | 0.8727 | |
| Hispanic or Latino | 0.2376 | | 0.3031 | | 0.3877 | |
| TRIMESTER² | | | | | | |
| First | N/A | | 0.8237 | | N/A | |
| Second | N/A | | 0.4575 | | N/A | |
| Third | N/A | | 0.8120 | | N/A | |

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.77C Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 243 | 237 | 18 | 14 | 241 | 236 |
| AGE | | | | | | |
| 15-17 | 31 | 29 | 3 | 2 | 31 | 29 |
| 18-25 | 76 | 75 | 9 | 6 | 111 | 105 |
| 26-44 | 199 | 202 | 15 | 12 | 198 | 202 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 224 | 211 | 18 | 11 | 222 | 210 |
| White | 200 | 191 | 10 | 10 | 199 | 190 |
| Black or African American | 73 | 78 | 13 | 4 | 72 | 78 |
| American Indian or Alaska Native | 14 | 12 | * | * | 14 | 12 |
| Native Hawaiian or Other Pacific Islander | * | 10 | * | * | * | 10 |
| Asian | 39 | 41 | * | * | 38 | 41 |
| Two or More Races | 18 | 20 | * | * | 18 | 20 |
| Hispanic or Latino | 92 | 103 | 6 | 9 | 92 | 102 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 15 | 13 | N/A | N/A |
| Second | N/A | N/A | 7 | 4 | N/A | N/A |
| Third | N/A | N/A | 4 | 1 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.77D Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 0.32 | 0.32 | 0.70 | 0.55 | 0.33 | 0.34 |
| AGE | | | | | | |
| 15-17 | 0.45 | 0.43 | 3.56 | 2.73 | 0.45 | 0.43 |
| 18-25 | 0.47 | 0.45 | 0.89 | 0.67 | 0.49 | 0.47 |
| 26-44 | 0.44 | 0.46 | 0.95 | 0.84 | 0.46 | 0.48 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.35 | 0.34 | 0.83 | 0.52 | 0.37 | 0.36 |
| White | 0.42 | 0.42 | 0.67 | 0.64 | 0.44 | 0.44 |
| Black or African American | 0.74 | 0.78 | 3.03 | 1.25 | 0.76 | 0.81 |
| American Indian or Alaska Native | 3.74 | 3.17 | * | * | 3.88 | 3.33 |
| Native Hawaiian or Other Pacific Islander | * | 3.81 | * | * | * | 3.88 |
| Asian | 1.13 | 1.23 | * | * | 1.18 | 1.27 |
| Two or More Races | 2.36 | 2.18 | * | * | 2.48 | 2.28 |
| Hispanic or Latino | 0.74 | 0.80 | 1.10 | 1.97 | 0.78 | 0.85 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 1.81 | 1.68 | N/A | N/A |
| Second | N/A | N/A | 0.71 | 0.45 | N/A | N/A |
| Third | N/A | N/A | 0.51 | 0.12 | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.77P Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 0.4455 | | 0.3335 | | 0.4354 | |
| AGE | | | | | | |
| 15-17 | 0.0266 | | 0.6414 | | 0.0324 | |
| 18-25 | 0.4135 | | 0.1623 | | 0.3583 | |
| 26-44 | 0.2206 | | 0.7774 | | 0.2121 | |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.6917 | | 0.1803 | | 0.5956 | |
| White | 0.8324 | | 0.8462 | | 0.6947 | |
| Black or African American | 0.2698 | | 0.1067 | | 0.2536 | |
| American Indian or Alaska Native | 0.0943 | | * | | 0.1241 | |
| Native Hawaiian or Other Pacific Islander | * | | * | | * | |
| Asian | 0.6942 | | * | | 0.6014 | |
| Two or More Races | 0.6519 | | * | | 0.6375 | |
| Hispanic or Latino | 0.2699 | | 0.6694 | | 0.3732 | |
| TRIMESTER² | | | | | | |
| First | N/A | | 0.9304 | | N/A | |
| Second | N/A | | 0.4126 | | N/A | |
| Third | N/A | | 0.0453 | | N/A | |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.78C Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2007-2008 and 2009-2010

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 100 | 103 | 6 | 6 | 100 | 102 |
| AGE | | | | | | |
| 15-17 | 15 | 12 | 1 | 1 | 15 | 12 |
| 18-25 | 43 | 42 | 3 | 5 | 49 | 46 |
| 26-44 | 84 | 91 | 5 | 4 | 83 | 90 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 94 | 97 | 6 | 5 | 93 | 96 |
| White | 87 | 87 | 6 | 4 | 87 | 87 |
| Black or African American | 26 | 33 | 2 | 2 | 25 | 33 |
| American Indian or Alaska Native | 7 | 6 | * | * | 7 | 6 |
| Native Hawaiian or Other Pacific Islander | 3 | 3 | * | * | 3 | 3 |
| Asian | 10 | 14 | * | * | 10 | 14 |
| Two or More Races | 9 | 6 | * | * | 8 | 6 |
| Hispanic or Latino | 33 | 33 | 1 | 4 | 33 | 33 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 6 | 6 | N/A | N/A |
| Second | N/A | N/A | 1 | 3 | N/A | N/A |
| Third | N/A | N/A | 1 | * | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.78D Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2007-2008 and 2009-2010

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 0.16 | 0.16 | 0.23 | 0.25 | 0.16 | 0.17 |
| AGE | | | | | | |
| 15-17 | 0.23 | 0.19 | 1.66 | 1.92 | 0.23 | 0.19 |
| 18-25 | 0.26 | 0.25 | 0.31 | 0.48 | 0.28 | 0.26 |
| 26-44 | 0.21 | 0.23 | 0.32 | 0.29 | 0.22 | 0.24 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.17 | 0.18 | 0.29 | 0.24 | 0.18 | 0.19 |
| White | 0.22 | 0.22 | 0.39 | 0.29 | 0.23 | 0.23 |
| Black or African American | 0.29 | 0.38 | 0.43 | 0.68 | 0.30 | 0.39 |
| American Indian or Alaska Native | 2.38 | 1.69 | * | * | 2.48 | 1.77 |
| Native Hawaiian or Other Pacific Islander | 1.08 | 1.12 | * | * | 1.10 | 1.14 |
| Asian | 0.30 | 0.41 | * | * | 0.32 | 0.43 |
| Two or More Races | 1.18 | 0.79 | * | * | 1.25 | 0.83 |
| Hispanic or Latino | 0.31 | 0.31 | 0.17 | 0.90 | 0.33 | 0.32 |
| TRIMESTER² | | | | | | |
| First | N/A | N/A | 0.73 | 0.76 | N/A | N/A |
| Second | N/A | N/A | 0.10 | 0.32 | N/A | N/A |
| Third | N/A | N/A | 0.17 | * | N/A | N/A |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.

Table 6.78P Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2009-2010 Versus 2007-2008

| Demographic/Pregnancy Characteristic | Total ¹ | | PREGNANCY STATUS | | | |
|---|--------------------|-----------|------------------|-----------|--------------|-----------|
| | | | Pregnant | | Not Pregnant | |
| | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 | 2007-2008 | 2009-2010 |
| TOTAL | 0.9178 | | 0.7414 | | 0.9184 | |
| AGE | | | | | | |
| 15-17 | 0.3065 | | 0.3327 | | 0.2840 | |
| 18-25 | 0.0439 | | 0.4656 | | 0.0275 | |
| 26-44 | 0.4274 | | 0.6377 | | 0.3807 | |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 0.9448 | | 0.7179 | | 0.9925 | |
| White | 0.7500 | | 0.5389 | | 0.8489 | |
| Black or African American | 0.3337 | | 0.4927 | | 0.2964 | |
| American Indian or Alaska Native | 0.9036 | | * | | 0.9340 | |
| Native Hawaiian or Other Pacific Islander | 0.5993 | | * | | 0.6500 | |
| Asian | 0.8049 | | * | | 0.8327 | |
| Two or More Races | 0.5248 | | * | | 0.5121 | |
| Hispanic or Latino | 0.9234 | | 0.2059 | | 0.9415 | |
| TRIMESTER² | | | | | | |
| First | N/A | | 0.7096 | | N/A | |
| Second | N/A | | 0.3436 | | N/A | |
| Third | N/A | | * | | N/A | |

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2007-2010.