

**Table 7.1A Types of Illicit Drug Use in Lifetime among Persons Aged 12 or Older: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	108,255 <sup>b</sup>	110,205 <sup>b</sup>	110,057 <sup>b</sup>	112,085 <sup>b</sup>	111,774 <sup>b</sup>	114,275 <sup>b</sup>	117,325	118,705	119,508
Marijuana and Hashish	94,946 <sup>b</sup>	96,611 <sup>b</sup>	96,772 <sup>b</sup>	97,545 <sup>b</sup>	97,825 <sup>b</sup>	100,518 <sup>b</sup>	102,404 <sup>b</sup>	104,446	106,232
Cocaine	33,910 <sup>b</sup>	34,891 <sup>b</sup>	34,153 <sup>b</sup>	33,673 <sup>b</sup>	35,298 <sup>a</sup>	35,882	36,773	36,599	37,210
Crack	8,402	7,949 <sup>b</sup>	7,840 <sup>b</sup>	7,928 <sup>a</sup>	8,554	8,581	8,445	8,359	9,158
Heroin	3,668	3,744	3,145 <sup>b</sup>	3,534	3,785	3,780	3,788	3,683	4,126
Hallucinogens	34,314 <sup>b</sup>	34,363 <sup>b</sup>	34,333 <sup>b</sup>	33,728 <sup>b</sup>	35,281 <sup>a</sup>	34,215 <sup>b</sup>	35,963	37,256	37,476
LSD	24,516	24,424	23,398	22,433	23,346	22,656	23,547	23,635	23,320
PCP	7,418 <sup>b</sup>	7,107 <sup>a</sup>	6,762	6,603	6,618	6,140	6,631	6,239	6,258
Ecstasy	10,150 <sup>b</sup>	10,904 <sup>b</sup>	11,130 <sup>b</sup>	11,495 <sup>b</sup>	12,262 <sup>b</sup>	12,426 <sup>b</sup>	12,924 <sup>b</sup>	14,234 <sup>b</sup>	15,901
Inhalants	22,870	22,995 <sup>a</sup>	22,798	22,745	22,879	22,477	22,274	22,448	21,660
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	47,958 <sup>b</sup>	49,001 <sup>b</sup>	49,157 <sup>a</sup>	49,571 <sup>a</sup>	50,965	50,415	51,970	51,771	51,641
Pain Relievers	29,611 <sup>b</sup>	31,207 <sup>b</sup>	31,768 <sup>b</sup>	32,692 <sup>a</sup>	33,472	33,060 <sup>a</sup>	34,861	35,046	34,776
OxyContin <sup>®</sup>	1,924 <sup>b</sup>	2,832 <sup>b</sup>	3,072 <sup>b</sup>	3,481 <sup>b</sup>	4,098 <sup>b</sup>	4,354 <sup>b</sup>	4,842 <sup>b</sup>	5,829	6,121
Tranquilizers	19,267 <sup>b</sup>	20,220 <sup>b</sup>	19,852 <sup>b</sup>	21,041	21,303	20,208 <sup>b</sup>	21,476	21,755	22,103
Stimulants <sup>3</sup>	23,496 <sup>a</sup>	23,004	22,297	20,983	22,468	21,654	21,206	21,930	21,660
Methamphetamine <sup>3</sup>	15,365 <sup>b</sup>	15,139 <sup>b</sup>	14,512 <sup>a</sup>	12,663	14,206	13,065	12,598	12,837	13,012
Sedatives	9,960 <sup>b</sup>	9,510 <sup>b</sup>	9,891 <sup>b</sup>	8,982 <sup>b</sup>	8,822 <sup>a</sup>	8,396	8,882 <sup>a</sup>	8,605 <sup>a</sup>	7,631
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	70,300 <sup>b</sup>	71,128 <sup>b</sup>	70,657 <sup>b</sup>	71,822 <sup>b</sup>	72,906 <sup>b</sup>	73,494 <sup>a</sup>	75,573	75,780	76,203

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.1B Types of Illicit Drug Use in Lifetime among Persons Aged 12 or Older: Percentages, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLICIT DRUGS<sup>1</sup></b>	46.0	46.4	45.8 <sup>a</sup>	46.1	45.4 <sup>b</sup>	46.1	47.0	47.1	47.1
Marijuana and Hashish	40.4 <sup>b</sup>	40.6 <sup>a</sup>	40.2 <sup>b</sup>	40.1 <sup>b</sup>	39.8 <sup>b</sup>	40.6 <sup>a</sup>	41.0	41.5	41.9
Cocaine	14.4	14.7	14.2	13.8 <sup>a</sup>	14.3	14.5	14.7	14.5	14.7
Crack	3.6	3.3	3.3	3.3	3.5	3.5	3.4	3.3	3.6
Heroin	1.6	1.6	1.3 <sup>a</sup>	1.5	1.5	1.5	1.5	1.5	1.6
Hallucinogens	14.6	14.5	14.3	13.9 <sup>a</sup>	14.3	13.8 <sup>b</sup>	14.4	14.8	14.8
LSD	10.4 <sup>b</sup>	10.3 <sup>b</sup>	9.7	9.2	9.5	9.1	9.4	9.4	9.2
PCP	3.2 <sup>b</sup>	3.0 <sup>b</sup>	2.8	2.7	2.7	2.5	2.7	2.5	2.5
Ecstasy	4.3 <sup>b</sup>	4.6 <sup>b</sup>	4.6 <sup>b</sup>	4.7 <sup>b</sup>	5.0 <sup>b</sup>	5.0 <sup>b</sup>	5.2 <sup>b</sup>	5.7 <sup>b</sup>	6.3
Inhalants	9.7 <sup>b</sup>	9.7 <sup>b</sup>	9.5 <sup>b</sup>	9.4 <sup>b</sup>	9.3 <sup>b</sup>	9.1	8.9	8.9	8.5
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	20.4	20.6	20.4	20.4	20.7	20.3	20.8	20.6	20.4
Pain Relievers	12.6 <sup>b</sup>	13.1	13.2	13.4	13.6	13.3	14.0	13.9	13.7
OxyContin <sup>®</sup>	0.8 <sup>b</sup>	1.2 <sup>b</sup>	1.3 <sup>b</sup>	1.4 <sup>b</sup>	1.7 <sup>b</sup>	1.8 <sup>b</sup>	1.9 <sup>b</sup>	2.3	2.4
Tranquilizers	8.2	8.5	8.3	8.7	8.7	8.2 <sup>a</sup>	8.6	8.6	8.7
Stimulants <sup>3</sup>	10.0 <sup>b</sup>	9.7 <sup>b</sup>	9.3 <sup>a</sup>	8.6	9.1 <sup>a</sup>	8.7	8.5	8.7	8.5
Methamphetamine <sup>3</sup>	6.5 <sup>b</sup>	6.4 <sup>b</sup>	6.0 <sup>b</sup>	5.2	5.8 <sup>a</sup>	5.3	5.0	5.1	5.1
Sedatives	4.2 <sup>b</sup>	4.0 <sup>b</sup>	4.1 <sup>b</sup>	3.7 <sup>b</sup>	3.6 <sup>b</sup>	3.4	3.6 <sup>a</sup>	3.4 <sup>a</sup>	3.0
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	29.9	29.9	29.4	29.5	29.6	29.7	30.3	30.1	30.0

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.2A Types of Illicit Drug Use in the Past Year among Persons Aged 12 or Older: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	35,132 <sup>b</sup>	34,993 <sup>b</sup>	34,807 <sup>b</sup>	35,041 <sup>b</sup>	35,775 <sup>b</sup>	35,692 <sup>b</sup>	35,525 <sup>b</sup>	37,954	38,806
Marijuana and Hashish	25,755 <sup>b</sup>	25,231 <sup>b</sup>	25,451 <sup>b</sup>	25,375 <sup>b</sup>	25,378 <sup>b</sup>	25,085 <sup>b</sup>	25,768 <sup>b</sup>	28,521	29,206
Cocaine	5,902 <sup>b</sup>	5,908 <sup>b</sup>	5,658 <sup>b</sup>	5,523 <sup>b</sup>	6,069 <sup>b</sup>	5,738 <sup>b</sup>	5,255 <sup>b</sup>	4,797	4,499
Crack	1,554 <sup>b</sup>	1,406 <sup>b</sup>	1,304 <sup>b</sup>	1,381 <sup>b</sup>	1,479 <sup>b</sup>	1,451 <sup>b</sup>	1,109	1,016	871
Heroin	404 <sup>a</sup>	314 <sup>b</sup>	398 <sup>a</sup>	379 <sup>b</sup>	560	366 <sup>b</sup>	453	605	618
Hallucinogens	4,749	3,936 <sup>a</sup>	3,878 <sup>b</sup>	3,809 <sup>b</sup>	3,956 <sup>a</sup>	3,762 <sup>b</sup>	3,678 <sup>b</sup>	4,509	4,517
LSD	999	558 <sup>b</sup>	592 <sup>b</sup>	563 <sup>b</sup>	666 <sup>a</sup>	620 <sup>b</sup>	802	779	874
PCP	235 <sup>b</sup>	219 <sup>b</sup>	210 <sup>b</sup>	164 <sup>a</sup>	187 <sup>a</sup>	137	99	122	95
Ecstasy	3,167 <sup>b</sup>	2,119 <sup>b</sup>	1,915 <sup>b</sup>	1,960 <sup>b</sup>	2,130 <sup>b</sup>	2,132 <sup>b</sup>	2,139 <sup>b</sup>	2,799	2,645
Inhalants	2,084	2,075	2,255	2,187	2,218	2,080	2,047	2,090	2,030
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	14,795 <sup>a</sup>	15,163	14,849 <sup>a</sup>	15,346	16,482	16,280	15,166	16,006	16,031
Pain Relievers	10,992 <sup>b</sup>	11,671	11,256 <sup>a</sup>	11,815	12,649	12,466	11,885	12,405	12,213
OxyContin <sup>®</sup>	--	--	1,213 <sup>b</sup>	1,226 <sup>b</sup>	1,323 <sup>b</sup>	1,422 <sup>b</sup>	1,459 <sup>b</sup>	1,677	1,869
Tranquilizers	4,849 <sup>a</sup>	5,051	5,068	5,249	5,058	5,282	5,103	5,460	5,581
Stimulants <sup>3</sup>	3,380 <sup>a</sup>	3,031	3,254	3,088	3,791 <sup>b</sup>	2,998	2,639	3,060	2,887
Methamphetamine <sup>3</sup>	1,755 <sup>b</sup>	1,602 <sup>b</sup>	1,808 <sup>b</sup>	1,603 <sup>b</sup>	1,889 <sup>b</sup>	1,343 <sup>a</sup>	850	1,165	954
Sedatives	981	831	737	750	926	864	621 <sup>a</sup>	811	907
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	20,423	20,305	19,658	20,109	21,254	21,144	19,990	21,000	20,576

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.2B Types of Illicit Drug Use in the Past Year among Persons Aged 12 or Older: Percentages, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	14.9	14.7	14.5 <sup>a</sup>	14.4 <sup>b</sup>	14.5 <sup>a</sup>	14.4 <sup>b</sup>	14.2 <sup>b</sup>	15.1	15.3
Marijuana and Hashish	11.0	10.6 <sup>b</sup>	10.6 <sup>b</sup>	10.4 <sup>b</sup>	10.3 <sup>b</sup>	10.1 <sup>b</sup>	10.3 <sup>b</sup>	11.3	11.5
Cocaine	2.5 <sup>b</sup>	2.5 <sup>b</sup>	2.4 <sup>b</sup>	2.3 <sup>b</sup>	2.5 <sup>b</sup>	2.3 <sup>b</sup>	2.1 <sup>b</sup>	1.9	1.8
Crack	0.7 <sup>b</sup>	0.6 <sup>b</sup>	0.5 <sup>b</sup>	0.6 <sup>b</sup>	0.6 <sup>b</sup>	0.6 <sup>b</sup>	0.4	0.4	0.3
Heroin	0.2	0.1 <sup>b</sup>	0.2 <sup>a</sup>	0.2 <sup>a</sup>	0.2	0.1 <sup>b</sup>	0.2	0.2	0.2
Hallucinogens	2.0 <sup>a</sup>	1.7	1.6	1.6 <sup>a</sup>	1.6	1.5 <sup>b</sup>	1.5 <sup>b</sup>	1.8	1.8
LSD	0.4 <sup>a</sup>	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.3 <sup>a</sup>	0.3 <sup>b</sup>	0.3	0.3	0.3
PCP	0.1 <sup>b</sup>	0.1 <sup>b</sup>	0.1 <sup>b</sup>	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.1	0.0	0.0	0.0
Ecstasy	1.3 <sup>b</sup>	0.9 <sup>a</sup>	0.8 <sup>b</sup>	0.8 <sup>b</sup>	0.9 <sup>b</sup>	0.9 <sup>b</sup>	0.9 <sup>b</sup>	1.1	1.0
Inhalants	0.9	0.9	0.9 <sup>a</sup>	0.9	0.9	0.8	0.8	0.8	0.8
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	6.3	6.4	6.2	6.3	6.7	6.6	6.1	6.4	6.3
Pain Relievers	4.7	4.9	4.7	4.9	5.1	5.0	4.8	4.9	4.8
OxyContin <sup>®</sup>	--	--	0.5 <sup>b</sup>	0.5 <sup>b</sup>	0.5 <sup>b</sup>	0.6 <sup>b</sup>	0.6 <sup>b</sup>	0.7	0.7
Tranquilizers	2.1	2.1	2.1	2.2	2.1	2.1	2.0	2.2	2.2
Stimulants <sup>3</sup>	1.4 <sup>b</sup>	1.3	1.4 <sup>a</sup>	1.3	1.5 <sup>b</sup>	1.2	1.1	1.2	1.1
Methamphetamine <sup>3</sup>	0.7 <sup>b</sup>	0.7 <sup>b</sup>	0.8 <sup>b</sup>	0.7 <sup>b</sup>	0.8 <sup>b</sup>	0.5 <sup>b</sup>	0.3	0.5	0.4
Sedatives	0.4	0.3	0.3	0.3	0.4	0.3	0.2 <sup>a</sup>	0.3	0.4
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	8.7 <sup>a</sup>	8.5	8.2	8.3	8.6 <sup>a</sup>	8.5	8.0	8.3	8.1

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.3A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	19,522 <sup>b</sup>	19,470 <sup>b</sup>	19,071 <sup>b</sup>	19,720 <sup>b</sup>	20,357 <sup>b</sup>	19,857 <sup>b</sup>	20,077 <sup>b</sup>	21,813	22,622
Marijuana and Hashish	14,584 <sup>b</sup>	14,638 <sup>b</sup>	14,576 <sup>b</sup>	14,626 <sup>b</sup>	14,813 <sup>b</sup>	14,448 <sup>b</sup>	15,203 <sup>b</sup>	16,718	17,373
Cocaine	2,020 <sup>b</sup>	2,281 <sup>b</sup>	2,021 <sup>b</sup>	2,397 <sup>b</sup>	2,421 <sup>b</sup>	2,075 <sup>b</sup>	1,855 <sup>a</sup>	1,637	1,466
Crack	567 <sup>a</sup>	604 <sup>a</sup>	467	682 <sup>b</sup>	702 <sup>b</sup>	610 <sup>a</sup>	359	492	378
Heroin	166	119 <sup>a</sup>	166	136 <sup>a</sup>	338	153	213	195	239
Hallucinogens	1,196	1,042	929 <sup>b</sup>	1,088	1,006	996	1,060	1,258	1,208
LSD	112	133	141	104	130	145	154	158	154
PCP	58	56	49	48	30	41	24	53	35
Ecstasy	676	470 <sup>b</sup>	450 <sup>b</sup>	502 <sup>a</sup>	528 <sup>a</sup>	503 <sup>a</sup>	555	760	695
Inhalants	635	570	638	611	761	616	640	560	685
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	6,287	6,451	6,110 <sup>a</sup>	6,491	7,095	6,895	6,224 <sup>a</sup>	6,953	6,967
Pain Relievers	4,377 <sup>a</sup>	4,693	4,404 <sup>a</sup>	4,658	5,220	5,174	4,747	5,257	5,100
OxyContin <sup>®</sup>	--	--	325 <sup>b</sup>	334 <sup>b</sup>	276 <sup>b</sup>	369 <sup>a</sup>	435	510	564
Tranquilizers	1,804	1,830	1,616 <sup>b</sup>	1,817	1,766 <sup>a</sup>	1,835	1,800 <sup>a</sup>	2,010	2,160
Stimulants <sup>3</sup>	1,303	1,310	1,312	1,188	1,385 <sup>a</sup>	1,053	904	1,290	1,077
Methamphetamine <sup>3</sup>	683 <sup>b</sup>	726 <sup>b</sup>	706 <sup>b</sup>	628 <sup>b</sup>	731 <sup>b</sup>	529	314	502	353
Sedatives	436	294	265	272	385	346	234	370	374
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	8,777	8,849	8,247	8,963	9,615	9,270	8,565	9,157	9,017

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.3B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older: Percentages, 2002-2010**

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	8.3 <sup>a</sup>	8.2 <sup>b</sup>	7.9 <sup>b</sup>	8.1 <sup>b</sup>	8.3 <sup>a</sup>	8.0 <sup>b</sup>	8.0 <sup>b</sup>	8.7	8.9
Marijuana and Hashish	6.2 <sup>b</sup>	6.2 <sup>b</sup>	6.1 <sup>b</sup>	6.0 <sup>b</sup>	6.0 <sup>b</sup>	5.8 <sup>b</sup>	6.1 <sup>b</sup>	6.6	6.9
Cocaine	0.9 <sup>b</sup>	1.0 <sup>b</sup>	0.8 <sup>b</sup>	1.0 <sup>b</sup>	1.0 <sup>b</sup>	0.8 <sup>b</sup>	0.7 <sup>a</sup>	0.7	0.6
Crack	0.2 <sup>a</sup>	0.3 <sup>a</sup>	0.2	0.3 <sup>b</sup>	0.3 <sup>b</sup>	0.2 <sup>a</sup>	0.1	0.2	0.1
Heroin	0.1	0.1 <sup>a</sup>	0.1	0.1 <sup>a</sup>	0.1	0.1	0.1	0.1	0.1
Hallucinogens	0.5	0.4	0.4 <sup>a</sup>	0.4	0.4	0.4	0.4	0.5	0.5
LSD	0.0	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1
PCP	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Ecstasy	0.3	0.2 <sup>a</sup>	0.2 <sup>b</sup>	0.2 <sup>a</sup>	0.2	0.2 <sup>a</sup>	0.2	0.3	0.3
Inhalants	0.3	0.2	0.3	0.3	0.3	0.2	0.3	0.2	0.3
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	2.7	2.7	2.5	2.7	2.9	2.8	2.5	2.8	2.7
Pain Relievers	1.9	2.0	1.8	1.9	2.1	2.1	1.9	2.1	2.0
OxyContin <sup>®</sup>	--	--	0.1 <sup>b</sup>	0.1 <sup>b</sup>	0.1 <sup>b</sup>	0.1 <sup>a</sup>	0.2	0.2	0.2
Tranquilizers	0.8	0.8	0.7 <sup>b</sup>	0.7	0.7	0.7	0.7	0.8	0.9
Stimulants <sup>3</sup>	0.6 <sup>a</sup>	0.6 <sup>a</sup>	0.5 <sup>a</sup>	0.5	0.6 <sup>a</sup>	0.4	0.4	0.5	0.4
Methamphetamine <sup>3</sup>	0.3 <sup>b</sup>	0.2	0.1	0.2	0.1				
Sedatives	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	3.7	3.7	3.4	3.7	3.9 <sup>a</sup>	3.7	3.4	3.6	3.6

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.4A Types of Illicit Drug Use in Lifetime among Persons Aged 12 to 17: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	7,642 <sup>b</sup>	7,631 <sup>b</sup>	7,566 <sup>b</sup>	7,022 <sup>b</sup>	7,009 <sup>b</sup>	6,621 <sup>a</sup>	6,520	6,589 <sup>a</sup>	6,251
Marijuana and Hashish	5,104 <sup>b</sup>	4,888 <sup>b</sup>	4,788 <sup>b</sup>	4,402 <sup>a</sup>	4,394 <sup>a</sup>	4,084	4,108	4,180	4,139
Cocaine	670 <sup>b</sup>	648 <sup>b</sup>	611 <sup>b</sup>	582 <sup>b</sup>	570 <sup>b</sup>	542 <sup>b</sup>	472 <sup>b</sup>	402	362
Crack	163 <sup>b</sup>	157 <sup>b</sup>	125 <sup>b</sup>	108 <sup>b</sup>	128 <sup>b</sup>	106 <sup>b</sup>	79	58	60
Heroin	94 <sup>b</sup>	80 <sup>a</sup>	78 <sup>a</sup>	60	60	54	68	56	46
Hallucinogens	1,419 <sup>b</sup>	1,250 <sup>b</sup>	1,160 <sup>a</sup>	998	986	896	973	1,012	999
LSD	670 <sup>b</sup>	397 <sup>b</sup>	313 <sup>a</sup>	288	218	200	268	249	231
PCP	233 <sup>b</sup>	204 <sup>b</sup>	175 <sup>b</sup>	182 <sup>b</sup>	128 <sup>b</sup>	104	102	99	76
Ecstasy	820 <sup>b</sup>	603	523 <sup>a</sup>	399 <sup>b</sup>	474 <sup>b</sup>	446 <sup>b</sup>	531	572	619
Inhalants	2,605 <sup>b</sup>	2,670 <sup>b</sup>	2,762 <sup>b</sup>	2,658 <sup>b</sup>	2,561 <sup>b</sup>	2,422 <sup>b</sup>	2,307 <sup>b</sup>	2,267 <sup>b</sup>	2,005
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	3,402 <sup>b</sup>	3,372 <sup>b</sup>	3,403 <sup>b</sup>	3,019 <sup>b</sup>	3,171 <sup>b</sup>	2,949 <sup>b</sup>	2,772	2,796 <sup>a</sup>	2,583
Pain Relievers	2,775 <sup>b</sup>	2,798 <sup>b</sup>	2,879 <sup>b</sup>	2,517 <sup>b</sup>	2,647 <sup>b</sup>	2,452 <sup>a</sup>	2,346	2,378	2,238
OxyContin <sup>®</sup>	224 <sup>b</sup>	254 <sup>b</sup>	305	276 <sup>a</sup>	342	360	341	378	366
Tranquilizers	834 <sup>a</sup>	874 <sup>b</sup>	816	766	740	748	718	693	722
Stimulants <sup>3</sup>	1,112 <sup>b</sup>	1,046 <sup>b</sup>	906 <sup>b</sup>	870 <sup>b</sup>	930 <sup>b</sup>	732 <sup>b</sup>	670 <sup>b</sup>	582	529
Methamphetamine <sup>3</sup>	416 <sup>b</sup>	382 <sup>b</sup>	346 <sup>b</sup>	319 <sup>b</sup>	341 <sup>b</sup>	222 <sup>a</sup>	195	185	157
Sedatives	238 <sup>a</sup>	249 <sup>b</sup>	251 <sup>b</sup>	232 <sup>a</sup>	224	214	203	224	177
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	5,286 <sup>b</sup>	5,319 <sup>b</sup>	5,355 <sup>b</sup>	4,944 <sup>b</sup>	4,949 <sup>b</sup>	4,637 <sup>b</sup>	4,488 <sup>b</sup>	4,474 <sup>b</sup>	4,151

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.4B Types of Illicit Drug Use in Lifetime among Persons Aged 12 to 17: Percentages, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	30.9 <sup>b</sup>	30.5 <sup>b</sup>	30.0 <sup>b</sup>	27.7 <sup>b</sup>	27.6 <sup>b</sup>	26.2	26.2	26.8	25.7
Marijuana and Hashish	20.6 <sup>b</sup>	19.6 <sup>b</sup>	19.0 <sup>b</sup>	17.4	17.3	16.2	16.5	17.0	17.0
Cocaine	2.7 <sup>b</sup>	2.6 <sup>b</sup>	2.4 <sup>b</sup>	2.3 <sup>b</sup>	2.2 <sup>b</sup>	2.1 <sup>b</sup>	1.9 <sup>a</sup>	1.6	1.5
Crack	0.7 <sup>b</sup>	0.6 <sup>b</sup>	0.5 <sup>b</sup>	0.4 <sup>a</sup>	0.5 <sup>b</sup>	0.4 <sup>b</sup>	0.3	0.2	0.2
Heroin	0.4 <sup>b</sup>	0.3 <sup>a</sup>	0.3 <sup>a</sup>	0.2	0.2	0.2	0.3	0.2	0.2
Hallucinogens	5.7 <sup>b</sup>	5.0 <sup>b</sup>	4.6	3.9	3.9	3.5 <sup>a</sup>	3.9	4.1	4.1
LSD	2.7 <sup>b</sup>	1.6 <sup>b</sup>	1.2 <sup>a</sup>	1.1	0.9	0.8	1.1	1.0	0.9
PCP	0.9 <sup>b</sup>	0.8 <sup>b</sup>	0.7 <sup>b</sup>	0.7 <sup>b</sup>	0.5 <sup>a</sup>	0.4	0.4	0.4	0.3
Ecstasy	3.3 <sup>b</sup>	2.4	2.1 <sup>a</sup>	1.6 <sup>b</sup>	1.9 <sup>b</sup>	1.8 <sup>b</sup>	2.1 <sup>a</sup>	2.3	2.5
Inhalants	10.5 <sup>b</sup>	10.7 <sup>b</sup>	11.0 <sup>b</sup>	10.5 <sup>b</sup>	10.1 <sup>b</sup>	9.6 <sup>b</sup>	9.3 <sup>b</sup>	9.2 <sup>b</sup>	8.2
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	13.7 <sup>b</sup>	13.5 <sup>b</sup>	13.5 <sup>b</sup>	11.9 <sup>b</sup>	12.5 <sup>b</sup>	11.7 <sup>b</sup>	11.1	11.4 <sup>a</sup>	10.6
Pain Relievers	11.2 <sup>b</sup>	11.2 <sup>b</sup>	11.4 <sup>b</sup>	9.9 <sup>a</sup>	10.4 <sup>b</sup>	9.7	9.4	9.7	9.2
OxyContin <sup>®</sup>	0.9 <sup>b</sup>	1.0 <sup>b</sup>	1.2 <sup>a</sup>	1.1 <sup>b</sup>	1.3	1.4	1.4	1.5	1.5
Tranquilizers	3.4	3.5 <sup>a</sup>	3.2	3.0	2.9	3.0	2.9	2.8	3.0
Stimulants <sup>3</sup>	4.5 <sup>b</sup>	4.2 <sup>b</sup>	3.6 <sup>b</sup>	3.4 <sup>b</sup>	3.7 <sup>b</sup>	2.9 <sup>b</sup>	2.7 <sup>a</sup>	2.4	2.2
Methamphetamine <sup>3</sup>	1.7 <sup>b</sup>	1.5 <sup>b</sup>	1.4 <sup>b</sup>	1.3 <sup>b</sup>	1.3 <sup>b</sup>	0.9 <sup>a</sup>	0.8	0.8	0.6
Sedatives	1.0 <sup>a</sup>	1.0 <sup>a</sup>	1.0 <sup>a</sup>	0.9	0.9	0.8	0.8	0.9	0.7
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	21.4 <sup>b</sup>	21.3 <sup>b</sup>	21.2 <sup>b</sup>	19.5 <sup>b</sup>	19.5 <sup>b</sup>	18.4 <sup>b</sup>	18.0 <sup>a</sup>	18.2 <sup>a</sup>	17.0

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.5A Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	5,494 <sup>b</sup>	5,455 <sup>b</sup>	5,300 <sup>b</sup>	5,041 <sup>a</sup>	4,971	4,726	4,726	4,809	4,730
Marijuana and Hashish	3,905 <sup>b</sup>	3,745 <sup>b</sup>	3,656 <sup>a</sup>	3,383	3,348	3,147 <sup>a</sup>	3,238	3,339	3,400
Cocaine	508 <sup>b</sup>	438 <sup>b</sup>	411 <sup>b</sup>	423 <sup>b</sup>	411 <sup>b</sup>	383 <sup>b</sup>	305 <sup>a</sup>	240	239
Crack	100 <sup>b</sup>	95 <sup>b</sup>	63 <sup>b</sup>	54 <sup>b</sup>	73 <sup>b</sup>	65 <sup>b</sup>	33	28	18
Heroin	51	36	48	37	37	23	42	34	28
Hallucinogens	945 <sup>b</sup>	778	753	665	650	627 <sup>a</sup>	690	735	747
LSD	314 <sup>b</sup>	149	157	150	106	125	183	142	135
PCP	110 <sup>b</sup>	89 <sup>b</sup>	72 <sup>a</sup>	85 <sup>b</sup>	59	55	61 <sup>a</sup>	47	35
Ecstasy	547	329 <sup>b</sup>	310 <sup>b</sup>	256 <sup>b</sup>	306 <sup>b</sup>	329 <sup>b</sup>	353 <sup>b</sup>	416	466
Inhalants	1,078 <sup>b</sup>	1,119 <sup>b</sup>	1,168 <sup>b</sup>	1,144 <sup>b</sup>	1,110 <sup>b</sup>	994	978	956	881
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	2,285 <sup>b</sup>	2,295 <sup>b</sup>	2,228 <sup>b</sup>	2,101 <sup>b</sup>	2,181 <sup>b</sup>	2,034 <sup>b</sup>	1,913	1,885	1,812
Pain Relievers	1,870 <sup>b</sup>	1,917 <sup>b</sup>	1,865 <sup>b</sup>	1,741 <sup>b</sup>	1,818 <sup>b</sup>	1,681 <sup>a</sup>	1,608	1,616	1,522
OxyContin <sup>®</sup>	--	--	194	187	199	232	223	223	234
Tranquilizers	568 <sup>a</sup>	586 <sup>b</sup>	532	497	483	506	482	460	466
Stimulants <sup>3</sup>	665 <sup>b</sup>	578 <sup>b</sup>	507 <sup>b</sup>	506 <sup>b</sup>	547 <sup>b</sup>	416 <sup>a</sup>	374	333	324
Methamphetamine <sup>3</sup>	245 <sup>b</sup>	185 <sup>b</sup>	177 <sup>b</sup>	178 <sup>b</sup>	185 <sup>b</sup>	118	94	85	98
Sedatives	138	131	133	118	110	111	113	126	111
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	3,338 <sup>b</sup>	3,351 <sup>b</sup>	3,269 <sup>b</sup>	3,129 <sup>b</sup>	3,146 <sup>b</sup>	2,914	2,850	2,835	2,753

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.5B Types of Illicit Drug Use in the Past Year among Persons Aged 12 to 17: Percentages, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	22.2 <sup>b</sup>	21.8 <sup>b</sup>	21.0 <sup>b</sup>	19.9	19.6	18.7	19.0	19.5	19.4
Marijuana and Hashish	15.8 <sup>b</sup>	15.0 <sup>a</sup>	14.5	13.3	13.2	12.5 <sup>b</sup>	13.0 <sup>a</sup>	13.6	14.0
Cocaine	2.1 <sup>b</sup>	1.8 <sup>b</sup>	1.6 <sup>b</sup>	1.7 <sup>b</sup>	1.6 <sup>b</sup>	1.5 <sup>b</sup>	1.2	1.0	1.0
Crack	0.4 <sup>b</sup>	0.4 <sup>b</sup>	0.3 <sup>b</sup>	0.2 <sup>b</sup>	0.3 <sup>b</sup>	0.3 <sup>b</sup>	0.1	0.1	0.1
Heroin	0.2	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.1
Hallucinogens	3.8 <sup>b</sup>	3.1	3.0	2.6 <sup>a</sup>	2.6 <sup>a</sup>	2.5 <sup>b</sup>	2.8	3.0	3.1
LSD	1.3 <sup>b</sup>	0.6	0.6	0.6	0.4	0.5	0.7	0.6	0.6
PCP	0.4 <sup>b</sup>	0.4 <sup>b</sup>	0.3 <sup>a</sup>	0.3 <sup>b</sup>	0.2	0.2	0.2 <sup>a</sup>	0.2	0.1
Ecstasy	2.2	1.3 <sup>b</sup>	1.2 <sup>b</sup>	1.0 <sup>b</sup>	1.2 <sup>b</sup>	1.3 <sup>b</sup>	1.4 <sup>b</sup>	1.7	1.9
Inhalants	4.4 <sup>b</sup>	4.5 <sup>b</sup>	4.6 <sup>b</sup>	4.5 <sup>b</sup>	4.4 <sup>b</sup>	3.9	3.9	3.9	3.6
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	9.2 <sup>b</sup>	9.2 <sup>b</sup>	8.8 <sup>b</sup>	8.3 <sup>a</sup>	8.6 <sup>b</sup>	8.1	7.7	7.7	7.4
Pain Relievers	7.6 <sup>b</sup>	7.7 <sup>b</sup>	7.4 <sup>b</sup>	6.9	7.2 <sup>b</sup>	6.7	6.5	6.6	6.2
OxyContin <sup>®</sup>	--	--	0.8	0.7	0.8	0.9	0.9	0.9	1.0
Tranquilizers	2.3 <sup>a</sup>	2.3 <sup>a</sup>	2.1	2.0	1.9	2.0	1.9	1.9	1.9
Stimulants <sup>3</sup>	2.7 <sup>b</sup>	2.3 <sup>b</sup>	2.0 <sup>b</sup>	2.0 <sup>b</sup>	2.2 <sup>b</sup>	1.6 <sup>a</sup>	1.5	1.4	1.3
Methamphetamine <sup>3</sup>	1.0 <sup>b</sup>	0.7 <sup>b</sup>	0.7 <sup>b</sup>	0.7 <sup>b</sup>	0.7 <sup>b</sup>	0.5	0.4	0.3	0.4
Sedatives	0.6	0.5	0.5	0.5	0.4	0.4	0.5	0.5	0.5
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	13.5 <sup>b</sup>	13.4 <sup>b</sup>	13.0 <sup>b</sup>	12.3 <sup>a</sup>	12.4 <sup>b</sup>	11.5	11.4	11.5	11.3

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.6A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLICIT DRUGS<sup>1</sup></b>	2,878 <sup>b</sup>	2,811 <sup>b</sup>	2,674 <sup>a</sup>	2,511	2,481	2,409	2,304	2,471	2,466
Marijuana and Hashish	2,023 <sup>b</sup>	1,971 <sup>a</sup>	1,909	1,730	1,689	1,686	1,657	1,799	1,796
Cocaine	146 <sup>b</sup>	153 <sup>b</sup>	126 <sup>b</sup>	142 <sup>b</sup>	110 <sup>b</sup>	104 <sup>a</sup>	97 <sup>a</sup>	69	60
Crack	26 <sup>a</sup>	32 <sup>b</sup>	16 <sup>a</sup>	25 <sup>b</sup>	12 <sup>a</sup>	16 <sup>a</sup>	7	8	4
Heroin	11	18	15	15	16	3	14	13	8
Hallucinogens	255	260	211	205	181	175	242	217	215
LSD	61	51	38	32	31	30	50	35	46
PCP	23 <sup>a</sup>	20 <sup>a</sup>	11	17	12	9	14	12	6
Ecstasy	124	91	77 <sup>a</sup>	74 <sup>b</sup>	82 <sup>a</sup>	65 <sup>b</sup>	105	115	124
Inhalants	301	320	312	313	340 <sup>a</sup>	295	268	245	263
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	986 <sup>b</sup>	997 <sup>b</sup>	914 <sup>b</sup>	838 <sup>a</sup>	844 <sup>a</sup>	829	710	772	725
Pain Relievers	794 <sup>b</sup>	801 <sup>b</sup>	751 <sup>b</sup>	686	690	669	567	653	611
OxyContin <sup>®</sup>	--	--	66	34	28	47	49	69	47
Tranquilizers	188 <sup>a</sup>	226 <sup>b</sup>	161	161	126	182 <sup>a</sup>	150	143	134
Stimulants <sup>3</sup>	208 <sup>b</sup>	222 <sup>b</sup>	180 <sup>b</sup>	172 <sup>b</sup>	171 <sup>b</sup>	127	113	129	107
Methamphetamine <sup>3</sup>	71 <sup>b</sup>	70 <sup>b</sup>	58 <sup>a</sup>	72 <sup>b</sup>	53 <sup>a</sup>	34	29	36	25
Sedatives	47	47	26	27	45	34	35	40	35
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	1,421 <sup>b</sup>	1,417 <sup>b</sup>	1,328 <sup>b</sup>	1,253 <sup>a</sup>	1,251 <sup>a</sup>	1,181	1,091	1,119	1,095

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.6B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17: Percentages, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	11.6 <sup>b</sup>	11.2 <sup>b</sup>	10.6	9.9	9.8	9.5	9.3 <sup>a</sup>	10.0	10.1
Marijuana and Hashish	8.2 <sup>a</sup>	7.9	7.6	6.8	6.7 <sup>a</sup>	6.7 <sup>a</sup>	6.7 <sup>a</sup>	7.3	7.4
Cocaine	0.6 <sup>b</sup>	0.6 <sup>b</sup>	0.5 <sup>b</sup>	0.6 <sup>b</sup>	0.4 <sup>b</sup>	0.4 <sup>a</sup>	0.4 <sup>a</sup>	0.3	0.2
Crack	0.1 <sup>a</sup>	0.1 <sup>b</sup>	0.1 <sup>a</sup>	0.1 <sup>b</sup>	0.0 <sup>a</sup>	0.1 <sup>a</sup>	0.0	0.0	0.0
Heroin	0.0	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.0
Hallucinogens	1.0	1.0	0.8	0.8	0.7	0.7	1.0	0.9	0.9
LSD	0.2	0.2	0.2	0.1	0.1	0.1	0.2	0.1	0.2
PCP	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.0	0.1	0.0	0.0	0.1	0.1	0.0
Ecstasy	0.5	0.4	0.3 <sup>a</sup>	0.3 <sup>b</sup>	0.3 <sup>a</sup>	0.3 <sup>b</sup>	0.4	0.5	0.5
Inhalants	1.2	1.3	1.2	1.2	1.3	1.2	1.1	1.0	1.1
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	4.0 <sup>b</sup>	4.0 <sup>b</sup>	3.6 <sup>b</sup>	3.3	3.3	3.3	2.9	3.1	3.0
Pain Relievers	3.2 <sup>b</sup>	3.2 <sup>b</sup>	3.0 <sup>a</sup>	2.7	2.7	2.7	2.3	2.7	2.5
OxyContin <sup>®</sup>	--	--	0.3	0.1	0.1	0.2	0.2	0.3	0.2
Tranquilizers	0.8 <sup>a</sup>	0.9 <sup>b</sup>	0.6	0.6	0.5	0.7	0.6	0.6	0.5
Stimulants <sup>3</sup>	0.8 <sup>b</sup>	0.9 <sup>b</sup>	0.7 <sup>b</sup>	0.7 <sup>a</sup>	0.7 <sup>a</sup>	0.5	0.5	0.5	0.4
Methamphetamine <sup>3</sup>	0.3 <sup>b</sup>	0.3 <sup>b</sup>	0.2 <sup>a</sup>	0.3 <sup>b</sup>	0.2 <sup>a</sup>	0.1	0.1	0.1	0.1
Sedatives	0.2	0.2	0.1	0.1	0.2	0.1	0.1	0.2	0.1
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	5.7 <sup>b</sup>	5.7 <sup>b</sup>	5.3 <sup>b</sup>	4.9	4.9	4.7	4.4	4.5	4.5

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.7A Types of Illicit Drug Use in Lifetime among Persons Aged 18 to 25: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	18,562 <sup>b</sup>	19,195	19,043	19,222	19,311	18,795 <sup>b</sup>	18,644 <sup>b</sup>	19,510	19,419
Marijuana and Hashish	16,680 <sup>b</sup>	17,103	16,983	17,032	17,157	16,620 <sup>b</sup>	16,593 <sup>b</sup>	17,544	17,415
Cocaine	4,786	4,755	4,893 <sup>a</sup>	4,905 <sup>a</sup>	5,130 <sup>b</sup>	4,917 <sup>a</sup>	4,730	4,964 <sup>b</sup>	4,529
Crack	1,182 <sup>b</sup>	1,193 <sup>b</sup>	1,135 <sup>b</sup>	1,145 <sup>b</sup>	1,176 <sup>b</sup>	980	904	961	876
Heroin	487 <sup>a</sup>	516	529	496 <sup>a</sup>	511	493 <sup>a</sup>	457 <sup>b</sup>	573	607
Hallucinogens	7,517 <sup>b</sup>	7,377 <sup>b</sup>	6,855 <sup>b</sup>	6,820 <sup>b</sup>	6,614	6,287	5,836 <sup>a</sup>	6,365	6,261
LSD	4,928 <sup>b</sup>	4,450 <sup>b</sup>	3,900 <sup>b</sup>	3,417 <sup>b</sup>	2,925 <sup>b</sup>	2,374	2,145	2,294	2,176
PCP	834 <sup>b</sup>	948 <sup>b</sup>	737 <sup>b</sup>	716 <sup>b</sup>	637 <sup>b</sup>	539 <sup>a</sup>	513	468	437
Ecstasy	4,674 <sup>b</sup>	4,683 <sup>b</sup>	4,439	4,464	4,382	4,191	3,992	4,168	4,216
Inhalants	4,867 <sup>b</sup>	4,729 <sup>b</sup>	4,516 <sup>b</sup>	4,333 <sup>b</sup>	4,089 <sup>b</sup>	3,690 <sup>a</sup>	3,411	3,581	3,363
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	8,778 <sup>b</sup>	9,360 <sup>a</sup>	9,564	9,990	10,051	9,790	9,626	9,819	9,780
Pain Relievers	6,854 <sup>b</sup>	7,513 <sup>b</sup>	7,813	8,279	8,342	8,128	8,042	8,156	8,098
OxyContin <sup>®</sup>	820 <sup>b</sup>	1,145 <sup>b</sup>	1,376 <sup>b</sup>	1,616 <sup>b</sup>	1,676 <sup>b</sup>	1,706 <sup>b</sup>	1,736 <sup>b</sup>	2,084	2,129
Tranquilizers	3,473 <sup>b</sup>	3,898 <sup>b</sup>	3,919 <sup>b</sup>	4,356	4,274	4,147 <sup>a</sup>	4,033 <sup>b</sup>	4,291	4,472
Stimulants <sup>3</sup>	3,796 <sup>a</sup>	3,873 <sup>b</sup>	3,892 <sup>b</sup>	4,008 <sup>b</sup>	3,922 <sup>b</sup>	3,574	3,460	3,555	3,510
Methamphetamine <sup>3</sup>	2,301 <sup>b</sup>	2,192 <sup>b</sup>	2,247 <sup>b</sup>	2,190 <sup>b</sup>	2,102 <sup>b</sup>	1,844 <sup>b</sup>	1,544 <sup>a</sup>	1,511	1,353
Sedatives	648 <sup>b</sup>	565 <sup>a</sup>	627 <sup>b</sup>	650 <sup>b</sup>	609 <sup>b</sup>	583 <sup>a</sup>	506	523	464
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	12,436	12,770	12,630	12,800 <sup>a</sup>	12,940 <sup>a</sup>	12,555	12,064	12,499	12,346

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.7B Types of Illicit Drug Use in Lifetime among Persons Aged 18 to 25: Percentages, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	59.8 <sup>b</sup>	60.5 <sup>b</sup>	59.2 <sup>b</sup>	59.2 <sup>b</sup>	59.0 <sup>b</sup>	57.4	56.6	58.1	57.0
Marijuana and Hashish	53.8 <sup>b</sup>	53.9 <sup>b</sup>	52.8 <sup>a</sup>	52.4	52.4	50.8	50.4	52.2	51.1
Cocaine	15.4 <sup>b</sup>	15.0 <sup>b</sup>	15.2 <sup>b</sup>	15.1 <sup>b</sup>	15.7 <sup>b</sup>	15.0 <sup>b</sup>	14.4 <sup>a</sup>	14.8 <sup>b</sup>	13.3
Crack	3.8 <sup>b</sup>	3.8 <sup>b</sup>	3.5 <sup>b</sup>	3.5 <sup>b</sup>	3.6 <sup>b</sup>	3.0 <sup>a</sup>	2.7	2.9	2.6
Heroin	1.6	1.6	1.6	1.5	1.6	1.5	1.4 <sup>a</sup>	1.7	1.8
Hallucinogens	24.2 <sup>b</sup>	23.3 <sup>b</sup>	21.3 <sup>b</sup>	21.0 <sup>b</sup>	20.2 <sup>b</sup>	19.2	17.7	19.0	18.4
LSD	15.9 <sup>b</sup>	14.0 <sup>b</sup>	12.1 <sup>b</sup>	10.5 <sup>b</sup>	8.9 <sup>b</sup>	7.3 <sup>b</sup>	6.5	6.8	6.4
PCP	2.7 <sup>b</sup>	3.0 <sup>b</sup>	2.3 <sup>b</sup>	2.2 <sup>b</sup>	1.9 <sup>b</sup>	1.6 <sup>a</sup>	1.6	1.4	1.3
Ecstasy	15.1 <sup>b</sup>	14.8 <sup>b</sup>	13.8 <sup>b</sup>	13.7 <sup>b</sup>	13.4 <sup>a</sup>	12.8	12.1	12.4	12.4
Inhalants	15.7 <sup>b</sup>	14.9 <sup>b</sup>	14.0 <sup>b</sup>	13.3 <sup>b</sup>	12.5 <sup>b</sup>	11.3 <sup>b</sup>	10.4	10.7 <sup>a</sup>	9.9
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	28.3	29.5	29.7	30.8 <sup>b</sup>	30.7 <sup>b</sup>	29.9 <sup>a</sup>	29.2	29.2	28.7
Pain Relievers	22.1 <sup>b</sup>	23.7	24.3	25.5 <sup>b</sup>	25.5 <sup>b</sup>	24.8	24.4	24.3	23.8
OxyContin <sup>®</sup>	2.6 <sup>b</sup>	3.6 <sup>b</sup>	4.3 <sup>b</sup>	5.0 <sup>b</sup>	5.1 <sup>b</sup>	5.2 <sup>b</sup>	5.3 <sup>b</sup>	6.2	6.2
Tranquilizers	11.2 <sup>b</sup>	12.3	12.2 <sup>a</sup>	13.4	13.1	12.7	12.2 <sup>a</sup>	12.8	13.1
Stimulants <sup>3</sup>	12.2 <sup>b</sup>	12.2 <sup>b</sup>	12.1 <sup>b</sup>	12.3 <sup>b</sup>	12.0 <sup>b</sup>	10.9	10.5	10.6	10.3
Methamphetamine <sup>3</sup>	7.4 <sup>b</sup>	6.9 <sup>b</sup>	7.0 <sup>b</sup>	6.7 <sup>b</sup>	6.4 <sup>b</sup>	5.6 <sup>b</sup>	4.7 <sup>b</sup>	4.5 <sup>a</sup>	4.0
Sedatives	2.1 <sup>b</sup>	1.8 <sup>b</sup>	1.9 <sup>b</sup>	2.0 <sup>b</sup>	1.9 <sup>b</sup>	1.8 <sup>b</sup>	1.5	1.6	1.4
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	40.1 <sup>b</sup>	40.2 <sup>b</sup>	39.2 <sup>b</sup>	39.4 <sup>b</sup>	39.5 <sup>b</sup>	38.4 <sup>b</sup>	36.6	37.2	36.2

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.8A Types of Illicit Drug Use in the Past Year among Persons Aged 18 to 25: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	11,000 <sup>b</sup>	10,976 <sup>b</sup>	10,913 <sup>b</sup>	11,117 <sup>b</sup>	11,252 <sup>b</sup>	10,864 <sup>b</sup>	11,021 <sup>b</sup>	12,094	11,935
Marijuana and Hashish	9,238 <sup>b</sup>	9,044 <sup>b</sup>	8,963 <sup>b</sup>	9,111 <sup>b</sup>	9,169 <sup>b</sup>	8,985 <sup>b</sup>	9,087 <sup>b</sup>	10,259	10,161
Cocaine	2,087 <sup>b</sup>	2,098 <sup>b</sup>	2,134 <sup>b</sup>	2,247 <sup>b</sup>	2,259 <sup>b</sup>	2,084 <sup>b</sup>	1,820 <sup>a</sup>	1,773 <sup>a</sup>	1,576
Crack	266 <sup>b</sup>	275 <sup>b</sup>	263 <sup>b</sup>	328 <sup>b</sup>	293 <sup>b</sup>	248 <sup>a</sup>	214	169	177
Heroin	122 <sup>b</sup>	96 <sup>b</sup>	118 <sup>b</sup>	159	147 <sup>a</sup>	142 <sup>a</sup>	145 <sup>a</sup>	175	210
Hallucinogens	2,610	2,112 <sup>b</sup>	1,944 <sup>b</sup>	2,051 <sup>b</sup>	2,147 <sup>a</sup>	2,048 <sup>b</sup>	2,108 <sup>b</sup>	2,409	2,422
LSD	553	336 <sup>b</sup>	317 <sup>b</sup>	322 <sup>b</sup>	379 <sup>b</sup>	375 <sup>b</sup>	503	515	544
PCP	95 <sup>b</sup>	117 <sup>b</sup>	100 <sup>b</sup>	57	76 <sup>a</sup>	58	30	34	35
Ecstasy	1,784 <sup>b</sup>	1,164 <sup>b</sup>	999 <sup>b</sup>	998 <sup>b</sup>	1,233 <sup>b</sup>	1,154 <sup>b</sup>	1,287 <sup>a</sup>	1,448	1,498
Inhalants	685	662	674	685	577	526	533	626	589
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	4,452 <sup>b</sup>	4,637	4,807	4,915	5,121	4,913	4,868	5,027	4,862
Pain Relievers	3,540	3,803	3,834	4,030	4,070 <sup>a</sup>	3,970	3,941	4,000	3,769
OxyContin <sup>®</sup>	--	--	562 <sup>b</sup>	559 <sup>b</sup>	574 <sup>b</sup>	566 <sup>b</sup>	598 <sup>b</sup>	759	746
Tranquilizers	1,509 <sup>b</sup>	1,682	1,658	1,796	1,785	1,699	1,756	1,787	1,790
Stimulants <sup>3</sup>	1,233	1,197	1,280	1,242	1,375	1,134	1,113	1,223	1,233
Methamphetamine <sup>3</sup>	619 <sup>b</sup>	594 <sup>b</sup>	619 <sup>b</sup>	576 <sup>b</sup>	552 <sup>b</sup>	404 <sup>b</sup>	269	301	266
Sedatives	162	156	163	191	185	172	175	159	162
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	6,277	6,266	6,201	6,467	6,627	6,370	6,227	6,652	6,368

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.8B Types of Illicit Drug Use in the Past Year among Persons Aged 18 to 25: Percentages, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	35.5	34.6	33.9	34.2	34.4	33.2 <sup>b</sup>	33.5 <sup>a</sup>	36.0	35.0
Marijuana and Hashish	29.8	28.5 <sup>a</sup>	27.8 <sup>b</sup>	28.0 <sup>b</sup>	28.0 <sup>b</sup>	27.5 <sup>b</sup>	27.6 <sup>b</sup>	30.6	29.8
Cocaine	6.7 <sup>b</sup>	6.6 <sup>b</sup>	6.6 <sup>b</sup>	6.9 <sup>b</sup>	6.9 <sup>b</sup>	6.4 <sup>b</sup>	5.5 <sup>b</sup>	5.3 <sup>a</sup>	4.6
Crack	0.9 <sup>b</sup>	0.9 <sup>b</sup>	0.8 <sup>b</sup>	1.0 <sup>b</sup>	0.9 <sup>b</sup>	0.8 <sup>a</sup>	0.6	0.5	0.5
Heroin	0.4 <sup>a</sup>	0.3 <sup>b</sup>	0.4 <sup>b</sup>	0.5	0.4	0.4	0.4	0.5	0.6
Hallucinogens	8.4 <sup>b</sup>	6.7	6.0 <sup>b</sup>	6.3 <sup>a</sup>	6.6	6.3 <sup>a</sup>	6.4 <sup>a</sup>	7.2	7.1
LSD	1.8	1.1 <sup>b</sup>	1.0 <sup>b</sup>	1.0 <sup>b</sup>	1.2 <sup>b</sup>	1.1 <sup>b</sup>	1.5	1.5	1.6
PCP	0.3 <sup>b</sup>	0.4 <sup>b</sup>	0.3 <sup>b</sup>	0.2	0.2 <sup>a</sup>	0.2	0.1	0.1	0.1
Ecstasy	5.8 <sup>b</sup>	3.7 <sup>b</sup>	3.1 <sup>b</sup>	3.1 <sup>b</sup>	3.8 <sup>a</sup>	3.5 <sup>b</sup>	3.9	4.3	4.4
Inhalants	2.2 <sup>b</sup>	2.1 <sup>a</sup>	2.1 <sup>a</sup>	2.1 <sup>a</sup>	1.8	1.6	1.6	1.9	1.7
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	14.4	14.6	14.9	15.1	15.6 <sup>b</sup>	15.0	14.8	15.0	14.3
Pain Relievers	11.4	12.0 <sup>a</sup>	11.9 <sup>a</sup>	12.4 <sup>b</sup>	12.4 <sup>b</sup>	12.1 <sup>b</sup>	12.0 <sup>a</sup>	11.9 <sup>a</sup>	11.1
OxyContin <sup>®</sup>	--	--	1.7 <sup>b</sup>	1.7 <sup>b</sup>	1.8 <sup>a</sup>	1.7 <sup>b</sup>	1.8 <sup>a</sup>	2.3	2.2
Tranquilizers	4.9	5.3	5.2	5.5	5.5	5.2	5.3	5.3	5.3
Stimulants <sup>3</sup>	4.0	3.8	4.0	3.8	4.2 <sup>a</sup>	3.5	3.4	3.6	3.6
Methamphetamine <sup>3</sup>	2.0 <sup>b</sup>	1.9 <sup>b</sup>	1.9 <sup>b</sup>	1.8 <sup>b</sup>	1.7 <sup>b</sup>	1.2 <sup>b</sup>	0.8	0.9	0.8
Sedatives	0.5	0.5	0.5	0.6	0.6	0.5	0.5	0.5	0.5
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	20.2 <sup>b</sup>	19.7	19.3	19.9 <sup>a</sup>	20.2 <sup>b</sup>	19.5	18.9	19.8 <sup>a</sup>	18.7

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.9A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	6,264 <sup>b</sup>	6,435 <sup>b</sup>	6,239 <sup>b</sup>	6,543 <sup>b</sup>	6,482 <sup>b</sup>	6,464 <sup>b</sup>	6,465 <sup>b</sup>	7,124	7,328
Marijuana and Hashish	5,376 <sup>b</sup>	5,400 <sup>b</sup>	5,196 <sup>b</sup>	5,388 <sup>b</sup>	5,322 <sup>b</sup>	5,378 <sup>b</sup>	5,426 <sup>b</sup>	6,082	6,292
Cocaine	631 <sup>a</sup>	696 <sup>b</sup>	692 <sup>b</sup>	832 <sup>b</sup>	729 <sup>b</sup>	573	507	462	496
Crack	52	55	80	104	72	56	53	39	72
Heroin	39 <sup>a</sup>	25 <sup>b</sup>	43 <sup>a</sup>	59	54	48 <sup>a</sup>	74	65	89
Hallucinogens	589	545 <sup>a</sup>	477 <sup>b</sup>	503 <sup>b</sup>	563	475 <sup>b</sup>	549 <sup>a</sup>	617	673
LSD	36 <sup>b</sup>	67	83	60 <sup>a</sup>	58 <sup>a</sup>	70	97	96	105
PCP	15	30 <sup>a</sup>	17	11	14	15	4	5	5
Ecstasy	355	230 <sup>b</sup>	231 <sup>b</sup>	267 <sup>b</sup>	326	229 <sup>b</sup>	304 <sup>a</sup>	377	406
Inhalants	150	134	144	146	129	130	104	148	135
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	1,698 <sup>b</sup>	1,922	1,978	2,046	2,115	1,949	1,943	2,119	2,003
Pain Relievers	1,262 <sup>b</sup>	1,486	1,504	1,535	1,620	1,517	1,502	1,600	1,519
OxyContin <sup>®</sup>	--	--	142 <sup>b</sup>	146 <sup>b</sup>	139 <sup>b</sup>	155 <sup>b</sup>	129 <sup>b</sup>	175 <sup>a</sup>	237
Tranquilizers	487	525	566	606	647	571	569	616	566
Stimulants <sup>3</sup>	409	412	479	468	456	352	348	421	410
Methamphetamine <sup>3</sup>	183 <sup>b</sup>	197 <sup>b</sup>	220 <sup>b</sup>	225 <sup>b</sup>	183 <sup>b</sup>	116	74	83	83
Sedatives	49	58	49	74	54	72	64	84	57
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	2,459 <sup>a</sup>	2,665	2,619	2,850	2,904	2,643	2,586	2,794	2,710

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.9B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25: Percentages, 2002-2010**

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	20.2 <sup>a</sup>	20.3 <sup>a</sup>	19.4 <sup>b</sup>	20.1 <sup>a</sup>	19.8 <sup>b</sup>	19.7 <sup>b</sup>	19.6 <sup>b</sup>	21.2	21.5
Marijuana and Hashish	17.3 <sup>a</sup>	17.0 <sup>b</sup>	16.1 <sup>b</sup>	16.6 <sup>b</sup>	16.3 <sup>b</sup>	16.4 <sup>b</sup>	16.5 <sup>b</sup>	18.1	18.5
Cocaine	2.0 <sup>b</sup>	2.2 <sup>b</sup>	2.1 <sup>b</sup>	2.6 <sup>b</sup>	2.2 <sup>b</sup>	1.7	1.5	1.4	1.5
Crack	0.2	0.2	0.3	0.3	0.2	0.2	0.2	0.1	0.2
Heroin	0.1 <sup>a</sup>	0.1 <sup>b</sup>	0.1 <sup>a</sup>	0.2	0.2	0.1	0.2	0.2	0.3
Hallucinogens	1.9	1.7	1.5 <sup>b</sup>	1.5 <sup>a</sup>	1.7	1.5 <sup>b</sup>	1.7	1.8	2.0
LSD	0.1 <sup>b</sup>	0.2	0.3	0.2	0.2 <sup>a</sup>	0.2	0.3	0.3	0.3
PCP	0.0	0.1 <sup>a</sup>	0.1	0.0	0.0	0.0	0.0	0.0	0.0
Ecstasy	1.1	0.7 <sup>b</sup>	0.7 <sup>b</sup>	0.8 <sup>b</sup>	1.0	0.7 <sup>b</sup>	0.9	1.1	1.2
Inhalants	0.5	0.4	0.4	0.5	0.4	0.4	0.3	0.4	0.4
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	5.5	6.1	6.1	6.3	6.5	6.0	5.9	6.3	5.9
Pain Relievers	4.1	4.7	4.7	4.7	4.9	4.6	4.6	4.8	4.5
OxyContin <sup>®</sup>	--	--	0.4 <sup>b</sup>	0.4 <sup>b</sup>	0.4 <sup>b</sup>	0.5 <sup>a</sup>	0.4 <sup>b</sup>	0.5	0.7
Tranquilizers	1.6	1.7	1.8	1.9	2.0	1.7	1.7	1.8	1.7
Stimulants <sup>3</sup>	1.3	1.3	1.5	1.4	1.4	1.1	1.1	1.3	1.2
Methamphetamine <sup>3</sup>	0.6 <sup>b</sup>	0.6 <sup>b</sup>	0.7 <sup>b</sup>	0.7 <sup>b</sup>	0.6 <sup>b</sup>	0.4	0.2	0.2	0.2
Sedatives	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	7.9	8.4	8.1	8.8 <sup>a</sup>	8.9 <sup>a</sup>	8.1	7.8	8.3	8.0

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.10A Types of Illicit Drug Use in Lifetime among Persons Aged 26 or Older: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLICIT DRUGS<sup>1</sup></b>	82,051 <sup>b</sup>	83,380 <sup>b</sup>	83,448 <sup>b</sup>	85,841 <sup>b</sup>	85,454 <sup>b</sup>	88,858 <sup>b</sup>	92,161	92,607	93,838
Marijuana and Hashish	73,162 <sup>b</sup>	74,620 <sup>b</sup>	75,001 <sup>b</sup>	76,111 <sup>b</sup>	76,275 <sup>b</sup>	79,815 <sup>b</sup>	81,703 <sup>a</sup>	82,721	84,678
Cocaine	28,453 <sup>b</sup>	29,488 <sup>b</sup>	28,649 <sup>b</sup>	28,185 <sup>b</sup>	29,598 <sup>b</sup>	30,423 <sup>a</sup>	31,572	31,232	32,320
Crack	7,057 <sup>a</sup>	6,599 <sup>b</sup>	6,580 <sup>b</sup>	6,675 <sup>b</sup>	7,250 <sup>a</sup>	7,495	7,462	7,340	8,223
Heroin	3,087	3,148	2,538 <sup>b</sup>	2,979	3,214	3,233	3,263	3,054	3,473
Hallucinogens	25,378 <sup>b</sup>	25,736 <sup>b</sup>	26,318 <sup>b</sup>	25,910 <sup>b</sup>	27,682 <sup>b</sup>	27,032 <sup>b</sup>	29,155	29,879	30,216
LSD	18,917 <sup>b</sup>	19,577	19,185 <sup>a</sup>	18,728 <sup>b</sup>	20,202	20,083	21,133	21,092	20,913
PCP	6,352	5,955	5,851	5,705	5,853	5,497	6,016	5,672	5,744
Ecstasy	4,656 <sup>b</sup>	5,619 <sup>b</sup>	6,168 <sup>b</sup>	6,632 <sup>b</sup>	7,406 <sup>b</sup>	7,788 <sup>b</sup>	8,400 <sup>b</sup>	9,494 <sup>b</sup>	11,066
Inhalants	15,398	15,596	15,521	15,754	16,229	16,365	16,555	16,599	16,292
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	35,779 <sup>b</sup>	36,269 <sup>b</sup>	36,190 <sup>b</sup>	36,562 <sup>b</sup>	37,743	37,676	39,572	39,156	39,277
Pain Relievers	19,983 <sup>b</sup>	20,896 <sup>b</sup>	21,076 <sup>b</sup>	21,896 <sup>b</sup>	22,484 <sup>a</sup>	22,481 <sup>a</sup>	24,473	24,513	24,439
OxyContin <sup>®</sup>	881 <sup>b</sup>	1,433 <sup>b</sup>	1,392 <sup>b</sup>	1,589 <sup>b</sup>	2,080 <sup>b</sup>	2,289 <sup>b</sup>	2,766 <sup>b</sup>	3,368	3,626
Tranquilizers	14,959 <sup>b</sup>	15,447 <sup>a</sup>	15,117 <sup>b</sup>	15,919	16,289	15,313 <sup>a</sup>	16,725	16,771	16,909
Stimulants <sup>3</sup>	18,588	18,085	17,499	16,105 <sup>a</sup>	17,616	17,348	17,076	17,793	17,621
Methamphetamine <sup>3</sup>	12,648	12,565	11,920	10,154 <sup>a</sup>	11,762	10,999	10,859	11,141	11,503
Sedatives	9,075 <sup>b</sup>	8,696 <sup>b</sup>	9,012 <sup>b</sup>	8,101 <sup>a</sup>	7,989 <sup>a</sup>	7,599	8,173 <sup>a</sup>	7,858	6,990
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	52,578 <sup>b</sup>	53,040 <sup>b</sup>	52,672 <sup>b</sup>	54,078 <sup>b</sup>	55,018 <sup>b</sup>	56,302 <sup>b</sup>	59,022	58,808	59,707

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.10B Types of Illicit Drug Use in Lifetime among Persons Aged 26 or Older: Percentages, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	45.7 <sup>b</sup>	46.1 <sup>b</sup>	45.6 <sup>b</sup>	46.3 <sup>a</sup>	45.5 <sup>b</sup>	46.8	48.0	47.8	48.1
Marijuana and Hashish	40.8 <sup>b</sup>	41.2 <sup>b</sup>	41.0 <sup>b</sup>	41.1 <sup>b</sup>	40.6 <sup>b</sup>	42.0	42.6	42.7	43.4
Cocaine	15.9	16.3	15.6	15.2 <sup>b</sup>	15.8	16.0	16.4	16.1	16.6
Crack	3.9	3.6 <sup>a</sup>	3.6 <sup>a</sup>	3.6 <sup>a</sup>	3.9	3.9	3.9	3.8	4.2
Heroin	1.7	1.7	1.4 <sup>a</sup>	1.6	1.7	1.7	1.7	1.6	1.8
Hallucinogens	14.1 <sup>b</sup>	14.2 <sup>b</sup>	14.4 <sup>a</sup>	14.0 <sup>b</sup>	14.7	14.2 <sup>b</sup>	15.2	15.4	15.5
LSD	10.5	10.8	10.5	10.1	10.8	10.6	11.0	10.9	10.7
PCP	3.5 <sup>b</sup>	3.3	3.2	3.1	3.1	2.9	3.1	2.9	2.9
Ecstasy	2.6 <sup>b</sup>	3.1 <sup>b</sup>	3.4 <sup>b</sup>	3.6 <sup>b</sup>	3.9 <sup>b</sup>	4.1 <sup>b</sup>	4.4 <sup>b</sup>	4.9 <sup>b</sup>	5.7
Inhalants	8.6	8.6	8.5	8.5	8.6	8.6	8.6	8.6	8.3
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	19.9	20.0	19.8	19.7	20.1	19.8	20.6	20.2	20.1
Pain Relievers	11.1 <sup>b</sup>	11.5 <sup>a</sup>	11.5 <sup>a</sup>	11.8	12.0	11.8	12.7	12.7	12.5
OxyContin <sup>®</sup>	0.5 <sup>b</sup>	0.8 <sup>b</sup>	0.8 <sup>b</sup>	0.9 <sup>b</sup>	1.1 <sup>b</sup>	1.2 <sup>b</sup>	1.4 <sup>b</sup>	1.7	1.9
Tranquilizers	8.3	8.5	8.3	8.6	8.7	8.1	8.7	8.7	8.7
Stimulants <sup>3</sup>	10.4 <sup>b</sup>	10.0 <sup>b</sup>	9.6	8.7	9.4	9.1	8.9	9.2	9.0
Methamphetamine <sup>3</sup>	7.1 <sup>b</sup>	6.9 <sup>b</sup>	6.5 <sup>a</sup>	5.5	6.3	5.8	5.7	5.8	5.9
Sedatives	5.1 <sup>b</sup>	4.8 <sup>b</sup>	4.9 <sup>b</sup>	4.4 <sup>b</sup>	4.3 <sup>a</sup>	4.0	4.3 <sup>a</sup>	4.1	3.6
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	29.3 <sup>a</sup>	29.3 <sup>a</sup>	28.8 <sup>b</sup>	29.2 <sup>a</sup>	29.3 <sup>a</sup>	29.7	30.7	30.4	30.6

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.11A Types of Illicit Drug Use in the Past Year among Persons Aged 26 or Older: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	18,639 <sup>b</sup>	18,562 <sup>b</sup>	18,594 <sup>b</sup>	18,883 <sup>b</sup>	19,552 <sup>b</sup>	20,102 <sup>b</sup>	19,778 <sup>b</sup>	21,051	22,140
Marijuana and Hashish	12,612 <sup>b</sup>	12,441 <sup>b</sup>	12,833 <sup>b</sup>	12,880 <sup>b</sup>	12,860 <sup>b</sup>	12,953 <sup>b</sup>	13,443 <sup>b</sup>	14,923	15,645
Cocaine	3,307 <sup>a</sup>	3,372 <sup>a</sup>	3,114	2,852	3,399 <sup>b</sup>	3,271 <sup>a</sup>	3,130	2,784	2,685
Crack	1,188 <sup>b</sup>	1,036 <sup>a</sup>	978 <sup>a</sup>	1,000 <sup>a</sup>	1,113 <sup>b</sup>	1,137 <sup>b</sup>	862	818	676
Heroin	231	182 <sup>a</sup>	232	184 <sup>a</sup>	376	200 <sup>a</sup>	266	396	380
Hallucinogens	1,194	1,046	1,182	1,093	1,160	1,088	879 <sup>b</sup>	1,366	1,348
LSD	131	73 <sup>a</sup>	117	91	181	120	115	122	195
PCP	31	13	38	22	52	25	*	40	24
Ecstasy	836	626	606	705	591	650	499	935	682
Inhalants	320 <sup>a</sup>	295 <sup>a</sup>	413	358	531	560	536	508	561
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	8,057 <sup>b</sup>	8,231 <sup>a</sup>	7,814 <sup>b</sup>	8,331 <sup>a</sup>	9,180	9,334	8,385 <sup>a</sup>	9,093	9,357
Pain Relievers	5,582 <sup>b</sup>	5,950 <sup>a</sup>	5,556 <sup>b</sup>	6,044 <sup>a</sup>	6,761	6,815	6,336	6,789	6,923
OxyContin <sup>®</sup>	--	--	457 <sup>b</sup>	480 <sup>b</sup>	549 <sup>b</sup>	624 <sup>a</sup>	638	695	890
Tranquilizers	2,772	2,783 <sup>a</sup>	2,878	2,956	2,791 <sup>a</sup>	3,077	2,865	3,213	3,325
Stimulants <sup>3</sup>	1,482	1,256	1,467	1,340	1,868 <sup>a</sup>	1,448	1,152	1,504	1,330
Methamphetamine <sup>3</sup>	891 <sup>a</sup>	823	1,012 <sup>b</sup>	849 <sup>a</sup>	1,151 <sup>b</sup>	822	487	779	590
Sedatives	681	544	441	441	632	580	334 <sup>b</sup>	526	634
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	10,808	10,689	10,189 <sup>a</sup>	10,513	11,481	11,860	10,912	11,514	11,454

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.11B Types of Illicit Drug Use in the Past Year among Persons Aged 26 or Older: Percentages, 2002-2010**

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	10.4 <sup>a</sup>	10.3 <sup>b</sup>	10.2 <sup>b</sup>	10.2 <sup>b</sup>	10.4 <sup>a</sup>	10.6	10.3 <sup>b</sup>	10.9	11.3
Marijuana and Hashish	7.0 <sup>b</sup>	6.9 <sup>b</sup>	7.0 <sup>b</sup>	6.9 <sup>b</sup>	6.8 <sup>b</sup>	6.8 <sup>b</sup>	7.0 <sup>b</sup>	7.7	8.0
Cocaine	1.8 <sup>b</sup>	1.9 <sup>b</sup>	1.7 <sup>a</sup>	1.5	1.8 <sup>b</sup>	1.7 <sup>a</sup>	1.6	1.4	1.4
Crack	0.7 <sup>b</sup>	0.6 <sup>b</sup>	0.5 <sup>a</sup>	0.5 <sup>b</sup>	0.6 <sup>b</sup>	0.6 <sup>b</sup>	0.4	0.4	0.3
Heroin	0.1	0.1 <sup>a</sup>	0.1	0.1 <sup>a</sup>	0.2	0.1 <sup>a</sup>	0.1	0.2	0.2
Hallucinogens	0.7	0.6	0.6	0.6	0.6	0.6	0.5 <sup>b</sup>	0.7	0.7
LSD	0.1	0.0 <sup>a</sup>	0.1	0.0	0.1	0.1	0.1	0.1	0.1
PCP	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Ecstasy	0.5	0.3	0.3	0.4	0.3	0.3	0.3	0.5	0.3
Inhalants	0.2	0.2 <sup>a</sup>	0.2	0.2	0.3	0.3	0.3	0.3	0.3
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	4.5	4.5	4.3 <sup>a</sup>	4.5	4.9	4.9	4.4	4.7	4.8
Pain Relievers	3.1 <sup>a</sup>	3.3	3.0 <sup>a</sup>	3.3	3.6	3.6	3.3	3.5	3.5
OxyContin <sup>®</sup>	--	--	0.2 <sup>b</sup>	0.3 <sup>b</sup>	0.3 <sup>a</sup>	0.3	0.3	0.4	0.5
Tranquilizers	1.5	1.5	1.6	1.6	1.5	1.6	1.5	1.7	1.7
Stimulants <sup>3</sup>	0.8	0.7	0.8	0.7	1.0 <sup>b</sup>	0.8	0.6	0.8	0.7
Methamphetamine <sup>3</sup>	0.5 <sup>b</sup>	0.5 <sup>a</sup>	0.6 <sup>b</sup>	0.5 <sup>a</sup>	0.6 <sup>b</sup>	0.4	0.3	0.4	0.3
Sedatives	0.4	0.3	0.2	0.2	0.3	0.3	0.2 <sup>a</sup>	0.3	0.3
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	6.0	5.9	5.6	5.7	6.1	6.2	5.7	5.9	5.9

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.12A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older: Numbers in Thousands, 2002-2010**

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	10,380 <sup>b</sup>	10,224 <sup>b</sup>	10,159 <sup>b</sup>	10,666 <sup>b</sup>	11,394 <sup>a</sup>	10,984 <sup>b</sup>	11,308 <sup>b</sup>	12,218	12,828
Marijuana and Hashish	7,185 <sup>b</sup>	7,267 <sup>b</sup>	7,471 <sup>b</sup>	7,508 <sup>b</sup>	7,801 <sup>b</sup>	7,384 <sup>b</sup>	8,120 <sup>a</sup>	8,837	9,286
Cocaine	1,243 <sup>a</sup>	1,431 <sup>b</sup>	1,204 <sup>a</sup>	1,423 <sup>b</sup>	1,582 <sup>b</sup>	1,399 <sup>b</sup>	1,251 <sup>a</sup>	1,105	910
Crack	488 <sup>a</sup>	517 <sup>a</sup>	370	553 <sup>a</sup>	618 <sup>b</sup>	537 <sup>a</sup>	300	445	303
Heroin	116	76	108	62	268	102	126	116	141
Hallucinogens	352	238	241	380	262	346	269	424	320
LSD	15	15	20	12	41	45	*	27	*
PCP	20	*	21	20	*	17	*	36	24
Ecstasy	197	149	142	161	120	209	146	267	165
Inhalants	183	117 <sup>a</sup>	181	152	291	191	268	167	288
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	3,602	3,532 <sup>a</sup>	3,218 <sup>b</sup>	3,607 <sup>a</sup>	4,136	4,117	3,571 <sup>a</sup>	4,062	4,239
Pain Relievers	2,321 <sup>a</sup>	2,406 <sup>a</sup>	2,149 <sup>b</sup>	2,437 <sup>a</sup>	2,909	2,987	2,678	3,004	2,971
OxyContin <sup>®</sup>	--	--	118 <sup>a</sup>	153	109 <sup>a</sup>	167	256	266	280
Tranquilizers	1,129	1,078 <sup>a</sup>	889 <sup>b</sup>	1,050 <sup>a</sup>	993 <sup>b</sup>	1,081 <sup>a</sup>	1,081 <sup>a</sup>	1,251	1,460
Stimulants <sup>3</sup>	685	676	653	548	757	573	444	741	559
Methamphetamine <sup>3</sup>	428 <sup>a</sup>	459 <sup>a</sup>	428 <sup>a</sup>	331	495 <sup>b</sup>	380	211	383	245
Sedatives	340	189	190	171	285	240	134	247	282
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	4,897	4,767	4,301 <sup>a</sup>	4,860	5,460	5,446	4,889	5,244	5,211

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.12B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older: Percentages, 2002-2010**

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	5.8 <sup>b</sup>	5.6 <sup>b</sup>	5.5 <sup>b</sup>	5.8 <sup>b</sup>	6.1	5.8 <sup>a</sup>	5.9 <sup>a</sup>	6.3	6.6
Marijuana and Hashish	4.0 <sup>b</sup>	4.0 <sup>b</sup>	4.1 <sup>b</sup>	4.1 <sup>b</sup>	4.2 <sup>a</sup>	3.9 <sup>b</sup>	4.2 <sup>a</sup>	4.6	4.8
Cocaine	0.7 <sup>b</sup>	0.8 <sup>b</sup>	0.7 <sup>a</sup>	0.8 <sup>b</sup>	0.8 <sup>b</sup>	0.7 <sup>b</sup>	0.7 <sup>a</sup>	0.6	0.5
Crack	0.3 <sup>a</sup>	0.3 <sup>a</sup>	0.2	0.3 <sup>a</sup>	0.3 <sup>b</sup>	0.3 <sup>a</sup>	0.2	0.2	0.2
Heroin	0.1	0.0	0.1	0.0	0.1	0.1	0.1	0.1	0.1
Hallucinogens	0.2	0.1	0.1	0.2	0.1	0.2	0.1	0.2	0.2
LSD	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	*
PCP	0.0	*	0.0	0.0	*	0.0	*	0.0	0.0
Ecstasy	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Inhalants	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	2.0	2.0	1.8 <sup>a</sup>	1.9	2.2	2.2	1.9	2.1	2.2
Pain Relievers	1.3	1.3	1.2 <sup>b</sup>	1.3	1.5	1.6	1.4	1.6	1.5
OxyContin <sup>®</sup>	--	--	0.1 <sup>a</sup>	0.1	0.1 <sup>a</sup>	0.1	0.1	0.1	0.1
Tranquilizers	0.6	0.6	0.5 <sup>b</sup>	0.6 <sup>a</sup>	0.5 <sup>a</sup>	0.6	0.6 <sup>a</sup>	0.6	0.7
Stimulants <sup>3</sup>	0.4	0.4	0.4	0.3	0.4	0.3	0.2	0.4	0.3
Methamphetamine <sup>3</sup>	0.2 <sup>a</sup>	0.3 <sup>b</sup>	0.2 <sup>a</sup>	0.2	0.3 <sup>b</sup>	0.2	0.1	0.2	0.1
Sedatives	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	2.7	2.6	2.3	2.6	2.9	2.9	2.5	2.7	2.7

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.13A Types of Illicit Drug Use in Lifetime among Persons Aged 18 or Older: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	100,612 <sup>b</sup>	102,574 <sup>b</sup>	102,491 <sup>b</sup>	105,063 <sup>b</sup>	104,765 <sup>b</sup>	107,654 <sup>b</sup>	110,805	112,117	113,257
Marijuana and Hashish	89,842 <sup>b</sup>	91,723 <sup>b</sup>	91,984 <sup>b</sup>	93,143 <sup>b</sup>	93,431 <sup>b</sup>	96,435 <sup>b</sup>	98,296 <sup>b</sup>	100,265	102,093
Cocaine	33,239 <sup>b</sup>	34,243 <sup>b</sup>	33,542 <sup>b</sup>	33,090 <sup>b</sup>	34,728 <sup>a</sup>	35,340	36,301	36,197	36,848
Crack	8,239	7,792 <sup>b</sup>	7,715 <sup>b</sup>	7,820 <sup>b</sup>	8,426	8,475	8,366	8,301	9,099
Heroin	3,574	3,664	3,067 <sup>b</sup>	3,475	3,725	3,726	3,720	3,627	4,080
Hallucinogens	32,895 <sup>b</sup>	33,113 <sup>b</sup>	33,173 <sup>b</sup>	32,730 <sup>b</sup>	34,296 <sup>a</sup>	33,319 <sup>b</sup>	34,990	36,244	36,477
LSD	23,845	24,027	23,085	22,145	23,128	22,457	23,278	23,386	23,089
PCP	7,185 <sup>a</sup>	6,903	6,587	6,421	6,490	6,035	6,529	6,140	6,181
Ecstasy	9,330 <sup>b</sup>	10,301 <sup>b</sup>	10,607 <sup>b</sup>	11,096 <sup>b</sup>	11,788 <sup>b</sup>	11,979 <sup>b</sup>	12,393 <sup>b</sup>	13,661 <sup>b</sup>	15,282
Inhalants	20,265	20,325	20,036	20,087	20,318	20,055	19,967	20,180	19,655
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	44,556 <sup>b</sup>	45,629 <sup>b</sup>	45,754 <sup>b</sup>	46,552 <sup>a</sup>	47,794	47,467	49,198	48,975	49,058
Pain Relievers	26,836 <sup>b</sup>	28,409 <sup>b</sup>	28,890 <sup>b</sup>	30,175 <sup>b</sup>	30,825 <sup>a</sup>	30,608 <sup>a</sup>	32,515	32,669	32,537
OxyContin <sup>®</sup>	1,700 <sup>b</sup>	2,579 <sup>b</sup>	2,767 <sup>b</sup>	3,205 <sup>b</sup>	3,756 <sup>b</sup>	3,995 <sup>b</sup>	4,502 <sup>b</sup>	5,451	5,755
Tranquilizers	18,433 <sup>b</sup>	19,346 <sup>b</sup>	19,036 <sup>b</sup>	20,275	20,563	19,460 <sup>b</sup>	20,758	21,062	21,381
Stimulants <sup>3</sup>	22,384	21,958	21,391	20,113	21,538	20,922	20,536	21,348	21,131
Methamphetamine <sup>3</sup>	14,949 <sup>b</sup>	14,756 <sup>b</sup>	14,166 <sup>a</sup>	12,344	13,865	12,843	12,404	12,652	12,856
Sedatives	9,723 <sup>b</sup>	9,261 <sup>b</sup>	9,640 <sup>b</sup>	8,751 <sup>b</sup>	8,598 <sup>a</sup>	8,183	8,679 <sup>a</sup>	8,381	7,454
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	65,014 <sup>b</sup>	65,809 <sup>b</sup>	65,302 <sup>b</sup>	66,879 <sup>b</sup>	67,957 <sup>b</sup>	68,857 <sup>a</sup>	71,086	71,307	72,052

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.13B Types of Illicit Drug Use in Lifetime among Persons Aged 18 or Older: Percentages, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	47.8 <sup>a</sup>	48.2	47.6 <sup>b</sup>	48.2	47.5 <sup>b</sup>	48.4	49.3	49.3	49.4
Marijuana and Hashish	42.7 <sup>b</sup>	43.1 <sup>a</sup>	42.7 <sup>b</sup>	42.8 <sup>b</sup>	42.3 <sup>b</sup>	43.3	43.7	44.1	44.5
Cocaine	15.8	16.1	15.6	15.2 <sup>a</sup>	15.7	15.9	16.1	15.9	16.1
Crack	3.9	3.7	3.6	3.6	3.8	3.8	3.7	3.7	4.0
Heroin	1.7	1.7	1.4 <sup>a</sup>	1.6	1.7	1.7	1.7	1.6	1.8
Hallucinogens	15.6	15.6	15.4	15.0 <sup>a</sup>	15.5	15.0 <sup>a</sup>	15.6	16.0	15.9
LSD	11.3 <sup>b</sup>	11.3 <sup>b</sup>	10.7	10.2	10.5	10.1	10.3	10.3	10.1
PCP	3.4 <sup>b</sup>	3.2 <sup>b</sup>	3.1	2.9	2.9	2.7	2.9	2.7	2.7
Ecstasy	4.4 <sup>b</sup>	4.8 <sup>b</sup>	4.9 <sup>b</sup>	5.1 <sup>b</sup>	5.3 <sup>b</sup>	5.4 <sup>b</sup>	5.5 <sup>b</sup>	6.0 <sup>b</sup>	6.7
Inhalants	9.6 <sup>b</sup>	9.6 <sup>b</sup>	9.3 <sup>a</sup>	9.2 <sup>a</sup>	9.2 <sup>a</sup>	9.0	8.9	8.9	8.6
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	21.2	21.5	21.3	21.4	21.7	21.3	21.9	21.6	21.4
Pain Relievers	12.8 <sup>b</sup>	13.4 <sup>a</sup>	13.4 <sup>a</sup>	13.9	14.0	13.8	14.5	14.4	14.2
OxyContin <sup>®</sup>	0.8 <sup>b</sup>	1.2 <sup>b</sup>	1.3 <sup>b</sup>	1.5 <sup>b</sup>	1.7 <sup>b</sup>	1.8 <sup>b</sup>	2.0 <sup>b</sup>	2.4	2.5
Tranquilizers	8.8	9.1	8.8	9.3	9.3	8.7	9.2	9.3	9.3
Stimulants <sup>3</sup>	10.6 <sup>b</sup>	10.3 <sup>b</sup>	9.9 <sup>a</sup>	9.2	9.8	9.4	9.1	9.4	9.2
Methamphetamine <sup>3</sup>	7.1 <sup>b</sup>	6.9 <sup>b</sup>	6.6 <sup>b</sup>	5.7	6.3 <sup>a</sup>	5.8	5.5	5.6	5.6
Sedatives	4.6 <sup>b</sup>	4.4 <sup>b</sup>	4.5 <sup>b</sup>	4.0 <sup>b</sup>	3.9 <sup>b</sup>	3.7	3.9 <sup>a</sup>	3.7 <sup>a</sup>	3.3
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	30.9	30.9	30.3 <sup>a</sup>	30.7	30.8	30.9	31.6	31.4	31.4

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.14A Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	29,638 <sup>b</sup>	29,538 <sup>b</sup>	29,506 <sup>b</sup>	29,999 <sup>b</sup>	30,804 <sup>b</sup>	30,966 <sup>b</sup>	30,799 <sup>b</sup>	33,145	34,075
Marijuana and Hashish	21,850 <sup>b</sup>	21,486 <sup>b</sup>	21,796 <sup>b</sup>	21,991 <sup>b</sup>	22,030 <sup>b</sup>	21,938 <sup>b</sup>	22,531 <sup>b</sup>	25,182	25,806
Cocaine	5,394 <sup>b</sup>	5,470 <sup>b</sup>	5,247 <sup>b</sup>	5,099 <sup>b</sup>	5,658 <sup>b</sup>	5,355 <sup>b</sup>	4,950 <sup>a</sup>	4,557	4,261
Crack	1,454 <sup>b</sup>	1,310 <sup>b</sup>	1,241 <sup>b</sup>	1,327 <sup>b</sup>	1,406 <sup>b</sup>	1,386 <sup>b</sup>	1,075	988	853
Heroin	353 <sup>b</sup>	278 <sup>b</sup>	350 <sup>b</sup>	343 <sup>b</sup>	523	342 <sup>b</sup>	411	571	590
Hallucinogens	3,804	3,158 <sup>b</sup>	3,125 <sup>b</sup>	3,144 <sup>b</sup>	3,307 <sup>a</sup>	3,135 <sup>b</sup>	2,988 <sup>b</sup>	3,774	3,770
LSD	684	409 <sup>b</sup>	434 <sup>b</sup>	413 <sup>b</sup>	560 <sup>a</sup>	495 <sup>b</sup>	619	637	739
PCP	126 <sup>a</sup>	130 <sup>a</sup>	138 <sup>a</sup>	79	128	82	38	75	59
Ecstasy	2,621 <sup>a</sup>	1,790 <sup>a</sup>	1,605 <sup>b</sup>	1,703 <sup>b</sup>	1,824 <sup>a</sup>	1,804 <sup>a</sup>	1,786 <sup>b</sup>	2,383	2,179
Inhalants	1,005	956	1,087	1,043	1,108	1,086	1,069	1,134	1,149
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	12,509 <sup>b</sup>	12,868 <sup>a</sup>	12,621 <sup>b</sup>	13,246	14,301	14,246	13,253	14,120	14,219
Pain Relievers	9,122 <sup>b</sup>	9,753 <sup>a</sup>	9,391 <sup>b</sup>	10,074	10,832	10,785	10,277	10,789	10,691
OxyContin <sup>®</sup>	--	--	1,019 <sup>b</sup>	1,039 <sup>b</sup>	1,124 <sup>b</sup>	1,190 <sup>b</sup>	1,235 <sup>b</sup>	1,454	1,636
Tranquilizers	4,281 <sup>b</sup>	4,465 <sup>a</sup>	4,536 <sup>a</sup>	4,752	4,575	4,776	4,621	5,000	5,115
Stimulants <sup>3</sup>	2,716	2,453	2,747	2,581	3,243 <sup>b</sup>	2,582	2,265	2,727	2,563
Methamphetamine <sup>3</sup>	1,510 <sup>b</sup>	1,417 <sup>b</sup>	1,631 <sup>b</sup>	1,425 <sup>b</sup>	1,704 <sup>b</sup>	1,226 <sup>a</sup>	756	1,079	856
Sedatives	843	700	604	632	817	752	509 <sup>a</sup>	685	796
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	17,086	16,954	16,389 <sup>a</sup>	16,980	18,108	18,230	17,140	18,165	17,822

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.14B Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older: Percentages, 2002-2010**

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	14.1 <sup>a</sup>	13.9 <sup>b</sup>	13.7 <sup>b</sup>	13.8 <sup>b</sup>	14.0 <sup>a</sup>	13.9 <sup>a</sup>	13.7 <sup>b</sup>	14.6	14.9
Marijuana and Hashish	10.4 <sup>b</sup>	10.1 <sup>b</sup>	10.1 <sup>b</sup>	10.1 <sup>b</sup>	10.0 <sup>b</sup>	9.9 <sup>b</sup>	10.0 <sup>b</sup>	11.1	11.3
Cocaine	2.6 <sup>b</sup>	2.6 <sup>b</sup>	2.4 <sup>b</sup>	2.3 <sup>b</sup>	2.6 <sup>b</sup>	2.4 <sup>b</sup>	2.2 <sup>b</sup>	2.0	1.9
Crack	0.7 <sup>b</sup>	0.6 <sup>b</sup>	0.5	0.4	0.4				
Heroin	0.2 <sup>a</sup>	0.1 <sup>b</sup>	0.2 <sup>a</sup>	0.2 <sup>b</sup>	0.2	0.2 <sup>a</sup>	0.2	0.3	0.3
Hallucinogens	1.8	1.5	1.5	1.4 <sup>a</sup>	1.5	1.4 <sup>a</sup>	1.3 <sup>b</sup>	1.7	1.6
LSD	0.3	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.3	0.2 <sup>b</sup>	0.3	0.3	0.3
PCP	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.1 <sup>a</sup>	0.0	0.1	0.0	0.0	0.0	0.0
Ecstasy	1.2 <sup>b</sup>	0.8	0.7 <sup>b</sup>	0.8 <sup>a</sup>	0.8	0.8	0.8 <sup>a</sup>	1.0	1.0
Inhalants	0.5	0.4	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	5.9	6.1	5.9	6.1	6.5	6.4	5.9	6.2	6.2
Pain Relievers	4.3	4.6	4.4	4.6	4.9	4.8	4.6	4.7	4.7
OxyContin <sup>®</sup>	--	--	0.5 <sup>b</sup>	0.5 <sup>b</sup>	0.5 <sup>b</sup>	0.5 <sup>b</sup>	0.5 <sup>a</sup>	0.6	0.7
Tranquilizers	2.0	2.1	2.1	2.2	2.1	2.1	2.1	2.2	2.2
Stimulants <sup>3</sup>	1.3	1.2	1.3	1.2	1.5 <sup>b</sup>	1.2	1.0	1.2	1.1
Methamphetamine <sup>3</sup>	0.7 <sup>b</sup>	0.7 <sup>b</sup>	0.8 <sup>b</sup>	0.7 <sup>b</sup>	0.8 <sup>b</sup>	0.6 <sup>a</sup>	0.3	0.5	0.4
Sedatives	0.4	0.3	0.3	0.3	0.4	0.3	0.2 <sup>a</sup>	0.3	0.3
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	8.1	8.0	7.6	7.8	8.2	8.2	7.6	8.0	7.8

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.15A Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older: Numbers in Thousands, 2002-2010**

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	16,644 <sup>b</sup>	16,659 <sup>b</sup>	16,397 <sup>b</sup>	17,209 <sup>b</sup>	17,876 <sup>b</sup>	17,448 <sup>b</sup>	17,773 <sup>b</sup>	19,342	20,156
Marijuana and Hashish	12,561 <sup>b</sup>	12,667 <sup>b</sup>	12,667 <sup>b</sup>	12,896 <sup>b</sup>	13,124 <sup>b</sup>	12,762 <sup>b</sup>	13,546 <sup>b</sup>	14,919	15,578
Cocaine	1,874 <sup>b</sup>	2,128 <sup>b</sup>	1,895 <sup>b</sup>	2,255 <sup>b</sup>	2,311 <sup>b</sup>	1,972 <sup>b</sup>	1,757 <sup>a</sup>	1,568	1,406
Crack	541	572	451	657 <sup>b</sup>	689 <sup>b</sup>	594 <sup>a</sup>	352	484	375
Heroin	155	101 <sup>a</sup>	151	121 <sup>a</sup>	322	150	199	181	230
Hallucinogens	941	783 <sup>a</sup>	719 <sup>b</sup>	883	825	821	818	1,041	993
LSD	51 <sup>b</sup>	82	103	72	99	115	104	123	108
PCP	35	35	37	31	18	32	*	40	29
Ecstasy	553	379 <sup>a</sup>	373 <sup>b</sup>	428	446	438	450	644	571
Inhalants	334	250	325	298	421	321	372	315	423
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	5,300 <sup>b</sup>	5,453 <sup>a</sup>	5,196 <sup>b</sup>	5,653	6,251	6,066	5,514 <sup>a</sup>	6,182	6,242
Pain Relievers	3,583 <sup>b</sup>	3,892 <sup>a</sup>	3,653 <sup>b</sup>	3,972	4,530	4,505	4,180	4,604	4,490
OxyContin <sup>®</sup>	--	--	260 <sup>b</sup>	299 <sup>b</sup>	248 <sup>b</sup>	322 <sup>a</sup>	386	441	517
Tranquilizers	1,617 <sup>a</sup>	1,604 <sup>a</sup>	1,455 <sup>b</sup>	1,657 <sup>a</sup>	1,640 <sup>a</sup>	1,652	1,650 <sup>a</sup>	1,866	2,026
Stimulants <sup>3</sup>	1,094	1,088	1,132	1,016	1,214	926	791	1,161	969
Methamphetamine <sup>3</sup>	612 <sup>b</sup>	655 <sup>b</sup>	648 <sup>b</sup>	556 <sup>b</sup>	677 <sup>b</sup>	496	285	466	328
Sedatives	389	247	239	245	340	312	199	330	339
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	7,356	7,432	6,920 <sup>b</sup>	7,710	8,364	8,089	7,474	8,038	7,922

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.15B Types of Illicit Drug Use in the Past Month among Persons Aged 18 or Older: Percentages, 2002-2010**

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	7.9 <sup>b</sup>	7.8 <sup>b</sup>	7.6 <sup>b</sup>	7.9 <sup>b</sup>	8.1 <sup>a</sup>	7.8 <sup>b</sup>	7.9 <sup>b</sup>	8.5	8.8
Marijuana and Hashish	6.0 <sup>b</sup>	6.0 <sup>b</sup>	5.9 <sup>b</sup>	5.9 <sup>b</sup>	5.9 <sup>b</sup>	5.7 <sup>b</sup>	6.0 <sup>b</sup>	6.6	6.8
Cocaine	0.9 <sup>b</sup>	1.0 <sup>b</sup>	0.9 <sup>b</sup>	1.0 <sup>b</sup>	1.0 <sup>b</sup>	0.9 <sup>b</sup>	0.8 <sup>a</sup>	0.7	0.6
Crack	0.3 <sup>a</sup>	0.3 <sup>a</sup>	0.2	0.3 <sup>b</sup>	0.3 <sup>b</sup>	0.3 <sup>a</sup>	0.2	0.2	0.2
Heroin	0.1	0.0 <sup>a</sup>	0.1	0.1 <sup>a</sup>	0.1	0.1	0.1	0.1	0.1
Hallucinogens	0.4	0.4	0.3 <sup>a</sup>	0.4	0.4	0.4	0.4	0.5	0.4
LSD	0.0 <sup>a</sup>	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.0
PCP	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Ecstasy	0.3	0.2 <sup>a</sup>	0.2 <sup>a</sup>	0.2	0.2	0.2	0.2	0.3	0.2
Inhalants	0.2	0.1	0.2	0.1	0.2	0.1	0.2	0.1	0.2
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	2.5	2.6	2.4 <sup>a</sup>	2.6	2.8	2.7	2.5	2.7	2.7
Pain Relievers	1.7 <sup>a</sup>	1.8	1.7 <sup>a</sup>	1.8	2.1	2.0	1.9	2.0	2.0
OxyContin <sup>®</sup>	--	--	0.1 <sup>b</sup>	0.1 <sup>b</sup>	0.1 <sup>b</sup>	0.1 <sup>a</sup>	0.2	0.2	0.2
Tranquilizers	0.8	0.8	0.7 <sup>b</sup>	0.8	0.7	0.7	0.7	0.8	0.9
Stimulants <sup>3</sup>	0.5	0.5	0.5	0.5	0.6 <sup>a</sup>	0.4	0.4	0.5	0.4
Methamphetamine <sup>3</sup>	0.3 <sup>b</sup>	0.2	0.1	0.2	0.1				
Sedatives	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	3.5	3.5	3.2	3.5	3.8	3.6	3.3	3.5	3.5

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.16A Types of Illicit Drug Use in Lifetime among Males Aged 12 to 17: Numbers in Thousands, 2002-2010**

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	4,035 <sup>b</sup>	3,968 <sup>b</sup>	3,956 <sup>b</sup>	3,679 <sup>b</sup>	3,653 <sup>b</sup>	3,489 <sup>a</sup>	3,310	3,506 <sup>b</sup>	3,247
Marijuana and Hashish	2,714 <sup>b</sup>	2,569 <sup>b</sup>	2,552 <sup>b</sup>	2,348	2,310	2,226	2,184	2,340	2,260
Cocaine	337 <sup>b</sup>	319 <sup>b</sup>	339 <sup>b</sup>	283 <sup>b</sup>	256 <sup>a</sup>	257 <sup>a</sup>	217	218	189
Crack	61	89 <sup>b</sup>	60	48	51	43	31	27	34
Heroin	34	34	43	32	25	28	37	25	23
Hallucinogens	740 <sup>b</sup>	618 <sup>a</sup>	616 <sup>a</sup>	518	510	477	501	533	514
LSD	364 <sup>b</sup>	206 <sup>b</sup>	174 <sup>a</sup>	177 <sup>a</sup>	124	109	146	139	120
PCP	106 <sup>b</sup>	100 <sup>b</sup>	97 <sup>b</sup>	97 <sup>b</sup>	53 <sup>a</sup>	52 <sup>a</sup>	56 <sup>a</sup>	44	27
Ecstasy	376 <sup>b</sup>	257	243	181 <sup>b</sup>	201 <sup>b</sup>	215	227	276	278
Inhalants	1,426 <sup>b</sup>	1,390 <sup>b</sup>	1,429 <sup>b</sup>	1,360 <sup>b</sup>	1,301 <sup>b</sup>	1,220 <sup>b</sup>	1,072	1,103 <sup>a</sup>	982
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	1,643 <sup>b</sup>	1,639 <sup>b</sup>	1,616 <sup>b</sup>	1,444 <sup>b</sup>	1,557 <sup>b</sup>	1,436 <sup>b</sup>	1,255	1,345 <sup>a</sup>	1,200
Pain Relievers	1,362 <sup>b</sup>	1,391 <sup>b</sup>	1,378 <sup>b</sup>	1,211 <sup>a</sup>	1,312 <sup>b</sup>	1,195 <sup>a</sup>	1,064	1,169	1,055
OxyContin <sup>®</sup>	112 <sup>b</sup>	146	162	136	158	172	159	188	183
Tranquilizers	385	398 <sup>a</sup>	353	342	315	323	316	301	324
Stimulants <sup>3</sup>	469 <sup>b</sup>	472 <sup>b</sup>	387 <sup>b</sup>	364 <sup>b</sup>	398 <sup>b</sup>	301 <sup>a</sup>	246	235	229
Methamphetamine <sup>3</sup>	150 <sup>b</sup>	160 <sup>b</sup>	140 <sup>b</sup>	133 <sup>b</sup>	126 <sup>a</sup>	81	66	62	76
Sedatives	84	126 <sup>a</sup>	128 <sup>b</sup>	110	97	89	89	102	76
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	2,726 <sup>b</sup>	2,711 <sup>b</sup>	2,721 <sup>b</sup>	2,488 <sup>b</sup>	2,522 <sup>b</sup>	2,369 <sup>b</sup>	2,164	2,239 <sup>a</sup>	2,028

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.16B Types of Illicit Drug Use in Lifetime among Males Aged 12 to 17: Percentages, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	31.9 <sup>b</sup>	31.1 <sup>b</sup>	30.7 <sup>b</sup>	28.4 <sup>b</sup>	28.2 <sup>a</sup>	27.1	26.0	27.9 <sup>a</sup>	26.1
Marijuana and Hashish	21.5 <sup>b</sup>	20.1 <sup>b</sup>	19.8 <sup>a</sup>	18.1	17.8	17.3	17.2	18.6	18.1
Cocaine	2.7 <sup>b</sup>	2.5 <sup>b</sup>	2.6 <sup>b</sup>	2.2 <sup>b</sup>	2.0 <sup>a</sup>	2.0	1.7	1.7	1.5
Crack	0.5	0.7 <sup>b</sup>	0.5	0.4	0.4	0.3	0.2	0.2	0.3
Heroin	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.2	0.2
Hallucinogens	5.8 <sup>b</sup>	4.8 <sup>a</sup>	4.8	4.0	3.9	3.7	3.9	4.2	4.1
LSD	2.9 <sup>b</sup>	1.6 <sup>b</sup>	1.4 <sup>a</sup>	1.4 <sup>a</sup>	1.0	0.8	1.1	1.1	1.0
PCP	0.8 <sup>b</sup>	0.8 <sup>b</sup>	0.8 <sup>b</sup>	0.7 <sup>b</sup>	0.4 <sup>a</sup>	0.4 <sup>a</sup>	0.4 <sup>a</sup>	0.3	0.2
Ecstasy	3.0 <sup>b</sup>	2.0	1.9	1.4 <sup>b</sup>	1.5 <sup>b</sup>	1.7 <sup>a</sup>	1.8	2.2	2.2
Inhalants	11.3 <sup>b</sup>	10.9 <sup>b</sup>	11.1 <sup>b</sup>	10.5 <sup>b</sup>	10.0 <sup>b</sup>	9.5 <sup>b</sup>	8.4	8.8	7.9
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	13.0 <sup>b</sup>	12.8 <sup>b</sup>	12.6 <sup>b</sup>	11.1 <sup>b</sup>	12.0 <sup>b</sup>	11.1 <sup>b</sup>	9.9	10.7 <sup>a</sup>	9.6
Pain Relievers	10.8 <sup>b</sup>	10.9 <sup>b</sup>	10.7 <sup>b</sup>	9.4	10.1 <sup>b</sup>	9.3	8.4	9.3	8.5
OxyContin <sup>®</sup>	0.9 <sup>b</sup>	1.1	1.3	1.0 <sup>a</sup>	1.2	1.3	1.2	1.5	1.5
Tranquilizers	3.0	3.1	2.7	2.6	2.4	2.5	2.5	2.4	2.6
Stimulants <sup>3</sup>	3.7 <sup>b</sup>	3.7 <sup>b</sup>	3.0 <sup>b</sup>	2.8 <sup>b</sup>	3.1 <sup>b</sup>	2.3 <sup>a</sup>	1.9	1.9	1.8
Methamphetamine <sup>3</sup>	1.2 <sup>b</sup>	1.3 <sup>b</sup>	1.1 <sup>b</sup>	1.0 <sup>a</sup>	1.0 <sup>a</sup>	0.6	0.5	0.5	0.6
Sedatives	0.7	1.0 <sup>a</sup>	1.0 <sup>b</sup>	0.8	0.8	0.7	0.7	0.8	0.6
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	21.6 <sup>b</sup>	21.2 <sup>b</sup>	21.1 <sup>b</sup>	19.2 <sup>b</sup>	19.4 <sup>b</sup>	18.4 <sup>b</sup>	17.0	17.8 <sup>a</sup>	16.3

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.17A Types of Illicit Drug Use in Lifetime among Females Aged 12 to 17: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLICIT DRUGS<sup>1</sup></b>	3,607 <sup>b</sup>	3,663 <sup>b</sup>	3,610 <sup>b</sup>	3,342 <sup>b</sup>	3,357 <sup>b</sup>	3,132	3,210 <sup>a</sup>	3,082	3,003
Marijuana and Hashish	2,391 <sup>b</sup>	2,318 <sup>b</sup>	2,236 <sup>b</sup>	2,055 <sup>a</sup>	2,084 <sup>a</sup>	1,858	1,924	1,840	1,879
Cocaine	333 <sup>b</sup>	330 <sup>b</sup>	272 <sup>b</sup>	300 <sup>b</sup>	314 <sup>b</sup>	285 <sup>b</sup>	255 <sup>b</sup>	185	173
Crack	102 <sup>b</sup>	68 <sup>b</sup>	66 <sup>b</sup>	60 <sup>b</sup>	77 <sup>b</sup>	64 <sup>b</sup>	48	30	26
Heroin	60 <sup>b</sup>	45 <sup>a</sup>	35	28	35	26	31	32	23
Hallucinogens	680 <sup>b</sup>	632 <sup>b</sup>	543	480	476	419	472	479	484
LSD	306 <sup>b</sup>	191 <sup>b</sup>	139	111	94	90	122	110	111
PCP	127 <sup>b</sup>	104 <sup>b</sup>	77	85 <sup>a</sup>	75	52	46	55	50
Ecstasy	444 <sup>b</sup>	346	280	218 <sup>b</sup>	273	232 <sup>b</sup>	304	297	341
Inhalants	1,179 <sup>a</sup>	1,281 <sup>b</sup>	1,333 <sup>b</sup>	1,298 <sup>b</sup>	1,261 <sup>b</sup>	1,202 <sup>b</sup>	1,235 <sup>b</sup>	1,164 <sup>a</sup>	1,023
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	1,759 <sup>b</sup>	1,733 <sup>b</sup>	1,787 <sup>b</sup>	1,575 <sup>b</sup>	1,613 <sup>b</sup>	1,513	1,517	1,450	1,383
Pain Relievers	1,412 <sup>b</sup>	1,407 <sup>b</sup>	1,500 <sup>b</sup>	1,305	1,335 <sup>a</sup>	1,256	1,282	1,208	1,183
OxyContin <sup>®</sup>	112 <sup>b</sup>	108 <sup>b</sup>	143	140	184	188	182	189	184
Tranquilizers	450	476 <sup>a</sup>	464	424	426	425	402	392	397
Stimulants <sup>3</sup>	643 <sup>b</sup>	574 <sup>b</sup>	519 <sup>b</sup>	506 <sup>b</sup>	532 <sup>b</sup>	431 <sup>b</sup>	424 <sup>b</sup>	346	300
Methamphetamine <sup>3</sup>	266 <sup>b</sup>	222 <sup>b</sup>	206 <sup>b</sup>	185 <sup>b</sup>	216 <sup>b</sup>	142 <sup>b</sup>	129 <sup>a</sup>	123 <sup>a</sup>	81
Sedatives	154 <sup>a</sup>	123	123	122	126	125	114	121	101
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	2,559 <sup>b</sup>	2,607 <sup>b</sup>	2,634 <sup>b</sup>	2,456 <sup>b</sup>	2,427 <sup>b</sup>	2,269	2,324 <sup>a</sup>	2,235	2,122

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.17B Types of Illicit Drug Use in Lifetime among Females Aged 12 to 17: Percentages, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLICIT DRUGS<sup>1</sup></b>	29.8 <sup>b</sup>	30.0 <sup>b</sup>	29.3 <sup>b</sup>	26.9 <sup>a</sup>	27.0 <sup>a</sup>	25.4	26.3	25.6	25.3
Marijuana and Hashish	19.7 <sup>b</sup>	19.0 <sup>b</sup>	18.1 <sup>b</sup>	16.6	16.8	15.0	15.8	15.3	15.8
Cocaine	2.8 <sup>b</sup>	2.7 <sup>b</sup>	2.2 <sup>b</sup>	2.4 <sup>b</sup>	2.5 <sup>b</sup>	2.3 <sup>b</sup>	2.1 <sup>a</sup>	1.5	1.5
Crack	0.8 <sup>b</sup>	0.6 <sup>b</sup>	0.5 <sup>b</sup>	0.5 <sup>b</sup>	0.6 <sup>b</sup>	0.5 <sup>b</sup>	0.4	0.2	0.2
Heroin	0.5 <sup>b</sup>	0.4 <sup>a</sup>	0.3	0.2	0.3	0.2	0.3	0.3	0.2
Hallucinogens	5.6 <sup>b</sup>	5.2 <sup>b</sup>	4.4	3.9	3.8	3.4 <sup>a</sup>	3.9	4.0	4.1
LSD	2.5 <sup>b</sup>	1.6 <sup>b</sup>	1.1	0.9	0.8	0.7	1.0	0.9	0.9
PCP	1.0 <sup>b</sup>	0.8 <sup>b</sup>	0.6	0.7 <sup>a</sup>	0.6	0.4	0.4	0.5	0.4
Ecstasy	3.7 <sup>a</sup>	2.8	2.3 <sup>a</sup>	1.8 <sup>b</sup>	2.2 <sup>a</sup>	1.9 <sup>b</sup>	2.5	2.5	2.9
Inhalants	9.7 <sup>a</sup>	10.5 <sup>b</sup>	10.8 <sup>b</sup>	10.5 <sup>b</sup>	10.1 <sup>b</sup>	9.7 <sup>a</sup>	10.1 <sup>b</sup>	9.7 <sup>a</sup>	8.6
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	14.5 <sup>b</sup>	14.2 <sup>b</sup>	14.5 <sup>b</sup>	12.7	13.0 <sup>a</sup>	12.3	12.4	12.0	11.6
Pain Relievers	11.7 <sup>b</sup>	11.5 <sup>b</sup>	12.2 <sup>b</sup>	10.5	10.7	10.2	10.5	10.0	10.0
OxyContin <sup>®</sup>	0.9 <sup>b</sup>	0.9 <sup>b</sup>	1.2	1.1 <sup>a</sup>	1.5	1.5	1.5	1.6	1.5
Tranquilizers	3.7	3.9	3.8	3.4	3.4	3.4	3.3	3.3	3.3
Stimulants <sup>3</sup>	5.3 <sup>b</sup>	4.7 <sup>b</sup>	4.2 <sup>b</sup>	4.1 <sup>b</sup>	4.3 <sup>b</sup>	3.5 <sup>b</sup>	3.5 <sup>b</sup>	2.9	2.5
Methamphetamine <sup>3</sup>	2.2 <sup>b</sup>	1.8 <sup>b</sup>	1.7 <sup>b</sup>	1.5 <sup>b</sup>	1.7 <sup>b</sup>	1.1 <sup>b</sup>	1.1	1.0	0.7
Sedatives	1.3 <sup>a</sup>	1.0	1.0	1.0	1.0	1.0	0.9	1.0	0.9
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	21.1 <sup>b</sup>	21.3 <sup>b</sup>	21.4 <sup>b</sup>	19.8 <sup>b</sup>	19.5 <sup>a</sup>	18.4	19.1	18.6	17.9

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.18A Types of Illicit Drug Use in the Past Year among Males Aged 12 to 17: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	2,842 <sup>b</sup>	2,776 <sup>b</sup>	2,653 <sup>b</sup>	2,557	2,525	2,499	2,419	2,539	2,419
Marijuana and Hashish	2,064 <sup>b</sup>	1,955	1,918	1,796	1,744	1,748	1,724	1,881	1,854
Cocaine	263 <sup>b</sup>	222 <sup>b</sup>	223 <sup>b</sup>	208 <sup>b</sup>	178 <sup>b</sup>	174 <sup>a</sup>	159	145	115
Crack	36 <sup>b</sup>	47 <sup>b</sup>	31 <sup>b</sup>	23 <sup>a</sup>	32 <sup>b</sup>	25 <sup>a</sup>	18	16	8
Heroin	20	17	32	17	14	16	27	17	13
Hallucinogens	501 <sup>b</sup>	386	389	352	318	339	376	404	382
LSD	183 <sup>b</sup>	73	94	92	51	76	116 <sup>a</sup>	89	73
PCP	46 <sup>b</sup>	44 <sup>b</sup>	39 <sup>b</sup>	44 <sup>b</sup>	26	32 <sup>a</sup>	35 <sup>b</sup>	26	13
Ecstasy	262	142 <sup>a</sup>	141 <sup>a</sup>	125 <sup>b</sup>	129 <sup>b</sup>	165	164	211	207
Inhalants	586 <sup>b</sup>	554 <sup>b</sup>	587 <sup>b</sup>	538 <sup>b</sup>	521 <sup>b</sup>	500 <sup>b</sup>	467 <sup>a</sup>	455 <sup>a</sup>	380
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	1,091 <sup>b</sup>	1,090 <sup>b</sup>	974 <sup>b</sup>	962 <sup>b</sup>	1,052 <sup>b</sup>	970 <sup>a</sup>	863	876	811
Pain Relievers	913 <sup>b</sup>	930 <sup>b</sup>	837 <sup>b</sup>	818 <sup>a</sup>	882 <sup>b</sup>	816 <sup>a</sup>	728	751	693
OxyContin <sup>®</sup>	--	--	99	101	95	107	122	109	113
Tranquilizers	249	258	226	229	195	215	225	208	200
Stimulants <sup>3</sup>	294 <sup>b</sup>	253 <sup>b</sup>	210 <sup>b</sup>	194 <sup>a</sup>	224 <sup>b</sup>	173	141	140	133
Methamphetamine <sup>3</sup>	97 <sup>b</sup>	82 <sup>a</sup>	69	78 <sup>a</sup>	74	39	33	29	44
Sedatives	45	67	59	51	56	42	46	70 <sup>a</sup>	41
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	1,683 <sup>b</sup>	1,637 <sup>b</sup>	1,537 <sup>b</sup>	1,479 <sup>b</sup>	1,540 <sup>b</sup>	1,444 <sup>a</sup>	1,377	1,370	1,268

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.18B Types of Illicit Drug Use in the Past Year among Males Aged 12 to 17: Percentages, 2002-2010**

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	22.5 <sup>b</sup>	21.7 <sup>b</sup>	20.6	19.7	19.5	19.4	19.0	20.2	19.4
Marijuana and Hashish	16.3 <sup>a</sup>	15.3	14.9	13.9	13.4 <sup>a</sup>	13.6 <sup>a</sup>	13.6 <sup>a</sup>	15.0	14.9
Cocaine	2.1 <sup>b</sup>	1.7 <sup>b</sup>	1.7 <sup>b</sup>	1.6 <sup>b</sup>	1.4 <sup>a</sup>	1.3 <sup>a</sup>	1.3	1.2	0.9
Crack	0.3 <sup>b</sup>	0.4 <sup>b</sup>	0.2 <sup>b</sup>	0.2	0.2 <sup>b</sup>	0.2 <sup>a</sup>	0.1	0.1	0.1
Heroin	0.2	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.1
Hallucinogens	4.0 <sup>b</sup>	3.0	3.0	2.7	2.5 <sup>a</sup>	2.6	3.0	3.2	3.1
LSD	1.5 <sup>b</sup>	0.6	0.7	0.7	0.4	0.6	0.9 <sup>a</sup>	0.7	0.6
PCP	0.4 <sup>b</sup>	0.3 <sup>b</sup>	0.3 <sup>b</sup>	0.3 <sup>b</sup>	0.2	0.2 <sup>a</sup>	0.3 <sup>a</sup>	0.2	0.1
Ecstasy	2.1	1.1 <sup>b</sup>	1.1 <sup>b</sup>	1.0 <sup>b</sup>	1.0 <sup>b</sup>	1.3	1.3	1.7	1.7
Inhalants	4.6 <sup>b</sup>	4.3 <sup>b</sup>	4.6 <sup>b</sup>	4.2 <sup>b</sup>	4.0 <sup>b</sup>	3.9 <sup>a</sup>	3.7	3.6	3.1
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	8.6 <sup>b</sup>	8.5 <sup>b</sup>	7.6 <sup>a</sup>	7.4 <sup>a</sup>	8.1 <sup>b</sup>	7.5 <sup>a</sup>	6.8	7.0	6.5
Pain Relievers	7.2 <sup>b</sup>	7.3 <sup>b</sup>	6.5 <sup>a</sup>	6.3	6.8 <sup>b</sup>	6.3	5.7	6.0	5.6
OxyContin <sup>®</sup>	--	--	0.8	0.8	0.7	0.8	1.0	0.9	0.9
Tranquilizers	2.0	2.0	1.8	1.8	1.5	1.7	1.8	1.7	1.6
Stimulants <sup>3</sup>	2.3 <sup>b</sup>	2.0 <sup>b</sup>	1.6 <sup>b</sup>	1.5 <sup>a</sup>	1.7 <sup>b</sup>	1.3	1.1	1.1	1.1
Methamphetamine <sup>3</sup>	0.8 <sup>b</sup>	0.6 <sup>a</sup>	0.5	0.6	0.6	0.3	0.3	0.2	0.4
Sedatives	0.4	0.5	0.5	0.4	0.4	0.3	0.4	0.6 <sup>a</sup>	0.3
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	13.3 <sup>b</sup>	12.8 <sup>b</sup>	11.9 <sup>b</sup>	11.4 <sup>a</sup>	11.9 <sup>b</sup>	11.2	10.8	10.9	10.2

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.19A Types of Illicit Drug Use in the Past Year among Females Aged 12 to 17: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	2,652 <sup>b</sup>	2,679 <sup>b</sup>	2,648 <sup>b</sup>	2,484 <sup>a</sup>	2,447	2,228	2,307	2,270	2,312
Marijuana and Hashish	1,841 <sup>b</sup>	1,790 <sup>b</sup>	1,738 <sup>a</sup>	1,587	1,604	1,399 <sup>a</sup>	1,514	1,457	1,547
Cocaine	245 <sup>b</sup>	216 <sup>b</sup>	188 <sup>a</sup>	215 <sup>b</sup>	233 <sup>b</sup>	209 <sup>b</sup>	146	95	124
Crack	64 <sup>b</sup>	48 <sup>b</sup>	32 <sup>a</sup>	31 <sup>a</sup>	41 <sup>b</sup>	40 <sup>b</sup>	16	12	10
Heroin	31	19	16	20	23	8	15	17	15
Hallucinogens	444 <sup>a</sup>	392	364	313	332	288 <sup>a</sup>	314	330	366
LSD	131 <sup>b</sup>	76	63	59	55	50	67	53	62
PCP	64 <sup>b</sup>	44 <sup>a</sup>	33	41	33	23	26	22	23
Ecstasy	285	188 <sup>a</sup>	169 <sup>b</sup>	131 <sup>b</sup>	177 <sup>b</sup>	163 <sup>b</sup>	189 <sup>a</sup>	205	259
Inhalants	492	565	581	606 <sup>a</sup>	590 <sup>a</sup>	494	512	501	501
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	1,194 <sup>b</sup>	1,205 <sup>b</sup>	1,254 <sup>b</sup>	1,139 <sup>a</sup>	1,129 <sup>a</sup>	1,064	1,050	1,009	1,001
Pain Relievers	957 <sup>a</sup>	987 <sup>b</sup>	1,028 <sup>b</sup>	922	936	864	880	866	829
OxyContin <sup>®</sup>	--	--	95	87	104	125	102	114	121
Tranquilizers	319	328	307	268	288	291	257	251	266
Stimulants <sup>3</sup>	371 <sup>b</sup>	325 <sup>b</sup>	297 <sup>b</sup>	312 <sup>b</sup>	324 <sup>b</sup>	242	233	193	191
Methamphetamine <sup>3</sup>	148 <sup>b</sup>	103 <sup>b</sup>	108 <sup>b</sup>	100 <sup>a</sup>	111 <sup>b</sup>	79	61	56	54
Sedatives	93	64	75	67	53	69	66	56	69
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	1,655 <sup>a</sup>	1,714 <sup>b</sup>	1,732 <sup>b</sup>	1,650 <sup>a</sup>	1,605	1,470	1,473	1,465	1,486

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.19B Types of Illicit Drug Use in the Past Year among Females Aged 12 to 17: Percentages, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	21.9 <sup>b</sup>	21.9 <sup>b</sup>	21.5 <sup>b</sup>	20.0	19.7	18.0 <sup>a</sup>	18.9	18.9	19.4
Marijuana and Hashish	15.2 <sup>b</sup>	14.6 <sup>a</sup>	14.1	12.8	12.9	11.3 <sup>b</sup>	12.4	12.1	13.0
Cocaine	2.0 <sup>b</sup>	1.8 <sup>b</sup>	1.5 <sup>a</sup>	1.7 <sup>b</sup>	1.9 <sup>b</sup>	1.7 <sup>b</sup>	1.2	0.8	1.0
Crack	0.5 <sup>b</sup>	0.4 <sup>b</sup>	0.3 <sup>a</sup>	0.3 <sup>a</sup>	0.3 <sup>b</sup>	0.3 <sup>b</sup>	0.1	0.1	0.1
Heroin	0.3	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.1
Hallucinogens	3.7	3.2	3.0	2.5	2.7	2.3 <sup>a</sup>	2.6	2.7	3.1
LSD	1.1 <sup>b</sup>	0.6	0.5	0.5	0.4	0.4	0.5	0.4	0.5
PCP	0.5 <sup>b</sup>	0.4 <sup>a</sup>	0.3	0.3	0.3	0.2	0.2	0.2	0.2
Ecstasy	2.4	1.5 <sup>a</sup>	1.4 <sup>b</sup>	1.1 <sup>b</sup>	1.4 <sup>b</sup>	1.3 <sup>b</sup>	1.6 <sup>a</sup>	1.7	2.2
Inhalants	4.1	4.6	4.7	4.9	4.7	4.0	4.2	4.2	4.2
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	9.9 <sup>b</sup>	9.9 <sup>b</sup>	10.2 <sup>b</sup>	9.2	9.1	8.6	8.6	8.4	8.4
Pain Relievers	7.9 <sup>a</sup>	8.1 <sup>a</sup>	8.3 <sup>b</sup>	7.4	7.5	7.0	7.2	7.2	7.0
OxyContin <sup>®</sup>	--	--	0.8	0.7	0.8	1.0	0.8	0.9	1.0
Tranquilizers	2.6	2.7	2.5	2.2	2.3	2.4	2.1	2.1	2.2
Stimulants <sup>3</sup>	3.1 <sup>b</sup>	2.7 <sup>b</sup>	2.4 <sup>b</sup>	2.5 <sup>b</sup>	2.6 <sup>b</sup>	2.0	1.9	1.6	1.6
Methamphetamine <sup>3</sup>	1.2 <sup>b</sup>	0.8 <sup>b</sup>	0.9 <sup>b</sup>	0.8 <sup>a</sup>	0.9 <sup>b</sup>	0.6	0.5	0.5	0.5
Sedatives	0.8	0.5	0.6	0.5	0.4	0.6	0.5	0.5	0.6
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	13.7	14.0 <sup>a</sup>	14.0 <sup>a</sup>	13.3	12.9	11.9	12.1	12.2	12.5

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.20A Types of Illicit Drug Use in the Past Month among Males Aged 12 to 17: Numbers in Thousands, 2002-2010**

<b>Drug</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>ILLCIT DRUGS<sup>1</sup></b>	1,560 <sup>b</sup>	1,457 <sup>a</sup>	1,360	1,313	1,272	1,291	1,201	1,335	1,300
Marijuana and Hashish	1,152	1,093	1,037	967	888 <sup>a</sup>	968	921	1,045	1,038
Cocaine	77 <sup>b</sup>	70 <sup>a</sup>	75 <sup>a</sup>	79 <sup>b</sup>	48	48	60	43	37
Crack	10	22	9	14	5	4	6	5	*
Heroin	6	8	11 <sup>a</sup>	11	3	3	8	8	2
Hallucinogens	130	138	120	98	86	102	145	128	116
LSD	37	27	23	24	10 <sup>a</sup>	24	34	21	29
PCP	11	14	7	8	6	9	8	8	4
Ecstasy	58	37	45	32	38	34	58	65	51
Inhalants	159	139	141	160	156	158	135	121	115
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	460 <sup>b</sup>	478 <sup>b</sup>	413 <sup>b</sup>	400 <sup>b</sup>	403 <sup>b</sup>	391 <sup>b</sup>	312	352	288
Pain Relievers	376 <sup>b</sup>	385 <sup>b</sup>	337 <sup>a</sup>	325 <sup>a</sup>	330 <sup>a</sup>	324 <sup>a</sup>	250	300	252
OxyContin <sup>®</sup>	--	--	29	17	18	25	33	40	27
Tranquilizers	70	111 <sup>b</sup>	76	84	46	84	62	58	61
Stimulants <sup>3</sup>	106 <sup>b</sup>	107 <sup>b</sup>	89 <sup>a</sup>	79	80 <sup>a</sup>	48	55	56	49
Methamphetamine <sup>3</sup>	40 <sup>a</sup>	32 <sup>b</sup>	24	39 <sup>b</sup>	24	10	16	18	8
Sedatives	6	27	4	9	21	15	8	25	14
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	705 <sup>b</sup>	682 <sup>b</sup>	626 <sup>b</sup>	616 <sup>b</sup>	603 <sup>b</sup>	582 <sup>a</sup>	534	540	471

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.20B Types of Illicit Drug Use in the Past Month among Males Aged 12 to 17: Percentages, 2002-2010**

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	12.3 <sup>b</sup>	11.4	10.6	10.1	9.8	10.0	9.5	10.6	10.4
Marijuana and Hashish	9.1	8.6	8.1	7.5	6.8 <sup>b</sup>	7.5	7.3 <sup>a</sup>	8.3	8.3
Cocaine	0.6 <sup>a</sup>	0.5 <sup>a</sup>	0.6 <sup>a</sup>	0.6 <sup>a</sup>	0.4	0.4	0.5	0.3	0.3
Crack	0.1	0.2	0.1	0.1	0.0	0.0	0.0	0.0	*
Heroin	0.0	0.1	0.1 <sup>a</sup>	0.1	0.0	0.0	0.1	0.1	0.0
Hallucinogens	1.0	1.1	0.9	0.8	0.7	0.8	1.1	1.0	0.9
LSD	0.3	0.2	0.2	0.2	0.1 <sup>a</sup>	0.2	0.3	0.2	0.2
PCP	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.0
Ecstasy	0.5	0.3	0.4	0.3	0.3	0.3	0.5	0.5	0.4
Inhalants	1.3	1.1	1.1	1.2	1.2	1.2	1.1	1.0	0.9
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	3.6 <sup>b</sup>	3.7 <sup>b</sup>	3.2 <sup>b</sup>	3.1 <sup>b</sup>	3.1 <sup>b</sup>	3.0 <sup>a</sup>	2.5	2.8	2.3
Pain Relievers	3.0 <sup>b</sup>	3.0 <sup>b</sup>	2.6 <sup>a</sup>	2.5	2.5	2.5	2.0	2.4	2.0
OxyContin <sup>®</sup>	--	--	0.2	0.1	0.1	0.2	0.3	0.3	0.2
Tranquilizers	0.6	0.9 <sup>b</sup>	0.6	0.7	0.4	0.6	0.5	0.5	0.5
Stimulants <sup>3</sup>	0.8 <sup>b</sup>	0.8 <sup>b</sup>	0.7 <sup>a</sup>	0.6	0.6	0.4	0.4	0.4	0.4
Methamphetamine <sup>3</sup>	0.3 <sup>a</sup>	0.3 <sup>b</sup>	0.2	0.3 <sup>b</sup>	0.2	0.1	0.1	0.1	0.1
Sedatives	0.0	0.2	0.0	0.1	0.2	0.1	0.1	0.2	0.1
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	5.6 <sup>b</sup>	5.3 <sup>b</sup>	4.9 <sup>b</sup>	4.8 <sup>b</sup>	4.6 <sup>a</sup>	4.5	4.2	4.3	3.8

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.21A Types of Illicit Drug Use in the Past Month among Females Aged 12 to 17: Numbers in Thousands, 2002-2010**

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	1,318 <sup>a</sup>	1,355 <sup>b</sup>	1,314 <sup>a</sup>	1,198	1,210	1,118	1,103	1,136	1,166
Marijuana and Hashish	871 <sup>a</sup>	878 <sup>a</sup>	872 <sup>a</sup>	763	801	718	735	754	758
Cocaine	68 <sup>b</sup>	83 <sup>b</sup>	51 <sup>a</sup>	63 <sup>b</sup>	61 <sup>b</sup>	56 <sup>b</sup>	37	27	24
Crack	16	10	6	12	7	12	1	3	3
Heroin	5	10	4	4	13	*	5	5	6
Hallucinogens	126	122	91	107	95	73	97	89	99
LSD	23	23	15	8	21	6	16	14	16
PCP	11	6	4	9	6	*	5	4	3
Ecstasy	66	54	32 <sup>b</sup>	41 <sup>a</sup>	44	31 <sup>b</sup>	47	50	73
Inhalants	143	181	171	153	185	136	134	124	147
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	527 <sup>a</sup>	519	501	439	441	438	398	420	437
Pain Relievers	418	416	414	361	360	345	317	353	359
OxyContin <sup>®</sup>	--	--	37	17	10	22	16	29	20
Tranquilizers	117 <sup>a</sup>	115 <sup>a</sup>	86	76	80	98	87	85	72
Stimulants <sup>3</sup>	102 <sup>a</sup>	115 <sup>b</sup>	91	93 <sup>a</sup>	90	80	58	73	58
Methamphetamine <sup>3</sup>	31	38 <sup>a</sup>	34	32	29	24	13	18	16
Sedatives	41 <sup>a</sup>	20	22	18	24	19	27	15	21
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	715	735 <sup>a</sup>	702	637	648	599	556	579	624

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

**Table 7.21B Types of Illicit Drug Use in the Past Month among Females Aged 12 to 17: Percentages, 2002-2010**

Drug	2002	2003	2004	2005	2006	2007	2008	2009	2010
<b>ILLCIT DRUGS<sup>1</sup></b>	10.9	11.1 <sup>a</sup>	10.6	9.7	9.7	9.1	9.1	9.4	9.8
Marijuana and Hashish	7.2	7.2	7.1	6.2	6.4	5.8	6.0	6.3	6.4
Cocaine	0.6 <sup>b</sup>	0.7 <sup>b</sup>	0.4 <sup>a</sup>	0.5 <sup>b</sup>	0.5 <sup>b</sup>	0.5 <sup>b</sup>	0.3	0.2	0.2
Crack	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0
Heroin	0.0	0.1	0.0	0.0	0.1	*	0.0	0.0	0.1
Hallucinogens	1.0	1.0	0.7	0.9	0.8	0.6	0.8	0.7	0.8
LSD	0.2	0.2	0.1	0.1	0.2	0.0	0.1	0.1	0.1
PCP	0.1	0.1	0.0	0.1	0.0	*	0.0	0.0	0.0
Ecstasy	0.5	0.4	0.3 <sup>b</sup>	0.3 <sup>a</sup>	0.4 <sup>a</sup>	0.3 <sup>b</sup>	0.4	0.4	0.6
Inhalants	1.2	1.5	1.4	1.2	1.5	1.1	1.1	1.0	1.2
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	4.4	4.2	4.1	3.5	3.6	3.5	3.3	3.5	3.7
Pain Relievers	3.5	3.4	3.4	2.9	2.9	2.8	2.6	2.9	3.0
OxyContin <sup>®</sup>	--	--	0.3	0.1	0.1	0.2	0.1	0.2	0.2
Tranquilizers	1.0 <sup>a</sup>	0.9 <sup>a</sup>	0.7	0.6	0.6	0.8	0.7	0.7	0.6
Stimulants <sup>3</sup>	0.8 <sup>a</sup>	0.9 <sup>b</sup>	0.7	0.8	0.7	0.6	0.5	0.6	0.5
Methamphetamine <sup>3</sup>	0.3	0.3 <sup>a</sup>	0.3	0.3	0.2	0.2	0.1	0.2	0.1
Sedatives	0.3 <sup>a</sup>	0.2	0.2	0.1	0.2	0.2	0.2	0.1	0.2
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	5.9	6.0	5.7	5.1	5.2	4.8	4.6	4.8	5.2

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. For the 2002 through 2005 survey years, a Bernoulli stochastic imputation procedure was used to generate adjusted estimates comparable with estimates for survey years 2006 and later. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.