

**Table 7.22A GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2010**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>LIFETIME USE</b>					
<b>TOTAL</b>	1,632	1,321 <sup>a</sup>	1,206 <sup>b</sup>	1,438	1,690
<b>AGE GROUP</b>					
12 to 17	40	21	20	30	21
18 to 25	473 <sup>b</sup>	306 <sup>a</sup>	204	254	216
26 or Older	1,119	994 <sup>a</sup>	982 <sup>b</sup>	1,154	1,452
<b>GENDER</b>					
Male	985	860 <sup>a</sup>	772 <sup>b</sup>	960	1,210
Female	647	461	434	478	480
<b>PAST YEAR USE</b>					
<b>TOTAL</b>	130	68 <sup>a</sup>	102	99	170
<b>AGE GROUP</b>					
12 to 17	16 <sup>a</sup>	7	8	14 <sup>a</sup>	3
18 to 25	46	42	25	30	30
26 or Older	68	19 <sup>a</sup>	70	55	137
<b>GENDER</b>					
Male	105	55	83	78	123
Female	26	14	19	21	47
<b>PAST MONTH USE</b>					
<b>TOTAL</b>	28	24	28	18	70
<b>AGE GROUP</b>					
12 to 17	5	3	1	5	*
18 to 25	6	21	5	4	6
26 or Older	17	*	22	*	63
<b>GENDER</b>					
Male	24	22	22	8	40
Female	*	*	*	9	30

\*Low precision, no estimate reported.

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.22B GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2010**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>LIFETIME USE</b>					
<b>TOTAL</b>	0.7	0.5	0.5 <sup>a</sup>	0.6	0.7
<b>AGE GROUP</b>					
12 to 17	0.2	0.1	0.1	0.1	0.1
18 to 25	1.4 <sup>b</sup>	0.9 <sup>a</sup>	0.6	0.8	0.6
26 or Older	0.6	0.5 <sup>a</sup>	0.5 <sup>a</sup>	0.6	0.7
<b>GENDER</b>					
Male	0.8	0.7 <sup>a</sup>	0.6 <sup>b</sup>	0.8	1.0
Female	0.5	0.4	0.3	0.4	0.4
<b>PAST YEAR USE</b>					
<b>TOTAL</b>	0.1	0.0 <sup>a</sup>	0.0	0.0	0.1
<b>AGE GROUP</b>					
12 to 17	0.1 <sup>a</sup>	0.0	0.0	0.1 <sup>a</sup>	0.0
18 to 25	0.1	0.1	0.1	0.1	0.1
26 or Older	0.0	0.0 <sup>a</sup>	0.0	0.0	0.1
<b>GENDER</b>					
Male	0.1	0.0	0.1	0.1	0.1
Female	0.0	0.0	0.0	0.0	0.0
<b>PAST MONTH USE</b>					
<b>TOTAL</b>	0.0	0.0	0.0	0.0	0.0
<b>AGE GROUP</b>					
12 to 17	0.0	0.0	0.0	0.0	*
18 to 25	0.0	0.1	0.0	0.0	0.0
26 or Older	0.0	*	0.0	*	0.0
<b>GENDER</b>					
Male	0.0	0.0	0.0	0.0	0.0
Female	*	*	*	0.0	0.0

\*Low precision, no estimate reported.

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.23A Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2010**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>LIFETIME USE</b>					
<b>TOTAL</b>	4,696 <sup>b</sup>	4,937 <sup>b</sup>	5,314 <sup>b</sup>	6,657 <sup>b</sup>	7,891
<b>AGE GROUP</b>					
12 to 17	700	672	615	623	663
18 to 25	2,772 <sup>b</sup>	2,861 <sup>b</sup>	3,069 <sup>b</sup>	3,656 <sup>a</sup>	3,994
26 or Older	1,225 <sup>b</sup>	1,403 <sup>b</sup>	1,631 <sup>b</sup>	2,379 <sup>b</sup>	3,234
<b>GENDER</b>					
Male	2,653 <sup>b</sup>	2,793 <sup>b</sup>	2,934 <sup>b</sup>	3,951 <sup>b</sup>	4,683
Female	2,043 <sup>b</sup>	2,144 <sup>b</sup>	2,381 <sup>b</sup>	2,706 <sup>b</sup>	3,208
<b>PAST YEAR USE</b>					
<b>TOTAL</b>	1,989 <sup>b</sup>	2,003 <sup>b</sup>	2,088 <sup>b</sup>	2,741	2,883
<b>AGE GROUP</b>					
12 to 17	432	395	384	394	413
18 to 25	1,210 <sup>b</sup>	1,215 <sup>b</sup>	1,209 <sup>b</sup>	1,585	1,757
26 or Older	348 <sup>b</sup>	393 <sup>b</sup>	494 <sup>a</sup>	763	713
<b>GENDER</b>					
Male	1,051 <sup>b</sup>	1,157 <sup>b</sup>	1,150 <sup>b</sup>	1,621	1,652
Female	938 <sup>b</sup>	847 <sup>b</sup>	937 <sup>b</sup>	1,120	1,231
<b>PAST MONTH USE</b>					
<b>TOTAL</b>	549 <sup>b</sup>	531 <sup>b</sup>	609 <sup>b</sup>	839	916
<b>AGE GROUP</b>					
12 to 17	118	134	103	118	126
18 to 25	336 <sup>b</sup>	347 <sup>a</sup>	342 <sup>b</sup>	460	496
26 or Older	95 <sup>b</sup>	51 <sup>b</sup>	164 <sup>a</sup>	261	293
<b>GENDER</b>					
Male	296 <sup>b</sup>	299 <sup>b</sup>	343 <sup>b</sup>	494	544
Female	253 <sup>a</sup>	232 <sup>b</sup>	266 <sup>a</sup>	345	372

\*Low precision, no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.23B Nonmedical Use of Adderall<sup>®</sup> in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2010**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>LIFETIME USE</b>					
<b>TOTAL</b>	1.9 <sup>b</sup>	2.0 <sup>b</sup>	2.1 <sup>b</sup>	2.6 <sup>b</sup>	3.1
<b>AGE GROUP</b>					
12 to 17	2.8	2.7	2.5	2.5	2.7
18 to 25	8.5 <sup>b</sup>	8.7 <sup>b</sup>	9.3 <sup>b</sup>	10.9	11.7
26 or Older	0.7 <sup>b</sup>	0.7 <sup>b</sup>	0.8 <sup>b</sup>	1.2 <sup>b</sup>	1.7
<b>GENDER</b>					
Male	2.2 <sup>b</sup>	2.3 <sup>b</sup>	2.4 <sup>b</sup>	3.2 <sup>b</sup>	3.8
Female	1.6 <sup>b</sup>	1.7 <sup>b</sup>	1.9 <sup>b</sup>	2.1 <sup>b</sup>	2.5
<b>PAST YEAR USE</b>					
<b>TOTAL</b>	0.8 <sup>b</sup>	0.8 <sup>b</sup>	0.8 <sup>b</sup>	1.1	1.1
<b>AGE GROUP</b>					
12 to 17	1.7	1.6	1.5	1.6	1.7
18 to 25	3.7 <sup>b</sup>	3.7 <sup>b</sup>	3.7 <sup>b</sup>	4.7	5.2
26 or Older	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.3 <sup>a</sup>	0.4	0.4
<b>GENDER</b>					
Male	0.9 <sup>b</sup>	1.0 <sup>b</sup>	0.9 <sup>b</sup>	1.3	1.3
Female	0.7 <sup>b</sup>	0.7 <sup>b</sup>	0.7 <sup>b</sup>	0.9	0.9
<b>PAST MONTH USE</b>					
<b>TOTAL</b>	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.3	0.4
<b>AGE GROUP</b>					
12 to 17	0.5	0.5	0.4	0.5	0.5
18 to 25	1.0 <sup>b</sup>	1.1 <sup>a</sup>	1.0 <sup>a</sup>	1.4	1.5
26 or Older	0.1 <sup>b</sup>	0.0 <sup>b</sup>	0.1 <sup>a</sup>	0.1	0.2
<b>GENDER</b>					
Male	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.3 <sup>b</sup>	0.4	0.4
Female	0.2 <sup>a</sup>	0.2 <sup>b</sup>	0.2 <sup>a</sup>	0.3	0.3

\*Low precision, no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.24A Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2010**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>LIFETIME USE</b>					
<b>TOTAL</b>	4,442 <sup>b</sup>	5,194 <sup>b</sup>	5,499 <sup>a</sup>	6,179	6,426
<b>AGE GROUP</b>					
12 to 17	313 <sup>b</sup>	297 <sup>a</sup>	265	285	230
18 to 25	1,186	1,088 <sup>a</sup>	1,168	1,275	1,277
26 or Older	2,942 <sup>b</sup>	3,808 <sup>b</sup>	4,066 <sup>a</sup>	4,619	4,919
<b>GENDER</b>					
Male	2,136 <sup>b</sup>	2,372 <sup>b</sup>	2,337 <sup>b</sup>	2,910	3,220
Female	2,305 <sup>b</sup>	2,822	3,162	3,269	3,206
<b>PAST YEAR USE</b>					
<b>TOTAL</b>	2,037	2,053	2,054	2,156	2,258
<b>AGE GROUP</b>					
12 to 17	173 <sup>a</sup>	151	138	142	123
18 to 25	535	479	474	526	448
26 or Older	1,330	1,423	1,442	1,488	1,688
<b>GENDER</b>					
Male	993	1,001	932	1,086	1,157
Female	1,044	1,053	1,122	1,070	1,102
<b>PAST MONTH USE</b>					
<b>TOTAL</b>	549	380 <sup>b</sup>	437 <sup>a</sup>	610	814
<b>AGE GROUP</b>					
12 to 17	45	31	28	35	35
18 to 25	97	112	129	131	108
26 or Older	407	238 <sup>b</sup>	280 <sup>b</sup>	444	671
<b>GENDER</b>					
Male	326	135 <sup>b</sup>	204 <sup>a</sup>	307	459
Female	223	245	233	303	356

\*Low precision, no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.24B Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2010**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>LIFETIME USE</b>					
<b>TOTAL</b>	1.8 <sup>b</sup>	2.1 <sup>b</sup>	2.2 <sup>a</sup>	2.5	2.5
<b>AGE GROUP</b>					
12 to 17	1.2 <sup>a</sup>	1.2	1.1	1.2	0.9
18 to 25	3.6	3.3	3.5	3.8	3.7
26 or Older	1.6 <sup>b</sup>	2.0 <sup>b</sup>	2.1 <sup>a</sup>	2.4	2.5
<b>GENDER</b>					
Male	1.8 <sup>b</sup>	2.0 <sup>b</sup>	1.9 <sup>b</sup>	2.4	2.6
Female	1.8 <sup>b</sup>	2.2	2.5	2.5	2.5
<b>PAST YEAR USE</b>					
<b>TOTAL</b>	0.8	0.8	0.8	0.9	0.9
<b>AGE GROUP</b>					
12 to 17	0.7	0.6	0.6	0.6	0.5
18 to 25	1.6	1.5	1.4	1.6	1.3
26 or Older	0.7	0.7	0.8	0.8	0.9
<b>GENDER</b>					
Male	0.8	0.8	0.8	0.9	0.9
Female	0.8	0.8	0.9	0.8	0.8
<b>PAST MONTH USE</b>					
<b>TOTAL</b>	0.2	0.2 <sup>b</sup>	0.2 <sup>a</sup>	0.2	0.3
<b>AGE GROUP</b>					
12 to 17	0.2	0.1	0.1	0.1	0.1
18 to 25	0.3	0.3	0.4	0.4	0.3
26 or Older	0.2	0.1 <sup>b</sup>	0.1 <sup>b</sup>	0.2	0.3
<b>GENDER</b>					
Male	0.3	0.1 <sup>b</sup>	0.2 <sup>a</sup>	0.3	0.4
Female	0.2	0.2	0.2	0.2	0.3

\*Low precision, no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.25A Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2010**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>LIFETIME USE</b>					
<b>TOTAL</b>	7,757	7,065 <sup>b</sup>	7,521	7,386	7,999
<b>AGE GROUP</b>					
12 to 17	930 <sup>b</sup>	856	795	749	755
18 to 25	2,133	2,034 <sup>a</sup>	2,175	2,305	2,306
26 or Older	4,694	4,175 <sup>a</sup>	4,550	4,332	4,938
<b>GENDER</b>					
Male	4,871	4,192 <sup>b</sup>	4,426	4,255 <sup>a</sup>	4,998
Female	2,885	2,873	3,095	3,131	3,001
<b>PAST YEAR USE</b>					
<b>TOTAL</b>	1,585	1,350 <sup>a</sup>	1,435	1,467	1,772
<b>AGE GROUP</b>					
12 to 17	485 <sup>b</sup>	435	405	371	371
18 to 25	511	479	463	481	484
26 or Older	589	436 <sup>b</sup>	567	614	917
<b>GENDER</b>					
Male	889	686 <sup>b</sup>	782	726 <sup>a</sup>	1,097
Female	696	664	653	740	675
<b>PAST MONTH USE</b>					
<b>TOTAL</b>	423	402	402	412	464
<b>AGE GROUP</b>					
12 to 17	122	125	105	108	93
18 to 25	107	130	105	124	126
26 or Older	193	147	193	181	246
<b>GENDER</b>					
Male	263	205	218	195	246
Female	159	197	183	217	218

\*Low precision, no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.25B Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2010**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>LIFETIME USE</b>					
<b>TOTAL</b>	3.2	2.9 <sup>a</sup>	3.0	2.9	3.2
<b>AGE GROUP</b>					
12 to 17	3.7 <sup>a</sup>	3.4	3.2	3.0	3.1
18 to 25	6.5	6.2	6.6	6.9	6.8
26 or Older	2.5	2.2	2.4	2.2	2.5
<b>GENDER</b>					
Male	4.1	3.5 <sup>a</sup>	3.7	3.5 <sup>a</sup>	4.0
Female	2.3	2.3	2.4	2.4	2.3
<b>PAST YEAR USE</b>					
<b>TOTAL</b>	0.6	0.5 <sup>a</sup>	0.6	0.6	0.7
<b>AGE GROUP</b>					
12 to 17	1.9 <sup>a</sup>	1.7	1.6	1.5	1.5
18 to 25	1.6	1.5	1.4	1.4	1.4
26 or Older	0.3	0.2 <sup>b</sup>	0.3	0.3	0.5
<b>GENDER</b>					
Male	0.7	0.6 <sup>a</sup>	0.6	0.6 <sup>a</sup>	0.9
Female	0.5	0.5	0.5	0.6	0.5
<b>PAST MONTH USE</b>					
<b>TOTAL</b>	0.2	0.2	0.2	0.2	0.2
<b>AGE GROUP</b>					
12 to 17	0.5	0.5	0.4	0.4	0.4
18 to 25	0.3	0.4	0.3	0.4	0.4
26 or Older	0.1	0.1	0.1	0.1	0.1
<b>GENDER</b>					
Male	0.2	0.2	0.2	0.2	0.2
Female	0.1	0.2	0.1	0.2	0.2

\*Low precision, no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.26A Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2010**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>LIFETIME USE</b>					
<b>TOTAL</b>	2,277	2,178	2,125	2,313	2,399
<b>AGE GROUP</b>					
12 to 17	83	70	73	78	80
18 to 25	914 <sup>b</sup>	793 <sup>b</sup>	626	586	549
26 or Older	1,279 <sup>a</sup>	1,315 <sup>a</sup>	1,426	1,650	1,771
<b>GENDER</b>					
Male	1,467	1,427	1,336 <sup>a</sup>	1,550	1,740
Female	810	751	788	763	659
<b>PAST YEAR USE</b>					
<b>TOTAL</b>	203	211	148 <sup>a</sup>	202	245
<b>AGE GROUP</b>					
12 to 17	33	32	34	35	51
18 to 25	74 <sup>a</sup>	121	71 <sup>a</sup>	92	135
26 or Older	95	58	43	75	59
<b>GENDER</b>					
Male	141	143	103	115	175
Female	62	68	46	88	70
<b>PAST MONTH USE</b>					
<b>TOTAL</b>	53	39	27	86	61
<b>AGE GROUP</b>					
12 to 17	11	12	10	17	14
18 to 25	14	17	16	26	22
26 or Older	28	10	*	43	24
<b>GENDER</b>					
Male	34	27	16	42	39
Female	19	12	10	43	21

\*Low precision, no estimate reported.

NOTE: Ketamine is also known as "Special K" or "Super K."

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.26B Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2010**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>LIFETIME USE</b>					
<b>TOTAL</b>	0.9	0.9	0.9	0.9	0.9
<b>AGE GROUP</b>					
12 to 17	0.3	0.3	0.3	0.3	0.3
18 to 25	2.8 <sup>b</sup>	2.4 <sup>b</sup>	1.9	1.7	1.6
26 or Older	0.7 <sup>a</sup>	0.7 <sup>a</sup>	0.7	0.9	0.9
<b>GENDER</b>					
Male	1.2	1.2	1.1 <sup>a</sup>	1.3	1.4
Female	0.6	0.6	0.6	0.6	0.5
<b>PAST YEAR USE</b>					
<b>TOTAL</b>	0.1	0.1	0.1 <sup>a</sup>	0.1	0.1
<b>AGE GROUP</b>					
12 to 17	0.1	0.1	0.1	0.1	0.2
18 to 25	0.2 <sup>a</sup>	0.4	0.2 <sup>a</sup>	0.3	0.4
26 or Older	0.1	0.0	0.0	0.0	0.0
<b>GENDER</b>					
Male	0.1	0.1	0.1	0.1	0.1
Female	0.0	0.1	0.0	0.1	0.1
<b>PAST MONTH USE</b>					
<b>TOTAL</b>	0.0	0.0	0.0	0.0	0.0
<b>AGE GROUP</b>					
12 to 17	0.0	0.0	0.0	0.1	0.1
18 to 25	0.0	0.1	0.0	0.1	0.1
26 or Older	0.0	0.0	*	0.0	0.0
<b>GENDER</b>					
Male	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0

\*Low precision, no estimate reported.

NOTE: Ketamine is also known as "Special K" or "Super K."

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.27A DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2010**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>LIFETIME USE</b>					
<b>TOTAL</b>	688 <sup>b</sup>	675 <sup>b</sup>	729 <sup>b</sup>	874 <sup>b</sup>	1,208
<b>AGE GROUP</b>					
12 to 17	45	44	49	58	61
18 to 25	308 <sup>b</sup>	269 <sup>b</sup>	278 <sup>b</sup>	360	440
26 or Older	335 <sup>b</sup>	362 <sup>b</sup>	402 <sup>b</sup>	455 <sup>a</sup>	707
<b>GENDER</b>					
Male	508 <sup>b</sup>	517 <sup>b</sup>	472 <sup>b</sup>	637 <sup>a</sup>	879
Female	180 <sup>a</sup>	158 <sup>b</sup>	257	237	328
<b>PAST YEAR USE</b>					
<b>TOTAL</b>	104 <sup>b</sup>	79 <sup>b</sup>	150	192	233
<b>AGE GROUP</b>					
12 to 17	16 <sup>a</sup>	19	29	38	38
18 to 25	65 <sup>a</sup>	52 <sup>b</sup>	73 <sup>a</sup>	109	116
26 or Older	24	*	47	45	79
<b>GENDER</b>					
Male	84 <sup>a</sup>	61 <sup>b</sup>	111	115	160
Female	20 <sup>a</sup>	18 <sup>a</sup>	39	76	73
<b>PAST MONTH USE</b>					
<b>TOTAL</b>	20 <sup>a</sup>	*	46	57	81
<b>AGE GROUP</b>					
12 to 17	*	*	8	11	17
18 to 25	15	9	19	12	21
26 or Older	*	*	19	34	43
<b>GENDER</b>					
Male	13	7	40	22	50
Female	7	*	*	35	31

\*Low precision, no estimate reported.

NOTE: DMT is also known as dimethyltryptamine.

NOTE: AMT is also known as alpha-methyltryptamine.

NOTE: 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy".

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.27B DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2010**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>LIFETIME USE</b>					
<b>TOTAL</b>	0.3 <sup>b</sup>	0.3 <sup>b</sup>	0.3 <sup>b</sup>	0.3 <sup>b</sup>	0.5
<b>AGE GROUP</b>					
12 to 17	0.2	0.2	0.2	0.2	0.3
18 to 25	0.9 <sup>a</sup>	0.8 <sup>b</sup>	0.8 <sup>b</sup>	1.1	1.3
26 or Older	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.2 <sup>b</sup>	0.2 <sup>a</sup>	0.4
<b>GENDER</b>					
Male	0.4 <sup>b</sup>	0.4 <sup>b</sup>	0.4 <sup>b</sup>	0.5 <sup>a</sup>	0.7
Female	0.1 <sup>a</sup>	0.1 <sup>b</sup>	0.2	0.2	0.3
<b>PAST YEAR USE</b>					
<b>TOTAL</b>	0.0 <sup>b</sup>	0.0 <sup>b</sup>	0.1	0.1	0.1
<b>AGE GROUP</b>					
12 to 17	0.1 <sup>a</sup>	0.1	0.1	0.2	0.2
18 to 25	0.2 <sup>a</sup>	0.2 <sup>b</sup>	0.2	0.3	0.3
26 or Older	0.0	*	0.0	0.0	0.0
<b>GENDER</b>					
Male	0.1 <sup>a</sup>	0.1 <sup>b</sup>	0.1	0.1	0.1
Female	0.0 <sup>a</sup>	0.0 <sup>a</sup>	0.0	0.1	0.1
<b>PAST MONTH USE</b>					
<b>TOTAL</b>	0.0 <sup>a</sup>	*	0.0	0.0	0.0
<b>AGE GROUP</b>					
12 to 17	*	*	0.0	0.0	0.1
18 to 25	0.0	0.0	0.1	0.0	0.1
26 or Older	*	*	0.0	0.0	0.0
<b>GENDER</b>					
Male	0.0	0.0	0.0	0.0	0.0
Female	0.0	*	*	0.0	0.0

\*Low precision, no estimate reported.

NOTE: DMT is also known as dimethyltryptamine.

NOTE: AMT is also known as alpha-methyltryptamine.

NOTE: 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy".

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.28A Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2010**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>LIFETIME USE</b>					
<b>TOTAL</b>	1,802 <sup>b</sup>	2,513 <sup>b</sup>	3,170 <sup>b</sup>	4,370 <sup>b</sup>	5,204
<b>AGE GROUP</b>					
12 to 17	228 <sup>b</sup>	297 <sup>b</sup>	393	414	398
18 to 25	1,185 <sup>b</sup>	1,597 <sup>b</sup>	1,941 <sup>b</sup>	2,673 <sup>a</sup>	2,973
26 or Older	389 <sup>b</sup>	619 <sup>b</sup>	836 <sup>b</sup>	1,283 <sup>b</sup>	1,833
<b>GENDER</b>					
Male	1,436 <sup>b</sup>	1,867 <sup>b</sup>	2,276 <sup>b</sup>	3,352 <sup>a</sup>	3,884
Female	366 <sup>b</sup>	646 <sup>b</sup>	894 <sup>b</sup>	1,018 <sup>b</sup>	1,320
<b>PAST YEAR USE</b>					
<b>TOTAL</b>	756 <sup>b</sup>	1,004	1,207	1,278	1,183
<b>AGE GROUP</b>					
12 to 17	148 <sup>b</sup>	191	263	242	229
18 to 25	548	655	736	777 <sup>a</sup>	645
26 or Older	60 <sup>b</sup>	158 <sup>a</sup>	207	259	309
<b>GENDER</b>					
Male	596 <sup>b</sup>	735 <sup>a</sup>	874	955	923
Female	160 <sup>b</sup>	269	333	323	260
<b>PAST MONTH USE</b>					
<b>TOTAL</b>	133	200	200	193	161
<b>AGE GROUP</b>					
12 to 17	28	42	59	49	44
18 to 25	102	134	131	95	93
26 or Older	*	24	10	49	25
<b>GENDER</b>					
Male	110	152	162	132	133
Female	23	49	38	60	28

\*Low precision, no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.

**Table 7.28B Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2010**

<b>Drug Use/Demographic Characteristic</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>	<b>2010</b>
<b>LIFETIME USE</b>					
<b>TOTAL</b>	0.7 <sup>b</sup>	1.0 <sup>b</sup>	1.3 <sup>b</sup>	1.7 <sup>b</sup>	2.1
<b>AGE GROUP</b>					
12 to 17	0.9 <sup>b</sup>	1.2 <sup>b</sup>	1.6	1.7	1.6
18 to 25	3.6 <sup>b</sup>	4.9 <sup>b</sup>	5.9 <sup>b</sup>	8.0 <sup>a</sup>	8.7
26 or Older	0.2 <sup>b</sup>	0.3 <sup>b</sup>	0.4 <sup>b</sup>	0.7 <sup>b</sup>	0.9
<b>GENDER</b>					
Male	1.2 <sup>b</sup>	1.6 <sup>b</sup>	1.9 <sup>b</sup>	2.7 <sup>a</sup>	3.1
Female	0.3 <sup>b</sup>	0.5 <sup>b</sup>	0.7 <sup>b</sup>	0.8 <sup>b</sup>	1.0
<b>PAST YEAR USE</b>					
<b>TOTAL</b>	0.3 <sup>b</sup>	0.4	0.5	0.5	0.5
<b>AGE GROUP</b>					
12 to 17	0.6 <sup>b</sup>	0.8	1.1	1.0	0.9
18 to 25	1.7	2.0	2.2	2.3 <sup>a</sup>	1.9
26 or Older	0.0 <sup>b</sup>	0.1 <sup>a</sup>	0.1	0.1	0.2
<b>GENDER</b>					
Male	0.5 <sup>b</sup>	0.6	0.7	0.8	0.7
Female	0.1 <sup>b</sup>	0.2	0.3	0.2	0.2
<b>PAST MONTH USE</b>					
<b>TOTAL</b>	0.1	0.1	0.1	0.1	0.1
<b>AGE GROUP</b>					
12 to 17	0.1	0.2	0.2	0.2	0.2
18 to 25	0.3	0.4	0.4	0.3	0.3
26 or Older	*	0.0	0.0	0.0	0.0
<b>GENDER</b>					
Male	0.1	0.1	0.1	0.1	0.1
Female	0.0	0.0	0.0	0.0	0.0

\*Low precision, no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2006-2010.