

Table 7.40C Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002-2010

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010
ILLICIT DRUGS^{1,2}	224	206	210	203	221	231	210	225	244
Marijuana and Hashish	158	140	158	143	154	155	156	161	184
Cocaine	119	115	114	108	127	125	101	105	100
Heroin	45	37	49	36	66	43	46	72	55
Hallucinogens	38	34	44	36	41	41	38	41	43
Inhalants	31	27	28	38	27	22	28	20	26
Nonmedical Use of Psychotherapeutics ^{2,3}	139	116	115	108	115	128	113	130	151
Pain Relievers	126	102	86	97	104	116	100	114	139
Tranquilizers	76	56	73	51	53	65	50	63	85
Stimulants ²	68	41	48	44	46	63	44	42	50
Sedatives	57	37	27	20	43	36	27	34	39
ALCOHOL	415	387	417	400	412	409	395	418	411
BOTH ILLICIT DRUGS AND ALCOHOL^{1,2}	145	138	143	141	141	145	128	158	130
ILLICIT DRUGS OR ALCOHOL^{1,2}	439	410	435	426	455	456	423	455	466

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

Table 7.40D Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 or Older: Standard Errors of Percentages, 2002-2010

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010
ILLICIT DRUGS^{1,2}	0.10	0.09	0.09	0.08	0.09	0.09	0.08	0.09	0.10
Marijuana and Hashish	0.07	0.06	0.07	0.06	0.06	0.06	0.06	0.06	0.07
Cocaine	0.05	0.05	0.05	0.04	0.05	0.05	0.04	0.04	0.04
Heroin	0.02	0.02	0.02	0.01	0.03	0.02	0.02	0.03	0.02
Hallucinogens	0.02	0.01	0.02	0.01	0.02	0.02	0.02	0.02	0.02
Inhalants	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01	0.01
Nonmedical Use of Psychotherapeutics ^{2,3}	0.06	0.05	0.05	0.04	0.05	0.05	0.05	0.05	0.06
Pain Relievers	0.05	0.04	0.04	0.04	0.04	0.05	0.04	0.05	0.05
Tranquilizers	0.03	0.02	0.03	0.02	0.02	0.03	0.02	0.02	0.03
Stimulants ²	0.03	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.02
Sedatives	0.02	0.02	0.01	0.01	0.02	0.01	0.01	0.01	0.02
ALCOHOL	0.18	0.16	0.17	0.16	0.17	0.16	0.16	0.17	0.16
BOTH ILLICIT DRUGS AND ALCOHOL^{1,2}	0.06	0.06	0.06	0.06	0.06	0.06	0.05	0.06	0.05
ILLICIT DRUGS OR ALCOHOL^{1,2}	0.19	0.17	0.18	0.18	0.18	0.18	0.17	0.18	0.18

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.

Table 7.40P Substance Dependence or Abuse for Specific Substances in the Past Year among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2010 Versus 2002-2009

Past Year Dependence or Abuse	2002	2003	2004	2005	2006	2007	2008	2009	2010
ILLICIT DRUGS^{1,2}	0.0967	0.5632	0.0725	0.9459	0.6958	0.7920	0.9835	0.8820	
Marijuana and Hashish	0.5344	0.9874	0.3387	0.3701	0.4710	0.0627	0.3694	0.5521	
Cocaine	0.0002	0.0001	0.0000	0.0001	0.0000	0.0001	0.0024	0.3903	
Heroin	0.0816	0.0205	0.3272	0.0504	0.7732	0.0488	0.3164	0.6359	
Hallucinogens	0.2932	0.3349	0.2235	0.8681	0.9360	0.7369	0.5686	0.6983	
Inhalants	0.4382	0.6170	0.0310	0.1433	0.6024	0.8571	0.6607	0.9107	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.3498	0.0978	0.2647	0.0822	0.1568	0.4159	0.3794	0.6994	
Pain Relievers	0.1319	0.0226	0.0059	0.0721	0.1915	0.3439	0.2946	0.7648	
Tranquilizers	0.8109	0.5841	0.4673	0.4092	0.3008	0.5341	0.5288	0.7306	
Stimulants ²	0.2033	0.4837	0.0537	0.3059	0.5147	0.4721	0.9941	0.7986	
Sedatives	0.9588	0.9079	0.5688	0.1596	0.5161	0.9345	0.4712	0.7824	
ALCOHOL	0.0069	0.0543	0.0030	0.0065	0.0105	0.0402	0.2053	0.1210	
BOTH ILLICIT DRUGS AND ALCOHOL^{1,2}	0.0035	0.0467	0.0001	0.0059	0.0246	0.0518	0.1386	0.0600	
ILLICIT DRUGS OR ALCOHOL^{1,2}	0.0148	0.1499	0.0134	0.0966	0.0666	0.2685	0.4705	0.3733	

*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

² Estimates in these designated rows do not include data from new methamphetamine use items added in 2005 and 2006.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2002-2010.