

2010-2011
National Survey on Drug Use and Health:
Model-Based Estimated Totals
(in Thousands)
(50 States and the District of Columbia)

Revised Mental Illness Estimates: The Substance Abuse and Mental Health Services Administration (SAMHSA) has been publishing State estimates of the prevalence (both percentages and estimated counts) of past year serious mental illness (SMI) and any mental illness (AMI) among adults aged 18 or older since the release of the 2008-2009 State small area estimation (SAE) report. Estimates were based on a model developed in 2008. In 2013, SAMHSA developed a more accurate model for the 2012 data. This revised model incorporates the NSDUH respondent's age and indicators of past year suicide thoughts and depression, along with the variables that were specified in the 2008 model (Kessler-6 [K6] questions on psychological distress and an abbreviated set of questions on impairment in carrying out activities from the World Health Organization Disability Assessment Schedule [WHODAS]), leading to more accurate estimates of SMI and AMI. The SMI and AMI tables in this document include estimates that have been revised based on the new model. Additionally, note that the 2010-2011 small area estimates for these two measures were produced using an updated 2010 sample weight (i.e., one that used 2010 population counts based on the 2010 census rather than the 2000 census in the poststratification adjustment) and updated SAE model predictors for both 2010 and 2011. Other mental health measures in these tables, such as major depressive episode (MDE) and serious thoughts of suicide, were not affected. For further information on the revised model, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>. For the further details on the revised weight and predictors used for these 2010-2011 SMI and AMI small area estimates, see the "2010-2011 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at <http://www.samhsa.gov/data/NSDUH/2k11State/NSDUHsae2011/Index.aspx>.

Table 1 *Illicit Drug Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs*

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	22,553	(21,889 - 23,231)	2,495	(2,399 - 2,594)	7,348	(7,148 - 7,553)	12,709	(12,119 - 13,323)	20,057	(19,406 - 20,726)
Northeast	4,334	(4,105 - 4,579)	446	(417 - 476)	1,433	(1,357 - 1,512)	2,455	(2,247 - 2,682)	3,888	(3,666 - 4,128)
Midwest	4,672	(4,465 - 4,883)	514	(488 - 542)	1,548	(1,474 - 1,624)	2,610	(2,421 - 2,810)	4,158	(3,957 - 4,364)
South	7,246	(6,934 - 7,570)	837	(792 - 885)	2,405	(2,281 - 2,534)	4,004	(3,716 - 4,313)	6,408	(6,107 - 6,725)
West	6,301	(5,965 - 6,650)	698	(652 - 747)	1,963	(1,855 - 2,075)	3,640	(3,338 - 3,966)	5,603	(5,276 - 5,944)
Alabama	299	(250 - 357)	35	(29 - 42)	91	(76 - 108)	173	(133 - 223)	264	(217 - 320)
Alaska	77	(65 - 90)	7	(5 - 8)	21	(18 - 25)	49	(39 - 61)	70	(59 - 83)
Arizona	509	(423 - 608)	61	(50 - 74)	140	(118 - 166)	307	(238 - 395)	448	(367 - 543)
Arkansas	182	(151 - 219)	22	(18 - 26)	56	(47 - 66)	105	(80 - 137)	160	(131 - 196)
California	3,213	(2,950 - 3,497)	380	(344 - 420)	1,044	(962 - 1,132)	1,789	(1,560 - 2,048)	2,833	(2,578 - 3,110)
Colorado	559	(480 - 648)	51	(43 - 61)	171	(150 - 192)	337	(270 - 418)	507	(432 - 594)
Connecticut	271	(229 - 321)	30	(25 - 36)	91	(78 - 106)	150	(116 - 194)	242	(201 - 290)
Delaware	67	(57 - 80)	8	(6 - 9)	24	(20 - 27)	36	(27 - 47)	59	(49 - 71)
District of Columbia	71	(61 - 83)	5	(4 - 6)	23	(20 - 27)	43	(34 - 55)	67	(56 - 78)
Florida	1,320	(1,201 - 1,452)	123	(110 - 138)	401	(366 - 438)	796	(687 - 920)	1,197	(1,080 - 1,327)
Georgia	517	(429 - 622)	72	(59 - 86)	181	(151 - 216)	264	(198 - 350)	445	(362 - 545)
Hawaii	95	(79 - 114)	10	(8 - 12)	27	(23 - 32)	58	(45 - 75)	85	(69 - 103)
Idaho	118	(99 - 140)	14	(11 - 17)	34	(28 - 40)	71	(55 - 90)	104	(86 - 126)
Illinois	910	(831 - 996)	102	(91 - 115)	318	(292 - 345)	490	(422 - 568)	808	(731 - 891)
Indiana	479	(402 - 567)	49	(40 - 59)	159	(136 - 184)	271	(210 - 349)	430	(357 - 516)
Iowa	161	(135 - 192)	20	(17 - 25)	62	(52 - 73)	79	(60 - 105)	141	(116 - 170)
Kansas	155	(128 - 188)	19	(16 - 24)	56	(46 - 67)	80	(60 - 107)	136	(110 - 167)
Kentucky	249	(206 - 299)	29	(24 - 35)	82	(68 - 97)	139	(105 - 181)	220	(180 - 269)
Louisiana	251	(212 - 297)	30	(25 - 36)	89	(76 - 104)	132	(101 - 172)	221	(184 - 265)
Maine	109	(92 - 128)	10	(8 - 12)	34	(30 - 39)	64	(51 - 82)	99	(83 - 118)
Maryland	355	(296 - 423)	40	(33 - 49)	125	(106 - 147)	189	(144 - 246)	314	(259 - 380)
Massachusetts	647	(558 - 748)	64	(54 - 75)	231	(203 - 260)	352	(278 - 444)	583	(497 - 682)
Michigan	889	(815 - 970)	89	(81 - 99)	256	(237 - 276)	543	(475 - 620)	800	(727 - 879)
Minnesota	362	(304 - 429)	39	(32 - 47)	124	(106 - 144)	199	(154 - 257)	323	(268 - 388)
Mississippi	195	(165 - 230)	24	(20 - 28)	63	(53 - 74)	108	(84 - 139)	171	(142 - 205)
Missouri	362	(304 - 431)	44	(36 - 52)	135	(115 - 157)	184	(139 - 242)	319	(263 - 385)
Montana	100	(86 - 115)	9	(8 - 11)	30	(27 - 34)	60	(49 - 74)	90	(78 - 105)

(continued)

Table 1 Illicit Drug Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	99	(82 - 120)	11	(9 - 14)	34	(29 - 41)	53	(40 - 71)	88	(71 - 108)
Nevada	218	(181 - 262)	25	(20 - 30)	62	(52 - 72)	131	(100 - 171)	193	(158 - 236)
New Hampshire	124	(105 - 145)	13	(11 - 16)	44	(39 - 49)	67	(52 - 86)	111	(93 - 131)
New Jersey	584	(491 - 694)	70	(58 - 85)	190	(161 - 223)	323	(250 - 416)	514	(426 - 619)
New Mexico	177	(150 - 208)	20	(17 - 24)	56	(49 - 65)	100	(78 - 128)	157	(131 - 187)
New York	1,525	(1,383 - 1,680)	152	(135 - 171)	488	(447 - 532)	885	(761 - 1,027)	1,373	(1,236 - 1,524)
North Carolina	692	(586 - 812)	76	(64 - 90)	244	(211 - 281)	371	(288 - 477)	616	(515 - 732)
North Dakota	31	(25 - 37)	3	(3 - 4)	12	(10 - 15)	15	(11 - 21)	27	(22 - 34)
Ohio	815	(743 - 895)	90	(81 - 101)	259	(239 - 281)	466	(401 - 540)	725	(655 - 803)
Oklahoma	274	(230 - 325)	30	(25 - 36)	82	(69 - 97)	162	(126 - 207)	244	(202 - 293)
Oregon	425	(363 - 496)	37	(31 - 43)	117	(102 - 132)	271	(218 - 336)	388	(328 - 458)
Pennsylvania	862	(783 - 949)	89	(79 - 101)	284	(259 - 312)	489	(420 - 568)	773	(696 - 859)
Rhode Island	130	(112 - 150)	11	(9 - 12)	45	(40 - 50)	75	(59 - 94)	120	(102 - 139)
South Carolina	282	(236 - 337)	31	(26 - 38)	90	(76 - 106)	161	(124 - 208)	251	(207 - 304)
South Dakota	40	(32 - 49)	5	(4 - 6)	13	(11 - 16)	21	(16 - 29)	35	(27 - 44)
Tennessee	402	(334 - 482)	44	(37 - 53)	126	(106 - 148)	232	(177 - 303)	358	(293 - 435)
Texas	1,442	(1,309 - 1,586)	198	(176 - 223)	501	(453 - 552)	743	(632 - 871)	1,243	(1,116 - 1,382)
Utah	108	(88 - 133)	15	(12 - 20)	39	(31 - 50)	54	(40 - 73)	93	(74 - 116)
Vermont	83	(72 - 94)	7	(6 - 8)	26	(23 - 29)	50	(40 - 60)	76	(65 - 87)
Virginia	523	(441 - 620)	57	(48 - 69)	185	(158 - 217)	281	(216 - 363)	466	(387 - 560)
Washington	670	(576 - 778)	65	(55 - 77)	211	(185 - 239)	394	(315 - 492)	605	(514 - 710)
West Virginia	125	(105 - 150)	13	(11 - 16)	42	(36 - 48)	70	(53 - 92)	112	(92 - 136)
Wisconsin	369	(306 - 444)	42	(34 - 50)	120	(101 - 141)	208	(158 - 272)	327	(267 - 400)
Wyoming	33	(27 - 39)	4	(3 - 4)	10	(9 - 13)	19	(14 - 24)	29	(24 - 35)

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

**Table 2 Marijuana Use in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages
Based on 2010 and 2011 NSDUHs**

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	29,523	(28,752 - 30,306)	3,484	(3,369 - 3,601)	10,385	(10,143 - 10,630)	15,654	(14,991 - 16,339)	26,039	(25,289 - 26,803)
Northeast	5,858	(5,609 - 6,122)	629	(594 - 665)	2,072	(1,993 - 2,151)	3,158	(2,935 - 3,397)	5,230	(4,986 - 5,488)
Midwest	6,110	(5,884 - 6,337)	701	(667 - 735)	2,229	(2,151 - 2,309)	3,180	(2,980 - 3,389)	5,409	(5,191 - 5,630)
South	9,377	(9,032 - 9,735)	1,165	(1,113 - 1,219)	3,373	(3,247 - 3,503)	4,839	(4,527 - 5,171)	8,212	(7,876 - 8,562)
West	8,178	(7,794 - 8,574)	989	(933 - 1,048)	2,711	(2,595 - 2,829)	4,478	(4,129 - 4,850)	7,188	(6,813 - 7,577)
Alabama	355	(305 - 414)	43	(36 - 51)	120	(103 - 138)	192	(152 - 242)	312	(264 - 368)
Alaska	99	(87 - 113)	10	(9 - 12)	31	(28 - 35)	58	(47 - 71)	89	(77 - 102)
Arizona	619	(526 - 724)	85	(72 - 99)	195	(170 - 223)	339	(264 - 434)	535	(447 - 635)
Arkansas	222	(188 - 261)	27	(23 - 32)	75	(66 - 86)	119	(92 - 154)	194	(163 - 232)
California	4,304	(3,992 - 4,637)	530	(485 - 578)	1,435	(1,343 - 1,531)	2,339	(2,062 - 2,649)	3,774	(3,471 - 4,101)
Colorado	677	(595 - 768)	74	(63 - 86)	231	(209 - 254)	372	(303 - 454)	603	(525 - 690)
Connecticut	399	(347 - 459)	47	(40 - 55)	138	(123 - 154)	214	(171 - 268)	353	(302 - 410)
Delaware	99	(86 - 113)	11	(10 - 13)	36	(32 - 40)	52	(41 - 64)	88	(76 - 101)
District of Columbia	97	(86 - 110)	6	(5 - 7)	34	(30 - 37)	58	(48 - 69)	91	(80 - 104)
Florida	1,742	(1,609 - 1,888)	187	(171 - 205)	595	(557 - 635)	960	(840 - 1,095)	1,555	(1,425 - 1,699)
Georgia	752	(648 - 871)	103	(88 - 122)	267	(232 - 304)	382	(300 - 483)	648	(550 - 763)
Hawaii	129	(110 - 149)	14	(12 - 17)	39	(34 - 45)	75	(60 - 94)	114	(97 - 134)
Idaho	145	(124 - 169)	19	(16 - 22)	45	(40 - 52)	81	(63 - 103)	126	(106 - 150)
Illinois	1,213	(1,124 - 1,308)	138	(125 - 153)	451	(422 - 481)	623	(547 - 710)	1,074	(989 - 1,167)
Indiana	548	(471 - 637)	60	(51 - 71)	207	(182 - 234)	281	(219 - 359)	488	(414 - 574)
Iowa	250	(216 - 288)	31	(26 - 36)	98	(86 - 111)	121	(95 - 154)	219	(187 - 256)
Kansas	221	(188 - 259)	30	(25 - 36)	88	(76 - 100)	103	(78 - 134)	190	(160 - 226)
Kentucky	315	(269 - 369)	41	(34 - 49)	109	(94 - 125)	166	(130 - 212)	275	(231 - 325)
Louisiana	317	(271 - 369)	40	(34 - 47)	116	(101 - 131)	161	(125 - 208)	277	(233 - 328)
Maine	160	(140 - 183)	15	(13 - 18)	49	(44 - 54)	96	(79 - 117)	145	(126 - 167)
Maryland	477	(409 - 556)	60	(50 - 71)	194	(171 - 219)	223	(170 - 293)	418	(353 - 494)
Massachusetts	891	(786 - 1,006)	90	(79 - 104)	323	(293 - 354)	477	(389 - 582)	800	(699 - 914)
Michigan	1,095	(1,018 - 1,178)	117	(106 - 128)	367	(346 - 388)	612	(541 - 690)	979	(903 - 1,060)
Minnesota	483	(416 - 556)	55	(47 - 65)	186	(164 - 209)	242	(190 - 307)	428	(365 - 499)
Mississippi	233	(201 - 270)	30	(25 - 35)	81	(71 - 93)	122	(97 - 154)	204	(173 - 239)
Missouri	500	(430 - 581)	62	(53 - 73)	181	(159 - 205)	257	(201 - 328)	438	(371 - 516)
Montana	135	(120 - 152)	15	(13 - 17)	43	(39 - 47)	78	(65 - 93)	121	(106 - 137)

(continued)

Table 2 *Marijuana Use in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)*

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Nebraska	137	(117 - 160)	16	(13 - 19)	53	(46 - 61)	68	(52 - 88)	121	(102 - 143)
Nevada	280	(240 - 327)	37	(31 - 44)	87	(75 - 99)	157	(123 - 198)	243	(205 - 288)
New Hampshire	159	(139 - 181)	19	(16 - 22)	59	(54 - 64)	81	(65 - 102)	140	(121 - 161)
New Jersey	786	(682 - 905)	101	(86 - 118)	276	(245 - 310)	409	(324 - 514)	685	(586 - 799)
New Mexico	222	(193 - 254)	31	(27 - 36)	77	(68 - 86)	114	(90 - 143)	191	(163 - 222)
New York	2,047	(1,890 - 2,215)	208	(188 - 229)	701	(658 - 746)	1,137	(999 - 1,293)	1,839	(1,686 - 2,004)
North Carolina	871	(753 - 1,001)	106	(90 - 123)	332	(294 - 372)	433	(340 - 550)	765	(654 - 891)
North Dakota	49	(43 - 57)	5	(4 - 6)	23	(20 - 26)	21	(16 - 28)	44	(38 - 51)
Ohio	1,103	(1,021 - 1,192)	125	(113 - 137)	383	(360 - 406)	595	(522 - 678)	978	(898 - 1,066)
Oklahoma	308	(263 - 360)	40	(34 - 48)	105	(90 - 121)	163	(127 - 208)	267	(225 - 317)
Oregon	519	(456 - 590)	54	(47 - 62)	161	(145 - 178)	305	(250 - 369)	466	(404 - 535)
Pennsylvania	1,156	(1,062 - 1,258)	124	(112 - 138)	432	(402 - 463)	599	(519 - 691)	1,031	(940 - 1,131)
Rhode Island	160	(141 - 180)	15	(12 - 17)	59	(53 - 64)	87	(71 - 105)	145	(127 - 165)
South Carolina	406	(349 - 470)	48	(41 - 56)	138	(122 - 155)	220	(174 - 277)	358	(304 - 419)
South Dakota	56	(47 - 66)	7	(6 - 9)	22	(19 - 26)	26	(20 - 35)	48	(40 - 58)
Tennessee	495	(419 - 582)	57	(48 - 68)	166	(144 - 190)	272	(211 - 348)	438	(365 - 522)
Texas	1,872	(1,723 - 2,033)	270	(244 - 299)	697	(646 - 749)	905	(779 - 1,051)	1,602	(1,458 - 1,758)
Utah	154	(130 - 182)	25	(20 - 31)	68	(57 - 81)	61	(45 - 83)	129	(107 - 156)
Vermont	101	(90 - 114)	9	(8 - 11)	35	(32 - 38)	57	(47 - 69)	92	(81 - 104)
Virginia	654	(564 - 758)	78	(66 - 92)	255	(225 - 288)	321	(251 - 410)	576	(490 - 676)
Washington	850	(746 - 965)	91	(78 - 106)	281	(254 - 309)	478	(389 - 583)	759	(659 - 870)
West Virginia	162	(139 - 187)	18	(15 - 21)	55	(48 - 61)	89	(70 - 113)	144	(122 - 169)
Wisconsin	454	(388 - 529)	54	(46 - 64)	170	(148 - 194)	230	(178 - 295)	400	(337 - 473)
Wyoming	44	(38 - 51)	5	(5 - 6)	17	(15 - 19)	22	(17 - 28)	39	(33 - 45)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

**Table 3 Marijuana Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages
Based on 2010 and 2011 NSDUHs**

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	17,741	(17,163 - 18,334)	1,885	(1,799 - 1,974)	6,419	(6,228 - 6,614)	9,437	(8,931 - 9,968)	15,856	(15,290 - 16,439)
Northeast	3,478	(3,286 - 3,683)	362	(336 - 389)	1,297	(1,227 - 1,371)	1,819	(1,650 - 2,004)	3,116	(2,931 - 3,316)
Midwest	3,674	(3,498 - 3,855)	383	(359 - 409)	1,346	(1,277 - 1,416)	1,946	(1,793 - 2,110)	3,291	(3,121 - 3,467)
South	5,480	(5,204 - 5,771)	604	(564 - 645)	2,053	(1,942 - 2,169)	2,823	(2,577 - 3,093)	4,876	(4,609 - 5,160)
West	5,109	(4,813 - 5,419)	537	(494 - 583)	1,723	(1,624 - 1,827)	2,849	(2,584 - 3,139)	4,572	(4,284 - 4,875)
Alabama	205	(168 - 249)	24	(19 - 30)	73	(61 - 88)	107	(80 - 144)	181	(146 - 223)
Alaska	64	(54 - 76)	5	(4 - 7)	20	(17 - 23)	39	(31 - 50)	59	(49 - 70)
Arizona	375	(304 - 461)	44	(35 - 56)	120	(99 - 143)	211	(156 - 285)	331	(264 - 412)
Arkansas	122	(98 - 151)	14	(11 - 18)	43	(36 - 52)	65	(47 - 89)	108	(85 - 136)
California	2,642	(2,408 - 2,895)	295	(261 - 334)	921	(843 - 1,004)	1,425	(1,223 - 1,658)	2,346	(2,121 - 2,593)
Colorado	458	(389 - 537)	42	(34 - 51)	153	(133 - 174)	264	(207 - 334)	416	(351 - 493)
Connecticut	216	(180 - 260)	25	(20 - 31)	82	(70 - 96)	109	(82 - 146)	191	(157 - 233)
Delaware	52	(44 - 62)	6	(4 - 7)	21	(18 - 24)	26	(20 - 34)	47	(39 - 56)
District of Columbia	59	(50 - 69)	3	(3 - 4)	21	(18 - 25)	34	(26 - 44)	55	(46 - 65)
Florida	1,009	(904 - 1,127)	92	(81 - 106)	358	(326 - 394)	558	(468 - 664)	917	(815 - 1,032)
Georgia	394	(320 - 484)	53	(42 - 67)	153	(125 - 185)	188	(136 - 260)	341	(272 - 426)
Hawaii	75	(61 - 91)	8	(6 - 10)	23	(20 - 28)	44	(33 - 58)	67	(54 - 83)
Idaho	90	(74 - 109)	10	(8 - 13)	27	(23 - 33)	53	(40 - 69)	80	(65 - 98)
Illinois	731	(662 - 806)	80	(70 - 92)	279	(255 - 303)	372	(314 - 439)	650	(584 - 723)
Indiana	347	(287 - 420)	35	(27 - 44)	132	(112 - 155)	181	(135 - 242)	313	(255 - 382)
Iowa	131	(107 - 159)	14	(11 - 18)	55	(45 - 65)	62	(45 - 85)	116	(94 - 143)
Kansas	126	(102 - 155)	15	(11 - 18)	47	(39 - 57)	64	(46 - 87)	111	(89 - 139)
Kentucky	182	(149 - 223)	21	(17 - 27)	69	(58 - 83)	91	(67 - 124)	161	(130 - 199)
Louisiana	179	(146 - 218)	20	(16 - 25)	71	(59 - 85)	88	(64 - 120)	159	(128 - 196)
Maine	91	(77 - 108)	9	(7 - 11)	31	(27 - 35)	51	(39 - 67)	82	(69 - 99)
Maryland	266	(218 - 324)	28	(21 - 36)	110	(92 - 131)	128	(94 - 174)	238	(193 - 293)
Massachusetts	558	(477 - 651)	56	(46 - 67)	217	(190 - 245)	285	(220 - 368)	502	(425 - 592)
Michigan	720	(655 - 791)	67	(59 - 76)	230	(212 - 249)	423	(365 - 489)	653	(590 - 722)
Minnesota	284	(236 - 340)	29	(23 - 36)	109	(92 - 127)	147	(111 - 194)	256	(210 - 309)
Mississippi	141	(117 - 171)	16	(12 - 20)	51	(42 - 61)	75	(56 - 100)	126	(102 - 154)
Missouri	300	(247 - 362)	35	(28 - 43)	120	(101 - 141)	145	(106 - 198)	265	(216 - 325)
Montana	85	(73 - 99)	8	(6 - 9)	27	(24 - 31)	51	(40 - 63)	78	(66 - 91)

(continued)

Table 3 Marijuana Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	83	(67 - 101)	9	(7 - 11)	29	(24 - 35)	45	(33 - 60)	74	(59 - 92)
Nevada	168	(135 - 208)	18	(14 - 23)	54	(45 - 63)	96	(71 - 130)	150	(119 - 187)
New Hampshire	98	(83 - 115)	12	(9 - 14)	38	(34 - 43)	48	(36 - 64)	86	(72 - 103)
New Jersey	469	(393 - 559)	56	(45 - 70)	170	(143 - 200)	243	(185 - 318)	412	(341 - 499)
New Mexico	138	(116 - 164)	15	(12 - 19)	48	(41 - 56)	75	(57 - 98)	123	(102 - 148)
New York	1,214	(1,103 - 1,337)	120	(106 - 137)	439	(403 - 478)	654	(558 - 766)	1,094	(986 - 1,213)
North Carolina	573	(476 - 687)	59	(48 - 73)	220	(188 - 256)	294	(221 - 391)	514	(421 - 624)
North Dakota	25	(20 - 31)	2	(2 - 3)	11	(9 - 13)	12	(9 - 17)	23	(18 - 29)
Ohio	620	(561 - 687)	66	(58 - 75)	223	(205 - 243)	331	(279 - 393)	555	(497 - 619)
Oklahoma	200	(162 - 245)	21	(17 - 27)	64	(53 - 78)	114	(85 - 153)	178	(143 - 222)
Oregon	356	(302 - 419)	30	(24 - 36)	104	(91 - 119)	222	(175 - 280)	326	(274 - 387)
Pennsylvania	654	(588 - 727)	69	(60 - 79)	253	(230 - 279)	331	(278 - 395)	585	(522 - 656)
Rhode Island	107	(92 - 124)	9	(7 - 11)	42	(37 - 47)	56	(44 - 71)	98	(83 - 115)
South Carolina	226	(186 - 273)	22	(18 - 28)	80	(67 - 95)	124	(93 - 163)	204	(166 - 248)
South Dakota	34	(27 - 42)	4	(3 - 5)	12	(10 - 15)	18	(13 - 25)	30	(23 - 38)
Tennessee	287	(234 - 350)	31	(24 - 39)	101	(83 - 122)	156	(116 - 208)	256	(206 - 317)
Texas	1,103	(987 - 1,231)	139	(120 - 160)	423	(381 - 469)	542	(447 - 655)	965	(854 - 1,088)
Utah	78	(60 - 99)	10	(7 - 13)	31	(24 - 40)	37	(26 - 54)	68	(52 - 88)
Vermont	71	(61 - 82)	6	(5 - 8)	25	(22 - 28)	40	(32 - 49)	65	(55 - 75)
Virginia	389	(320 - 471)	44	(35 - 54)	158	(132 - 187)	187	(138 - 253)	345	(281 - 423)
Washington	556	(475 - 650)	50	(41 - 61)	186	(163 - 211)	321	(253 - 405)	506	(428 - 597)
West Virginia	93	(77 - 113)	10	(8 - 12)	37	(32 - 43)	46	(34 - 63)	84	(68 - 102)
Wisconsin	274	(221 - 338)	28	(22 - 36)	99	(81 - 119)	146	(107 - 199)	245	(195 - 307)
Wyoming	23	(19 - 29)	2	(2 - 3)	9	(7 - 11)	12	(9 - 17)	21	(17 - 27)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 4 Perceptions of Great Risk of Smoking Marijuana Once a Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	82,620	(81,219 - 84,029)	7,048	(6,906 - 7,192)	6,253	(6,068 - 6,444)	69,318	(67,962 - 70,683)	75,572	(74,186 - 76,967)
Northeast	14,299	(13,827 - 14,774)	1,075	(1,028 - 1,123)	961	(906 - 1,018)	12,263	(11,809 - 12,723)	13,224	(12,759 - 13,694)
Midwest	17,264	(16,854 - 17,683)	1,580	(1,531 - 1,629)	1,263	(1,209 - 1,319)	14,422	(14,030 - 14,820)	15,684	(15,280 - 16,098)
South	33,939	(33,214 - 34,668)	2,881	(2,798 - 2,966)	2,634	(2,536 - 2,735)	28,424	(27,729 - 29,121)	31,058	(30,345 - 31,774)
West	17,118	(16,501 - 17,748)	1,512	(1,442 - 1,585)	1,396	(1,312 - 1,484)	14,209	(13,616 - 14,816)	15,605	(14,998 - 16,227)
Alabama	1,512	(1,391 - 1,637)	146	(132 - 160)	130	(114 - 148)	1,237	(1,122 - 1,355)	1,367	(1,247 - 1,490)
Alaska	125	(110 - 142)	15	(13 - 17)	10	(8 - 13)	100	(86 - 116)	111	(96 - 127)
Arizona	1,597	(1,432 - 1,774)	139	(120 - 161)	125	(105 - 147)	1,333	(1,178 - 1,497)	1,458	(1,297 - 1,631)
Arkansas	836	(762 - 913)	77	(68 - 86)	63	(54 - 72)	697	(627 - 768)	759	(687 - 834)
California	9,371	(8,868 - 9,888)	772	(715 - 831)	800	(732 - 873)	7,799	(7,316 - 8,298)	8,599	(8,104 - 9,109)
Colorado	876	(775 - 986)	82	(69 - 96)	61	(50 - 74)	733	(639 - 837)	794	(696 - 901)
Connecticut	868	(775 - 966)	60	(50 - 71)	50	(41 - 60)	758	(671 - 851)	808	(717 - 904)
Delaware	222	(201 - 245)	18	(16 - 21)	15	(12 - 17)	190	(170 - 211)	204	(184 - 226)
District of Columbia	134	(119 - 151)	7	(6 - 8)	12	(10 - 15)	115	(100 - 130)	127	(111 - 143)
Florida	6,127	(5,860 - 6,398)	415	(391 - 441)	387	(356 - 420)	5,325	(5,064 - 5,590)	5,712	(5,445 - 5,982)
Georgia	2,761	(2,521 - 3,010)	265	(235 - 297)	238	(205 - 275)	2,258	(2,035 - 2,490)	2,496	(2,263 - 2,739)
Hawaii	332	(300 - 367)	23	(20 - 27)	21	(18 - 25)	288	(257 - 321)	309	(277 - 343)
Idaho	388	(352 - 427)	40	(36 - 45)	33	(28 - 39)	315	(280 - 351)	348	(312 - 386)
Illinois	3,405	(3,235 - 3,578)	303	(283 - 323)	254	(233 - 276)	2,848	(2,685 - 3,016)	3,102	(2,935 - 3,273)
Indiana	1,709	(1,550 - 1,876)	167	(148 - 188)	131	(113 - 153)	1,410	(1,261 - 1,567)	1,542	(1,387 - 1,704)
Iowa	868	(792 - 949)	79	(71 - 88)	67	(57 - 78)	722	(650 - 797)	789	(714 - 868)
Kansas	729	(663 - 799)	68	(60 - 77)	57	(48 - 66)	605	(543 - 670)	661	(596 - 729)
Kentucky	1,347	(1,230 - 1,468)	130	(116 - 144)	114	(99 - 130)	1,103	(993 - 1,218)	1,217	(1,103 - 1,336)
Louisiana	1,512	(1,403 - 1,624)	129	(117 - 142)	132	(118 - 148)	1,251	(1,146 - 1,357)	1,383	(1,276 - 1,493)
Maine	273	(242 - 307)	23	(20 - 27)	16	(13 - 19)	234	(204 - 266)	250	(220 - 283)
Maryland	1,560	(1,412 - 1,715)	119	(103 - 135)	105	(89 - 123)	1,336	(1,197 - 1,482)	1,441	(1,296 - 1,593)
Massachusetts	1,356	(1,202 - 1,522)	101	(87 - 116)	94	(77 - 113)	1,161	(1,015 - 1,319)	1,255	(1,104 - 1,418)
Michigan	2,368	(2,243 - 2,497)	213	(199 - 228)	166	(151 - 182)	1,990	(1,869 - 2,114)	2,155	(2,032 - 2,282)
Minnesota	1,264	(1,145 - 1,390)	121	(107 - 137)	86	(73 - 102)	1,056	(945 - 1,175)	1,142	(1,027 - 1,266)
Mississippi	1,000	(928 - 1,074)	101	(91 - 111)	92	(81 - 104)	807	(740 - 875)	899	(829 - 971)
Missouri	1,624	(1,479 - 1,776)	130	(114 - 148)	125	(107 - 146)	1,368	(1,232 - 1,510)	1,493	(1,352 - 1,642)
Montana	201	(181 - 223)	17	(15 - 19)	14	(11 - 16)	171	(152 - 191)	184	(165 - 206)

(continued)

Table 4 Perceptions of Great Risk of Smoking Marijuana Once a Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	478	(435 - 525)	48	(43 - 54)	38	(33 - 44)	392	(351 - 435)	430	(387 - 476)
Nevada	618	(551 - 689)	54	(46 - 64)	42	(35 - 50)	522	(460 - 588)	564	(500 - 632)
New Hampshire	262	(231 - 296)	21	(18 - 25)	14	(11 - 17)	227	(197 - 259)	241	(211 - 274)
New Jersey	2,407	(2,183 - 2,641)	182	(158 - 207)	133	(111 - 158)	2,092	(1,879 - 2,315)	2,225	(2,006 - 2,455)
New Mexico	557	(507 - 610)	42	(36 - 48)	39	(33 - 45)	477	(429 - 527)	516	(466 - 567)
New York	5,347	(5,066 - 5,635)	393	(364 - 423)	395	(361 - 432)	4,559	(4,290 - 4,835)	4,954	(4,678 - 5,237)
North Carolina	2,467	(2,235 - 2,714)	201	(176 - 229)	179	(153 - 208)	2,087	(1,869 - 2,317)	2,266	(2,040 - 2,505)
North Dakota	189	(173 - 205)	17	(15 - 19)	19	(16 - 21)	153	(138 - 168)	171	(156 - 188)
Ohio	3,018	(2,866 - 3,174)	272	(255 - 290)	202	(186 - 218)	2,545	(2,397 - 2,696)	2,746	(2,596 - 2,900)
Oklahoma	1,071	(978 - 1,168)	95	(84 - 107)	86	(74 - 100)	890	(803 - 979)	976	(885 - 1,070)
Oregon	689	(606 - 780)	67	(58 - 76)	48	(40 - 58)	575	(496 - 661)	622	(541 - 712)
Pennsylvania	3,425	(3,249 - 3,604)	266	(247 - 286)	236	(215 - 260)	2,922	(2,754 - 3,096)	3,159	(2,986 - 3,336)
Rhode Island	233	(207 - 261)	17	(15 - 20)	15	(12 - 18)	201	(177 - 228)	216	(191 - 244)
South Carolina	1,422	(1,309 - 1,540)	106	(94 - 119)	103	(89 - 118)	1,214	(1,106 - 1,325)	1,317	(1,205 - 1,432)
South Dakota	222	(203 - 243)	21	(19 - 24)	18	(15 - 21)	183	(164 - 202)	201	(182 - 222)
Tennessee	1,810	(1,652 - 1,977)	153	(136 - 172)	129	(111 - 149)	1,527	(1,378 - 1,683)	1,656	(1,502 - 1,820)
Texas	7,525	(7,186 - 7,869)	720	(678 - 764)	646	(598 - 697)	6,158	(5,833 - 6,489)	6,805	(6,471 - 7,144)
Utah	858	(793 - 924)	115	(104 - 127)	102	(89 - 115)	641	(582 - 702)	742	(680 - 807)
Vermont	128	(114 - 144)	11	(10 - 13)	8	(7 - 10)	109	(95 - 123)	117	(103 - 132)
Virginia	2,054	(1,858 - 2,260)	156	(138 - 177)	165	(141 - 192)	1,732	(1,548 - 1,926)	1,897	(1,705 - 2,100)
Washington	1,357	(1,203 - 1,523)	131	(114 - 149)	90	(75 - 108)	1,135	(991 - 1,293)	1,226	(1,076 - 1,389)
West Virginia	578	(529 - 629)	43	(38 - 48)	39	(33 - 44)	497	(450 - 545)	535	(488 - 585)
Wisconsin	1,390	(1,251 - 1,537)	139	(122 - 156)	100	(85 - 117)	1,151	(1,021 - 1,290)	1,251	(1,116 - 1,396)
Wyoming	148	(134 - 162)	15	(14 - 17)	12	(10 - 14)	121	(108 - 134)	133	(119 - 147)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 5 Average Annual Marijuana Initiates, by Age Group and State: Estimated Numbers (in Thousands), Based on 2010 and 2011 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	2,829	(2,728 - 2,934)	1,316	(1,268 - 1,367)	1,309	(1,244 - 1,377)	204	(161 - 258)	1,513	(1,426 - 1,604)
Northeast	517	(486 - 551)	234	(219 - 250)	247	(227 - 268)	36	(25 - 52)	283	(258 - 311)
Midwest	594	(564 - 626)	276	(261 - 291)	279	(260 - 300)	39	(28 - 56)	319	(293 - 346)
South	968	(916 - 1,024)	449	(425 - 473)	451	(419 - 484)	69	(49 - 98)	520	(476 - 567)
West	749	(701 - 801)	358	(334 - 383)	332	(303 - 364)	59	(40 - 87)	391	(352 - 435)
Alabama	35	(30 - 42)	16	(13 - 19)	17	(13 - 21)	3	(1 - 5)	20	(15 - 25)
Alaska	7	(6 - 8)	4	(3 - 4)	3	(2 - 4)	1	(0 - 1)	4	(3 - 5)
Arizona	61	(52 - 72)	30	(24 - 36)	27	(21 - 34)	4	(2 - 8)	32	(25 - 39)
Arkansas	22	(18 - 26)	10	(8 - 12)	10	(8 - 13)	2	(1 - 3)	12	(9 - 15)
California	417	(378 - 461)	195	(176 - 216)	187	(163 - 215)	35	(21 - 60)	222	(191 - 260)
Colorado	53	(46 - 62)	28	(23 - 33)	22	(18 - 28)	3	(2 - 6)	26	(21 - 32)
Connecticut	33	(28 - 39)	16	(13 - 19)	15	(12 - 19)	2	(1 - 4)	17	(14 - 21)
Delaware	9	(7 - 10)	4	(3 - 5)	4	(3 - 5)	1	(0 - 1)	4	(4 - 6)
District of Columbia	7	(6 - 9)	2	(2 - 3)	5	(4 - 6)	1	(0 - 1)	5	(4 - 6)
Florida	152	(137 - 169)	67	(59 - 75)	73	(64 - 84)	13	(8 - 21)	86	(74 - 99)
Georgia	82	(70 - 98)	43	(35 - 52)	34	(26 - 44)	6	(3 - 11)	40	(31 - 51)
Hawaii	11	(9 - 14)	5	(4 - 6)	5	(4 - 7)	1	(1 - 2)	6	(5 - 8)
Idaho	14	(12 - 17)	7	(6 - 8)	6	(5 - 8)	1	(0 - 2)	7	(6 - 9)
Illinois	116	(105 - 128)	54	(48 - 60)	54	(47 - 62)	8	(5 - 13)	63	(54 - 72)
Indiana	60	(52 - 71)	27	(22 - 33)	30	(24 - 37)	4	(2 - 7)	33	(27 - 41)
Iowa	28	(23 - 32)	12	(10 - 14)	14	(11 - 18)	2	(1 - 3)	16	(13 - 19)
Kansas	27	(23 - 32)	13	(10 - 16)	12	(9 - 15)	2	(1 - 4)	14	(11 - 18)
Kentucky	33	(28 - 40)	17	(14 - 20)	14	(11 - 18)	3	(1 - 5)	17	(13 - 21)
Louisiana	37	(32 - 44)	17	(14 - 20)	18	(15 - 22)	3	(1 - 5)	21	(17 - 26)
Maine	11	(10 - 13)	5	(4 - 7)	5	(4 - 7)	1	(0 - 1)	6	(5 - 8)
Maryland	47	(39 - 56)	23	(19 - 29)	20	(15 - 26)	3	(2 - 6)	23	(18 - 30)
Massachusetts	71	(61 - 83)	33	(28 - 39)	33	(26 - 41)	5	(3 - 10)	38	(30 - 48)
Michigan	95	(87 - 104)	44	(39 - 49)	46	(40 - 52)	6	(3 - 9)	52	(45 - 59)
Minnesota	46	(38 - 53)	22	(18 - 27)	20	(16 - 26)	3	(2 - 6)	23	(18 - 29)
Mississippi	23	(20 - 27)	11	(9 - 14)	10	(8 - 13)	2	(1 - 3)	12	(10 - 15)
Missouri	47	(40 - 56)	24	(20 - 29)	19	(15 - 25)	4	(2 - 7)	23	(18 - 29)
Montana	12	(10 - 13)	5	(5 - 6)	5	(4 - 7)	1	(0 - 1)	6	(5 - 8)

(continued)

Table 5 Average Annual Marijuana Initiates, by Age Group and State: Estimated Numbers (in Thousands), Based on 2010 and 2011 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	16	(13 - 18)	7	(5 - 8)	8	(6 - 10)	1	(1 - 2)	9	(7 - 11)
Nevada	27	(22 - 32)	13	(10 - 16)	12	(9 - 15)	2	(1 - 4)	13	(11 - 17)
New Hampshire	13	(11 - 16)	7	(6 - 8)	6	(5 - 8)	1	(0 - 1)	7	(5 - 8)
New Jersey	82	(70 - 97)	41	(33 - 49)	36	(28 - 45)	6	(3 - 12)	42	(33 - 53)
New Mexico	22	(19 - 25)	11	(9 - 13)	9	(7 - 11)	2	(1 - 3)	11	(9 - 13)
New York	175	(159 - 194)	77	(68 - 86)	85	(74 - 98)	13	(8 - 21)	99	(85 - 114)
North Carolina	85	(72 - 99)	40	(33 - 49)	39	(31 - 49)	5	(3 - 9)	44	(35 - 55)
North Dakota	6	(5 - 8)	2	(2 - 3)	4	(3 - 5)	0	(0 - 1)	4	(3 - 5)
Ohio	102	(93 - 112)	47	(42 - 52)	48	(42 - 55)	7	(4 - 11)	55	(48 - 63)
Oklahoma	31	(26 - 37)	15	(12 - 18)	14	(11 - 17)	3	(1 - 5)	16	(13 - 21)
Oregon	34	(29 - 40)	16	(14 - 20)	15	(12 - 19)	2	(1 - 4)	18	(14 - 22)
Pennsylvania	113	(103 - 126)	48	(42 - 53)	59	(51 - 68)	7	(4 - 12)	66	(57 - 76)
Rhode Island	11	(10 - 13)	5	(4 - 6)	5	(4 - 7)	1	(0 - 2)	6	(5 - 8)
South Carolina	39	(33 - 45)	19	(15 - 22)	17	(14 - 22)	3	(1 - 5)	20	(16 - 25)
South Dakota	7	(6 - 9)	3	(3 - 4)	4	(3 - 5)	0	(0 - 1)	4	(3 - 5)
Tennessee	47	(39 - 56)	22	(18 - 27)	22	(17 - 28)	4	(2 - 7)	25	(20 - 32)
Texas	233	(212 - 257)	106	(94 - 119)	113	(98 - 129)	15	(9 - 25)	127	(111 - 147)
Utah	24	(20 - 29)	12	(10 - 15)	11	(8 - 14)	2	(1 - 3)	12	(10 - 16)
Vermont	6	(5 - 7)	3	(3 - 4)	3	(2 - 3)	0	(0 - 1)	3	(2 - 4)
Virginia	70	(59 - 82)	31	(26 - 37)	34	(27 - 43)	5	(3 - 9)	39	(31 - 49)
Washington	61	(52 - 72)	30	(25 - 36)	26	(21 - 33)	5	(3 - 8)	31	(25 - 39)
West Virginia	16	(13 - 18)	7	(6 - 9)	7	(6 - 9)	1	(1 - 2)	8	(7 - 10)
Wisconsin	44	(37 - 53)	21	(17 - 26)	20	(15 - 26)	3	(1 - 6)	23	(18 - 29)
Wyoming	5	(4 - 6)	2	(2 - 3)	2	(2 - 3)	0	(0 - 1)	2	(2 - 3)

NOTE: Average annual number of marijuana initiates = $X_1 \div 2$, where X_1 is the number of marijuana initiates in the past 24 months.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 6 *Illicit Drug Use Other Than Marijuana in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs*

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Total U.S.	8,518	(8,143 - 8,910)	1,059	(1,000 - 1,121)	2,547	(2,429 - 2,670)	4,912	(4,571 - 5,278)	7,459	(7,090 - 7,847)
Northeast	1,560	(1,432 - 1,700)	164	(149 - 181)	459	(422 - 499)	936	(824 - 1,064)	1,396	(1,270 - 1,534)
Midwest	1,802	(1,682 - 1,929)	224	(207 - 242)	561	(524 - 600)	1,017	(912 - 1,133)	1,578	(1,460 - 1,703)
South	2,994	(2,804 - 3,195)	383	(355 - 413)	870	(815 - 929)	1,741	(1,573 - 1,925)	2,611	(2,425 - 2,810)
West	2,163	(1,976 - 2,366)	288	(257 - 322)	656	(599 - 719)	1,219	(1,061 - 1,399)	1,875	(1,695 - 2,073)
Alabama	126	(98 - 160)	17	(14 - 22)	35	(27 - 46)	73	(52 - 103)	108	(82 - 143)
Alaska	20	(16 - 26)	2	(2 - 3)	6	(5 - 8)	12	(8 - 17)	18	(14 - 24)
Arizona	197	(151 - 254)	28	(22 - 35)	54	(41 - 71)	115	(80 - 166)	169	(126 - 226)
Arkansas	94	(74 - 119)	12	(10 - 15)	27	(21 - 33)	55	(39 - 78)	82	(63 - 106)
California	1,083	(939 - 1,249)	159	(135 - 186)	337	(291 - 388)	588	(470 - 735)	925	(786 - 1,087)
Colorado	165	(132 - 205)	17	(13 - 22)	54	(43 - 67)	94	(68 - 129)	148	(116 - 188)
Connecticut	104	(80 - 134)	11	(8 - 14)	29	(22 - 37)	64	(45 - 91)	93	(70 - 123)
Delaware	30	(23 - 38)	3	(2 - 4)	9	(7 - 12)	17	(12 - 25)	27	(20 - 35)
District of Columbia	23	(17 - 29)	1	(1 - 2)	6	(5 - 8)	15	(11 - 21)	21	(16 - 28)
Florida	490	(422 - 569)	55	(48 - 65)	124	(107 - 144)	310	(251 - 384)	434	(368 - 513)
Georgia	214	(165 - 276)	30	(23 - 38)	64	(49 - 84)	120	(83 - 174)	184	(138 - 245)
Hawaii	33	(25 - 43)	4	(3 - 5)	9	(7 - 12)	20	(14 - 29)	29	(22 - 39)
Idaho	47	(37 - 59)	6	(5 - 8)	14	(11 - 17)	27	(19 - 37)	41	(31 - 53)
Illinois	326	(282 - 377)	43	(37 - 51)	106	(92 - 122)	177	(141 - 223)	283	(241 - 332)
Indiana	195	(153 - 248)	23	(18 - 29)	62	(50 - 79)	110	(76 - 157)	172	(131 - 225)
Iowa	66	(51 - 85)	10	(8 - 13)	22	(17 - 29)	34	(23 - 50)	56	(41 - 75)
Kansas	73	(58 - 91)	10	(8 - 12)	23	(17 - 29)	40	(29 - 56)	63	(49 - 81)
Kentucky	107	(84 - 137)	14	(11 - 18)	33	(26 - 43)	60	(42 - 85)	93	(71 - 123)
Louisiana	127	(102 - 158)	16	(13 - 20)	40	(33 - 50)	70	(50 - 98)	111	(87 - 141)
Maine	37	(29 - 46)	4	(3 - 5)	12	(10 - 15)	21	(15 - 29)	33	(26 - 42)
Maryland	138	(107 - 178)	17	(13 - 22)	38	(29 - 49)	84	(59 - 120)	121	(91 - 161)
Massachusetts	189	(149 - 240)	20	(16 - 25)	65	(51 - 82)	104	(73 - 148)	169	(130 - 220)
Michigan	299	(260 - 344)	36	(31 - 42)	84	(73 - 96)	179	(145 - 221)	263	(224 - 308)
Minnesota	139	(110 - 175)	16	(12 - 20)	44	(35 - 56)	79	(57 - 111)	123	(95 - 159)
Mississippi	79	(63 - 98)	11	(9 - 14)	23	(18 - 30)	44	(32 - 61)	67	(52 - 87)
Missouri	139	(109 - 176)	19	(15 - 24)	46	(36 - 59)	74	(51 - 106)	120	(92 - 157)
Montana	26	(20 - 33)	3	(3 - 4)	8	(6 - 10)	15	(10 - 20)	22	(17 - 29)

(continued)

Table 6 Illicit Drug Use Other Than Marijuana in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	44	(34 - 56)	5	(4 - 7)	14	(11 - 18)	25	(17 - 35)	38	(29 - 50)
Nevada	78	(60 - 101)	10	(8 - 14)	20	(16 - 27)	47	(33 - 67)	67	(50 - 90)
New Hampshire	46	(36 - 57)	4	(3 - 6)	15	(12 - 18)	26	(19 - 37)	41	(32 - 53)
New Jersey	241	(188 - 308)	27	(21 - 34)	67	(52 - 87)	146	(104 - 206)	214	(163 - 280)
New Mexico	64	(51 - 81)	9	(7 - 11)	19	(15 - 23)	37	(26 - 52)	56	(43 - 72)
New York	530	(449 - 626)	54	(45 - 64)	145	(124 - 170)	331	(261 - 419)	476	(397 - 571)
North Carolina	237	(185 - 304)	30	(24 - 38)	73	(56 - 94)	135	(95 - 192)	207	(157 - 273)
North Dakota	13	(9 - 17)	2	(1 - 2)	4	(3 - 6)	7	(4 - 10)	11	(8 - 15)
Ohio	324	(283 - 371)	40	(34 - 46)	96	(84 - 109)	188	(153 - 230)	284	(244 - 330)
Oklahoma	123	(96 - 157)	13	(10 - 17)	36	(29 - 46)	73	(52 - 103)	110	(84 - 144)
Oregon	135	(107 - 170)	14	(11 - 18)	39	(31 - 49)	82	(58 - 113)	121	(94 - 155)
Pennsylvania	346	(299 - 401)	39	(33 - 46)	103	(89 - 120)	204	(164 - 252)	307	(261 - 361)
Rhode Island	43	(34 - 55)	3	(3 - 4)	13	(10 - 16)	27	(19 - 37)	40	(30 - 51)
South Carolina	117	(91 - 149)	14	(11 - 17)	32	(25 - 41)	71	(50 - 99)	103	(79 - 135)
South Dakota	18	(13 - 23)	3	(2 - 3)	6	(4 - 7)	9	(6 - 14)	15	(11 - 20)
Tennessee	193	(152 - 245)	24	(19 - 30)	60	(47 - 77)	110	(78 - 154)	170	(130 - 221)
Texas	623	(540 - 718)	96	(82 - 112)	181	(156 - 210)	346	(277 - 431)	527	(447 - 621)
Utah	63	(49 - 82)	9	(7 - 12)	22	(16 - 29)	33	(22 - 48)	54	(41 - 72)
Vermont	24	(19 - 30)	2	(2 - 3)	8	(7 - 10)	14	(10 - 19)	22	(17 - 28)
Virginia	219	(172 - 279)	23	(18 - 29)	69	(55 - 88)	127	(89 - 179)	196	(151 - 255)
Washington	238	(190 - 298)	25	(19 - 32)	72	(58 - 89)	142	(103 - 194)	214	(167 - 273)
West Virginia	54	(43 - 68)	6	(5 - 8)	17	(14 - 22)	31	(22 - 43)	48	(38 - 62)
Wisconsin	168	(133 - 212)	19	(15 - 24)	53	(42 - 67)	96	(69 - 133)	149	(115 - 193)
Wyoming	13	(10 - 17)	2	(1 - 2)	4	(3 - 5)	8	(5 - 11)	11	(9 - 15)

NOTE: Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 7 Cocaine Use in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	4,195	(3,947 - 4,458)	234	(208 - 264)	1,578	(1,485 - 1,678)	2,383	(2,161 - 2,626)	3,961	(3,715 - 4,223)
Northeast	895	(797 - 1,007)	40	(33 - 47)	321	(288 - 358)	535	(449 - 638)	856	(759 - 966)
Midwest	794	(712 - 884)	45	(38 - 53)	296	(266 - 330)	452	(379 - 539)	749	(669 - 837)
South	1,347	(1,213 - 1,497)	78	(67 - 91)	484	(435 - 538)	785	(663 - 930)	1,270	(1,139 - 1,416)
West	1,159	(1,028 - 1,305)	71	(58 - 87)	477	(425 - 536)	610	(501 - 743)	1,087	(959 - 1,231)
Alabama	36	(23 - 55)	2	(1 - 4)	12	(8 - 19)	21	(13 - 36)	34	(22 - 52)
Alaska	11	(8 - 15)	1	(0 - 1)	4	(3 - 6)	6	(4 - 10)	10	(7 - 14)
Arizona	92	(65 - 130)	6	(4 - 9)	36	(26 - 50)	51	(32 - 79)	87	(61 - 123)
Arkansas	26	(18 - 37)	2	(1 - 3)	9	(6 - 13)	15	(9 - 24)	24	(16 - 35)
California	623	(519 - 746)	41	(31 - 55)	253	(211 - 302)	329	(248 - 436)	582	(480 - 703)
Colorado	105	(80 - 137)	5	(3 - 7)	50	(38 - 65)	50	(33 - 75)	100	(75 - 131)
Connecticut	57	(41 - 79)	3	(2 - 4)	21	(15 - 29)	33	(21 - 50)	54	(39 - 75)
Delaware	14	(10 - 19)	1	(0 - 1)	5	(3 - 7)	8	(6 - 13)	13	(9 - 18)
District of Columbia	16	(11 - 22)	0	(0 - 0)	4	(3 - 5)	12	(8 - 18)	16	(11 - 22)
Florida	255	(210 - 311)	12	(9 - 16)	93	(77 - 113)	150	(113 - 199)	243	(199 - 299)
Georgia	90	(61 - 132)	5	(3 - 8)	29	(19 - 43)	56	(35 - 90)	85	(57 - 126)
Hawaii	17	(12 - 25)	1	(0 - 1)	5	(4 - 8)	11	(7 - 18)	17	(11 - 24)
Idaho	12	(8 - 17)	1	(1 - 2)	5	(4 - 8)	5	(3 - 10)	11	(7 - 16)
Illinois	192	(159 - 233)	10	(8 - 13)	65	(54 - 80)	117	(90 - 153)	183	(150 - 222)
Indiana	54	(36 - 80)	4	(2 - 6)	22	(15 - 33)	28	(16 - 48)	50	(33 - 75)
Iowa	25	(17 - 38)	2	(1 - 3)	11	(7 - 16)	13	(7 - 22)	23	(15 - 35)
Kansas	33	(23 - 46)	2	(2 - 3)	14	(10 - 19)	17	(10 - 26)	30	(21 - 43)
Kentucky	44	(31 - 63)	3	(2 - 4)	17	(12 - 25)	24	(15 - 38)	41	(29 - 60)
Louisiana	50	(35 - 71)	2	(1 - 3)	16	(11 - 22)	32	(20 - 50)	48	(33 - 68)
Maine	20	(15 - 27)	1	(1 - 1)	8	(6 - 10)	11	(8 - 17)	19	(14 - 26)
Maryland	73	(51 - 105)	3	(2 - 5)	23	(16 - 34)	46	(30 - 73)	70	(48 - 101)
Massachusetts	119	(88 - 161)	5	(4 - 8)	45	(33 - 60)	69	(46 - 103)	114	(83 - 155)
Michigan	122	(100 - 149)	7	(5 - 9)	46	(38 - 55)	69	(51 - 93)	115	(93 - 141)
Minnesota	56	(38 - 81)	3	(2 - 5)	21	(14 - 31)	32	(20 - 51)	53	(36 - 77)
Mississippi	34	(24 - 49)	2	(1 - 2)	10	(7 - 15)	22	(14 - 35)	32	(22 - 47)
Missouri	69	(49 - 98)	4	(3 - 6)	27	(19 - 38)	39	(24 - 61)	65	(46 - 93)
Montana	12	(8 - 16)	1	(0 - 1)	5	(4 - 7)	6	(4 - 9)	11	(8 - 15)

(continued)

Table 7 Cocaine Use in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	21	(15 - 29)	1	(1 - 2)	9	(6 - 12)	11	(7 - 17)	19	(14 - 28)
Nevada	41	(29 - 57)	2	(2 - 4)	16	(11 - 22)	23	(15 - 35)	38	(27 - 54)
New Hampshire	25	(19 - 33)	1	(1 - 2)	11	(8 - 14)	13	(9 - 19)	24	(17 - 32)
New Jersey	152	(111 - 208)	7	(5 - 10)	50	(37 - 69)	94	(63 - 141)	145	(105 - 200)
New Mexico	34	(25 - 46)	2	(2 - 4)	14	(10 - 18)	18	(12 - 27)	32	(23 - 43)
New York	303	(250 - 368)	13	(10 - 17)	106	(88 - 127)	184	(140 - 243)	290	(238 - 354)
North Carolina	128	(91 - 180)	7	(4 - 10)	45	(32 - 63)	76	(49 - 118)	121	(85 - 172)
North Dakota	5	(3 - 8)	0	(0 - 1)	2	(1 - 3)	2	(1 - 5)	5	(3 - 7)
Ohio	140	(113 - 173)	7	(6 - 10)	49	(40 - 59)	84	(62 - 113)	133	(107 - 165)
Oklahoma	38	(26 - 54)	3	(2 - 4)	15	(11 - 23)	20	(12 - 32)	35	(24 - 51)
Oregon	68	(52 - 90)	4	(3 - 5)	29	(22 - 38)	35	(24 - 52)	65	(49 - 86)
Pennsylvania	182	(148 - 225)	8	(6 - 11)	64	(52 - 79)	110	(82 - 148)	174	(140 - 217)
Rhode Island	23	(17 - 31)	1	(1 - 1)	9	(7 - 12)	13	(9 - 19)	22	(16 - 30)
South Carolina	52	(36 - 75)	2	(2 - 3)	16	(11 - 23)	34	(22 - 53)	50	(34 - 72)
South Dakota	6	(4 - 10)	0	(0 - 1)	3	(2 - 4)	3	(2 - 6)	6	(4 - 9)
Tennessee	64	(44 - 94)	4	(2 - 6)	21	(14 - 32)	39	(25 - 63)	61	(41 - 89)
Texas	306	(250 - 374)	25	(19 - 33)	122	(100 - 149)	159	(116 - 216)	281	(227 - 346)
Utah	30	(21 - 43)	2	(1 - 3)	14	(10 - 20)	14	(9 - 23)	28	(19 - 40)
Vermont	15	(11 - 19)	1	(0 - 1)	7	(5 - 9)	7	(5 - 11)	14	(11 - 18)
Virginia	100	(73 - 138)	5	(3 - 7)	37	(27 - 51)	58	(38 - 88)	96	(69 - 133)
Washington	109	(82 - 145)	5	(4 - 8)	44	(33 - 59)	59	(40 - 87)	103	(77 - 139)
West Virginia	22	(16 - 30)	1	(1 - 2)	9	(7 - 12)	11	(7 - 18)	20	(15 - 28)
Wisconsin	70	(50 - 98)	4	(3 - 6)	29	(20 - 40)	38	(24 - 59)	67	(47 - 94)
Wyoming	5	(4 - 8)	0	(0 - 1)	2	(2 - 3)	3	(1 - 5)	5	(3 - 8)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 8 Nonmedical Use of Pain Relievers in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	11,692	(11,261 - 12,139)	1,503	(1,433 - 1,576)	3,566	(3,431 - 3,706)	6,624	(6,235 - 7,036)	10,190	(9,768 - 10,629)
Northeast	1,948	(1,811 - 2,097)	221	(202 - 241)	622	(579 - 667)	1,106	(988 - 1,238)	1,728	(1,594 - 1,874)
Midwest	2,600	(2,458 - 2,748)	336	(315 - 359)	833	(790 - 878)	1,431	(1,307 - 1,566)	2,264	(2,125 - 2,410)
South	4,110	(3,887 - 4,346)	552	(517 - 588)	1,241	(1,175 - 1,311)	2,318	(2,122 - 2,531)	3,559	(3,342 - 3,789)
West	3,034	(2,804 - 3,282)	395	(360 - 432)	871	(807 - 939)	1,768	(1,564 - 1,999)	2,639	(2,415 - 2,884)
Alabama	175	(143 - 212)	25	(20 - 31)	53	(43 - 66)	96	(72 - 128)	150	(120 - 186)
Alaska	30	(25 - 36)	4	(3 - 5)	9	(7 - 11)	17	(13 - 22)	26	(21 - 32)
Arizona	302	(246 - 370)	43	(35 - 54)	81	(65 - 100)	179	(133 - 238)	259	(205 - 326)
Arkansas	133	(110 - 160)	18	(15 - 22)	40	(33 - 48)	74	(56 - 98)	115	(93 - 141)
California	1,436	(1,269 - 1,625)	190	(165 - 218)	405	(355 - 462)	841	(697 - 1,014)	1,247	(1,084 - 1,433)
Colorado	250	(207 - 302)	29	(23 - 36)	78	(65 - 94)	143	(108 - 189)	222	(180 - 272)
Connecticut	131	(105 - 162)	13	(11 - 17)	40	(32 - 49)	77	(57 - 104)	117	(92 - 148)
Delaware	42	(34 - 51)	4	(3 - 5)	14	(11 - 17)	24	(18 - 32)	38	(31 - 47)
District of Columbia	25	(20 - 30)	1	(1 - 2)	8	(6 - 9)	16	(12 - 21)	23	(19 - 29)
Florida	643	(567 - 729)	75	(65 - 86)	164	(142 - 188)	404	(337 - 485)	568	(494 - 654)
Georgia	301	(246 - 368)	44	(35 - 55)	94	(74 - 119)	163	(122 - 218)	257	(205 - 322)
Hawaii	42	(33 - 53)	5	(4 - 7)	11	(9 - 14)	26	(19 - 35)	37	(28 - 47)
Idaho	72	(60 - 87)	10	(8 - 12)	21	(17 - 26)	42	(32 - 54)	63	(51 - 77)
Illinois	433	(381 - 492)	55	(47 - 63)	145	(128 - 164)	233	(191 - 285)	378	(328 - 436)
Indiana	303	(249 - 367)	37	(30 - 46)	104	(86 - 125)	162	(120 - 216)	266	(214 - 328)
Iowa	92	(74 - 113)	14	(11 - 17)	32	(26 - 40)	46	(33 - 64)	78	(61 - 99)
Kansas	105	(87 - 127)	14	(12 - 18)	34	(27 - 41)	57	(43 - 75)	91	(74 - 112)
Kentucky	161	(133 - 194)	22	(18 - 28)	50	(40 - 61)	88	(67 - 116)	138	(112 - 171)
Louisiana	180	(151 - 214)	24	(19 - 29)	61	(51 - 73)	95	(73 - 124)	156	(128 - 190)
Maine	47	(38 - 58)	5	(4 - 7)	15	(12 - 18)	27	(20 - 36)	42	(33 - 52)
Maryland	186	(151 - 230)	21	(17 - 27)	57	(45 - 70)	109	(81 - 145)	165	(131 - 208)
Massachusetts	239	(196 - 291)	24	(19 - 31)	81	(66 - 100)	134	(101 - 177)	215	(173 - 266)
Michigan	424	(379 - 475)	52	(45 - 59)	129	(117 - 143)	243	(204 - 290)	372	(328 - 422)
Minnesota	202	(167 - 242)	26	(21 - 32)	66	(54 - 80)	110	(84 - 145)	176	(143 - 215)
Mississippi	106	(88 - 129)	17	(14 - 21)	32	(26 - 40)	57	(43 - 75)	89	(72 - 111)
Missouri	240	(200 - 287)	32	(26 - 39)	77	(63 - 93)	131	(100 - 170)	208	(169 - 254)
Montana	40	(33 - 48)	6	(5 - 7)	12	(10 - 14)	23	(17 - 30)	34	(28 - 42)

(continued)

Table 8 Nonmedical Use of Pain Relievers in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Nebraska	62	(51 - 76)	7	(6 - 9)	20	(16 - 24)	35	(27 - 46)	55	(44 - 68)
Nevada	123	(100 - 151)	17	(13 - 21)	33	(26 - 40)	74	(55 - 99)	107	(85 - 134)
New Hampshire	52	(43 - 62)	6	(5 - 8)	17	(15 - 21)	28	(21 - 37)	45	(37 - 56)
New Jersey	303	(248 - 371)	36	(28 - 46)	96	(78 - 117)	172	(129 - 228)	267	(215 - 332)
New Mexico	91	(75 - 111)	14	(11 - 18)	25	(20 - 31)	51	(38 - 69)	77	(61 - 96)
New York	653	(570 - 748)	70	(60 - 82)	197	(172 - 225)	386	(314 - 474)	583	(502 - 677)
North Carolina	312	(252 - 385)	46	(38 - 57)	91	(72 - 114)	175	(128 - 237)	266	(208 - 337)
North Dakota	21	(17 - 26)	3	(2 - 3)	7	(6 - 9)	11	(8 - 15)	19	(15 - 23)
Ohio	480	(431 - 534)	66	(58 - 75)	144	(130 - 160)	269	(227 - 319)	414	(367 - 467)
Oklahoma	157	(129 - 191)	21	(17 - 26)	47	(38 - 58)	89	(67 - 118)	136	(110 - 169)
Oregon	207	(170 - 250)	21	(17 - 26)	62	(51 - 73)	124	(94 - 163)	185	(150 - 228)
Pennsylvania	449	(398 - 506)	58	(50 - 66)	150	(131 - 170)	242	(200 - 292)	391	(342 - 448)
Rhode Island	46	(38 - 56)	4	(3 - 5)	16	(13 - 20)	26	(20 - 34)	42	(34 - 52)
South Carolina	176	(145 - 213)	21	(17 - 26)	53	(44 - 65)	102	(77 - 134)	155	(125 - 191)
South Dakota	25	(19 - 31)	4	(3 - 4)	7	(6 - 9)	14	(10 - 19)	21	(16 - 27)
Tennessee	264	(219 - 318)	34	(28 - 42)	88	(72 - 106)	142	(108 - 187)	230	(186 - 282)
Texas	873	(774 - 985)	132	(114 - 152)	265	(234 - 300)	476	(392 - 578)	741	(646 - 850)
Utah	94	(77 - 115)	15	(11 - 19)	31	(24 - 39)	49	(37 - 65)	80	(64 - 99)
Vermont	28	(23 - 33)	3	(2 - 4)	10	(8 - 12)	15	(11 - 20)	25	(20 - 30)
Virginia	302	(248 - 366)	36	(29 - 45)	101	(82 - 123)	165	(124 - 218)	266	(214 - 328)
Washington	323	(268 - 390)	39	(31 - 48)	97	(81 - 117)	187	(142 - 247)	285	(231 - 349)
West Virginia	75	(62 - 90)	9	(8 - 12)	23	(19 - 28)	42	(32 - 55)	65	(53 - 80)
Wisconsin	214	(175 - 262)	27	(22 - 34)	67	(54 - 83)	120	(90 - 160)	187	(149 - 234)
Wyoming	21	(18 - 26)	3	(2 - 3)	6	(5 - 8)	12	(9 - 16)	19	(15 - 23)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 9 Alcohol Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Total U.S.	132,390	(130,948 - 133,810)	3,323	(3,212 - 3,436)	20,866	(20,600 - 21,129)	108,201	(106,815 - 109,564)	129,067	(127,640 - 130,470)
Northeast	26,597	(26,041 - 27,146)	676	(638 - 716)	4,053	(3,958 - 4,146)	21,868	(21,350 - 22,380)	25,921	(25,366 - 26,472)
Midwest	30,163	(29,601 - 30,730)	723	(690 - 756)	4,767	(4,667 - 4,866)	24,673	(24,155 - 25,191)	29,441	(28,873 - 30,007)
South	45,406	(44,475 - 46,324)	1,133	(1,081 - 1,186)	7,233	(7,065 - 7,400)	37,040	(36,188 - 37,876)	44,273	(43,337 - 45,192)
West	30,223	(29,505 - 30,936)	791	(741 - 844)	4,812	(4,677 - 4,946)	24,620	(23,955 - 25,278)	29,432	(28,714 - 30,142)
Alabama	1,699	(1,574 - 1,825)	44	(37 - 53)	271	(251 - 291)	1,383	(1,265 - 1,503)	1,654	(1,531 - 1,780)
Alaska	303	(283 - 322)	7	(6 - 9)	50	(46 - 53)	245	(227 - 263)	295	(276 - 314)
Arizona	2,596	(2,403 - 2,792)	68	(57 - 82)	390	(360 - 419)	2,138	(1,956 - 2,319)	2,528	(2,335 - 2,719)
Arkansas	977	(898 - 1,058)	30	(25 - 36)	156	(144 - 168)	791	(717 - 867)	947	(868 - 1,027)
California	15,285	(14,730 - 15,837)	428	(386 - 473)	2,570	(2,470 - 2,667)	12,287	(11,763 - 12,809)	14,857	(14,304 - 15,406)
Colorado	2,562	(2,428 - 2,692)	60	(50 - 71)	394	(372 - 414)	2,108	(1,981 - 2,229)	2,502	(2,369 - 2,629)
Connecticut	1,830	(1,733 - 1,925)	48	(41 - 57)	265	(251 - 279)	1,517	(1,425 - 1,604)	1,782	(1,685 - 1,874)
Delaware	409	(384 - 434)	10	(8 - 12)	65	(61 - 68)	334	(310 - 358)	399	(374 - 424)
District of Columbia	331	(313 - 349)	5	(4 - 6)	68	(65 - 71)	258	(241 - 275)	327	(308 - 344)
Florida	8,083	(7,794 - 8,370)	171	(155 - 188)	1,100	(1,057 - 1,143)	6,812	(6,534 - 7,087)	7,912	(7,623 - 8,200)
Georgia	3,689	(3,440 - 3,939)	96	(80 - 115)	579	(537 - 620)	3,014	(2,778 - 3,249)	3,592	(3,346 - 3,839)
Hawaii	521	(484 - 558)	13	(11 - 16)	82	(77 - 88)	425	(390 - 460)	508	(471 - 544)
Idaho	619	(578 - 661)	16	(14 - 20)	89	(82 - 96)	514	(475 - 553)	603	(561 - 644)
Illinois	5,857	(5,661 - 6,050)	141	(127 - 156)	926	(895 - 956)	4,790	(4,604 - 4,973)	5,716	(5,521 - 5,908)
Indiana	2,714	(2,538 - 2,889)	61	(51 - 74)	442	(413 - 470)	2,210	(2,042 - 2,376)	2,652	(2,476 - 2,825)
Iowa	1,362	(1,280 - 1,444)	38	(32 - 44)	235	(221 - 248)	1,089	(1,011 - 1,166)	1,324	(1,241 - 1,404)
Kansas	1,193	(1,117 - 1,268)	31	(26 - 36)	204	(191 - 216)	958	(888 - 1,028)	1,162	(1,087 - 1,236)
Kentucky	1,456	(1,341 - 1,576)	43	(36 - 52)	235	(216 - 254)	1,178	(1,069 - 1,291)	1,413	(1,297 - 1,531)
Louisiana	1,743	(1,633 - 1,854)	52	(44 - 60)	306	(287 - 325)	1,386	(1,281 - 1,490)	1,692	(1,581 - 1,801)
Maine	621	(582 - 659)	12	(10 - 14)	82	(77 - 87)	527	(490 - 564)	609	(570 - 647)
Maryland	2,633	(2,469 - 2,795)	57	(48 - 68)	397	(372 - 421)	2,179	(2,024 - 2,330)	2,576	(2,412 - 2,736)
Massachusetts	3,531	(3,356 - 3,700)	78	(67 - 91)	538	(509 - 566)	2,915	(2,750 - 3,071)	3,453	(3,279 - 3,619)
Michigan	4,446	(4,301 - 4,590)	104	(94 - 115)	693	(670 - 715)	3,649	(3,510 - 3,787)	4,342	(4,198 - 4,485)
Minnesota	2,621	(2,482 - 2,759)	55	(46 - 65)	392	(369 - 413)	2,174	(2,042 - 2,301)	2,566	(2,426 - 2,700)
Mississippi	966	(895 - 1,039)	29	(25 - 35)	175	(162 - 188)	762	(695 - 830)	937	(866 - 1,009)
Missouri	2,475	(2,316 - 2,634)	71	(60 - 83)	399	(374 - 422)	2,005	(1,856 - 2,154)	2,404	(2,246 - 2,561)
Montana	498	(472 - 523)	13	(11 - 15)	76	(72 - 79)	410	(385 - 433)	485	(460 - 510)

(continued)

Table 9 Alcohol Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Nebraska	792	(743 - 840)	18	(16 - 22)	136	(128 - 144)	637	(591 - 682)	773	(724 - 821)
Nevada	1,185	(1,104 - 1,265)	28	(23 - 34)	161	(149 - 172)	996	(920 - 1,071)	1,157	(1,076 - 1,236)
New Hampshire	695	(657 - 732)	17	(15 - 20)	104	(99 - 109)	573	(537 - 608)	677	(639 - 714)
New Jersey	4,160	(3,915 - 4,400)	118	(101 - 138)	562	(526 - 596)	3,480	(3,247 - 3,704)	4,042	(3,798 - 4,278)
New Mexico	715	(660 - 772)	21	(17 - 25)	123	(114 - 132)	571	(519 - 624)	694	(639 - 750)
New York	9,030	(8,727 - 9,328)	247	(224 - 272)	1,451	(1,403 - 1,497)	7,332	(7,041 - 7,619)	8,783	(8,480 - 9,081)
North Carolina	3,834	(3,572 - 4,097)	95	(80 - 113)	626	(586 - 666)	3,112	(2,866 - 3,357)	3,738	(3,476 - 3,997)
North Dakota	326	(309 - 344)	7	(6 - 8)	66	(62 - 69)	254	(237 - 270)	320	(302 - 337)
Ohio	5,040	(4,873 - 5,206)	125	(113 - 137)	774	(749 - 799)	4,141	(3,982 - 4,299)	4,916	(4,749 - 5,081)
Oklahoma	1,355	(1,257 - 1,454)	38	(32 - 46)	243	(225 - 260)	1,074	(983 - 1,166)	1,317	(1,219 - 1,414)
Oregon	1,947	(1,843 - 2,048)	42	(36 - 49)	264	(248 - 279)	1,641	(1,541 - 1,736)	1,905	(1,801 - 2,004)
Pennsylvania	5,851	(5,657 - 6,043)	135	(122 - 150)	902	(868 - 935)	4,814	(4,629 - 4,997)	5,716	(5,522 - 5,907)
Rhode Island	552	(521 - 582)	12	(10 - 14)	94	(90 - 99)	446	(417 - 474)	540	(509 - 569)
South Carolina	1,843	(1,716 - 1,970)	43	(36 - 52)	286	(267 - 304)	1,513	(1,393 - 1,633)	1,799	(1,673 - 1,925)
South Dakota	373	(351 - 395)	10	(9 - 12)	63	(59 - 67)	299	(278 - 319)	362	(340 - 384)
Tennessee	2,192	(2,019 - 2,369)	55	(46 - 65)	355	(328 - 382)	1,781	(1,619 - 1,947)	2,137	(1,964 - 2,311)
Texas	10,005	(9,649 - 10,361)	266	(239 - 296)	1,682	(1,618 - 1,745)	8,057	(7,715 - 8,396)	9,739	(9,383 - 10,092)
Utah	553	(491 - 620)	20	(15 - 26)	117	(102 - 133)	417	(359 - 480)	533	(471 - 600)
Vermont	327	(309 - 344)	8	(7 - 9)	55	(52 - 58)	264	(247 - 280)	319	(301 - 336)
Virginia	3,585	(3,362 - 3,804)	78	(66 - 93)	587	(554 - 619)	2,919	(2,708 - 3,124)	3,506	(3,284 - 3,722)
Washington	3,208	(3,021 - 3,391)	69	(58 - 82)	461	(432 - 488)	2,678	(2,501 - 2,849)	3,139	(2,952 - 3,320)
West Virginia	607	(557 - 659)	19	(16 - 23)	102	(95 - 109)	486	(439 - 535)	588	(538 - 639)
Wisconsin	2,965	(2,807 - 3,118)	62	(52 - 74)	438	(414 - 460)	2,465	(2,313 - 2,607)	2,903	(2,744 - 3,052)
Wyoming	231	(216 - 246)	6	(5 - 7)	37	(34 - 39)	189	(174 - 203)	225	(210 - 240)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 10 Binge Alcohol Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	58,440	(57,404 - 59,485)	1,881	(1,799 - 1,966)	13,725	(13,463 - 13,988)	42,835	(41,853 - 43,829)	56,559	(55,527 - 57,600)
Northeast	11,518	(11,152 - 11,893)	381	(353 - 412)	2,673	(2,592 - 2,754)	8,464	(8,118 - 8,820)	11,137	(10,774 - 11,510)
Midwest	13,730	(13,383 - 14,082)	419	(394 - 446)	3,267	(3,190 - 3,344)	10,044	(9,713 - 10,384)	13,310	(12,965 - 13,661)
South	20,012	(19,451 - 20,581)	622	(583 - 663)	4,632	(4,508 - 4,757)	14,758	(14,220 - 15,306)	19,390	(18,832 - 19,956)
West	13,181	(12,710 - 13,662)	458	(417 - 503)	3,153	(3,040 - 3,268)	9,569	(9,122 - 10,031)	12,722	(12,255 - 13,200)
Alabama	776	(689 - 871)	25	(20 - 31)	162	(142 - 182)	589	(510 - 677)	751	(664 - 845)
Alaska	135	(122 - 150)	4	(3 - 6)	35	(31 - 38)	96	(84 - 110)	131	(117 - 146)
Arizona	1,195	(1,060 - 1,341)	39	(31 - 49)	260	(232 - 289)	897	(774 - 1,033)	1,156	(1,022 - 1,300)
Arkansas	454	(403 - 510)	17	(14 - 21)	97	(86 - 109)	340	(293 - 393)	437	(386 - 493)
California	6,743	(6,363 - 7,138)	249	(217 - 286)	1,654	(1,561 - 1,748)	4,840	(4,481 - 5,220)	6,494	(6,116 - 6,888)
Colorado	1,020	(922 - 1,125)	30	(24 - 37)	250	(227 - 273)	740	(652 - 837)	990	(893 - 1,093)
Connecticut	785	(707 - 869)	29	(23 - 36)	180	(164 - 195)	577	(505 - 655)	756	(679 - 839)
Delaware	174	(156 - 193)	6	(5 - 7)	42	(38 - 45)	127	(110 - 145)	168	(151 - 187)
District of Columbia	169	(155 - 185)	2	(2 - 3)	46	(42 - 49)	121	(107 - 136)	167	(152 - 182)
Florida	3,158	(2,956 - 3,371)	94	(82 - 106)	654	(614 - 694)	2,410	(2,219 - 2,614)	3,064	(2,864 - 3,277)
Georgia	1,540	(1,372 - 1,722)	51	(41 - 64)	376	(335 - 419)	1,112	(963 - 1,279)	1,488	(1,323 - 1,668)
Hawaii	250	(224 - 278)	8	(6 - 10)	56	(51 - 62)	186	(162 - 211)	242	(216 - 269)
Idaho	273	(243 - 306)	10	(8 - 12)	60	(53 - 67)	204	(176 - 234)	263	(234 - 296)
Illinois	2,697	(2,558 - 2,841)	84	(74 - 95)	638	(608 - 669)	1,974	(1,841 - 2,114)	2,613	(2,475 - 2,756)
Indiana	1,209	(1,084 - 1,344)	35	(28 - 44)	303	(274 - 332)	871	(759 - 995)	1,174	(1,050 - 1,307)
Iowa	670	(609 - 734)	23	(19 - 28)	176	(162 - 190)	471	(416 - 531)	647	(586 - 710)
Kansas	540	(487 - 596)	18	(15 - 23)	140	(127 - 153)	382	(334 - 434)	521	(469 - 578)
Kentucky	748	(670 - 834)	26	(21 - 32)	174	(157 - 192)	549	(476 - 629)	722	(645 - 807)
Louisiana	841	(764 - 923)	25	(21 - 31)	196	(179 - 214)	619	(548 - 697)	816	(739 - 897)
Maine	253	(227 - 282)	7	(6 - 9)	55	(50 - 61)	191	(166 - 218)	246	(220 - 275)
Maryland	1,003	(898 - 1,118)	30	(24 - 37)	245	(221 - 271)	728	(632 - 835)	974	(869 - 1,088)
Massachusetts	1,516	(1,379 - 1,661)	46	(38 - 55)	364	(334 - 393)	1,106	(979 - 1,243)	1,470	(1,334 - 1,614)
Michigan	2,100	(1,992 - 2,213)	62	(55 - 71)	480	(459 - 501)	1,558	(1,453 - 1,668)	2,038	(1,930 - 2,151)
Minnesota	1,155	(1,053 - 1,262)	35	(28 - 42)	277	(254 - 300)	843	(751 - 944)	1,120	(1,019 - 1,226)
Mississippi	459	(408 - 516)	15	(12 - 19)	108	(96 - 120)	337	(290 - 389)	444	(393 - 500)
Missouri	1,128	(1,017 - 1,246)	40	(33 - 49)	247	(223 - 273)	840	(739 - 950)	1,087	(978 - 1,205)
Montana	231	(212 - 252)	7	(6 - 9)	56	(51 - 60)	168	(150 - 188)	224	(204 - 245)

(continued)

Table 10 Binge Alcohol Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	365	(329 - 404)	11	(9 - 13)	93	(84 - 101)	262	(229 - 297)	355	(318 - 393)
Nevada	539	(480 - 602)	17	(13 - 23)	108	(96 - 120)	413	(360 - 472)	522	(464 - 584)
New Hampshire	272	(245 - 300)	10	(8 - 12)	70	(65 - 76)	192	(167 - 219)	262	(235 - 290)
New Jersey	1,795	(1,616 - 1,987)	71	(58 - 87)	388	(352 - 426)	1,335	(1,172 - 1,514)	1,724	(1,547 - 1,914)
New Mexico	339	(302 - 380)	12	(9 - 15)	82	(74 - 91)	245	(211 - 283)	327	(290 - 368)
New York	3,891	(3,667 - 4,124)	130	(114 - 148)	910	(863 - 958)	2,850	(2,638 - 3,074)	3,761	(3,538 - 3,994)
North Carolina	1,518	(1,349 - 1,703)	49	(40 - 61)	362	(323 - 402)	1,107	(955 - 1,277)	1,469	(1,300 - 1,651)
North Dakota	169	(154 - 184)	4	(3 - 5)	49	(46 - 53)	116	(103 - 130)	165	(150 - 180)
Ohio	2,224	(2,105 - 2,348)	67	(59 - 76)	523	(500 - 546)	1,634	(1,520 - 1,755)	2,157	(2,039 - 2,281)
Oklahoma	637	(570 - 710)	20	(16 - 26)	162	(145 - 180)	455	(394 - 522)	617	(550 - 689)
Oregon	725	(651 - 805)	20	(16 - 25)	166	(150 - 182)	539	(472 - 613)	705	(632 - 784)
Pennsylvania	2,620	(2,480 - 2,766)	77	(68 - 88)	603	(572 - 634)	1,940	(1,808 - 2,079)	2,543	(2,404 - 2,688)
Rhode Island	250	(226 - 274)	7	(6 - 8)	65	(59 - 70)	178	(157 - 201)	243	(220 - 267)
South Carolina	870	(778 - 968)	22	(17 - 27)	184	(165 - 203)	664	(579 - 757)	848	(756 - 946)
South Dakota	184	(166 - 202)	6	(5 - 8)	46	(42 - 51)	131	(116 - 148)	177	(160 - 195)
Tennessee	973	(856 - 1,102)	31	(24 - 39)	222	(196 - 249)	720	(615 - 840)	942	(825 - 1,069)
Texas	4,850	(4,590 - 5,119)	151	(131 - 172)	1,136	(1,079 - 1,195)	3,563	(3,313 - 3,825)	4,699	(4,440 - 4,968)
Utah	309	(268 - 354)	14	(10 - 19)	85	(73 - 99)	210	(175 - 251)	295	(255 - 340)
Vermont	137	(124 - 150)	4	(4 - 5)	38	(35 - 41)	94	(83 - 107)	132	(120 - 145)
Virginia	1,523	(1,369 - 1,688)	46	(38 - 57)	393	(357 - 430)	1,083	(945 - 1,235)	1,476	(1,324 - 1,640)
Washington	1,318	(1,185 - 1,461)	44	(36 - 54)	317	(289 - 345)	957	(836 - 1,091)	1,274	(1,142 - 1,415)
West Virginia	319	(285 - 356)	11	(9 - 14)	75	(67 - 82)	233	(202 - 268)	308	(274 - 345)
Wisconsin	1,289	(1,164 - 1,422)	34	(28 - 42)	294	(269 - 320)	960	(845 - 1,086)	1,255	(1,130 - 1,387)
Wyoming	102	(92 - 114)	4	(3 - 5)	26	(23 - 28)	73	(64 - 83)	99	(89 - 110)

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 11 *Perceptions of Great Risk of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs*

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	108,820	(107,448 - 110,185)	10,007	(9,845 - 10,170)	11,669	(11,440 - 11,900)	87,145	(85,846 - 88,436)	98,814	—
Northeast	19,153	(18,680 - 19,626)	1,658	(1,608 - 1,708)	1,910	(1,842 - 1,979)	15,585	(15,129 - 16,043)	17,495	(17,030 - 17,961)
Midwest	21,487	(21,050 - 21,925)	2,050	(2,001 - 2,100)	2,222	(2,160 - 2,286)	17,215	(16,797 - 17,633)	19,437	(19,006 - 19,869)
South	42,105	(41,383 - 42,825)	3,881	(3,795 - 3,967)	4,646	(4,533 - 4,759)	33,578	(32,884 - 34,271)	38,224	(37,517 - 38,929)
West	26,076	(25,470 - 26,681)	2,418	(2,341 - 2,495)	2,891	(2,789 - 2,994)	20,767	(20,186 - 21,348)	23,658	(23,063 - 24,253)
Alabama	1,751	(1,635 - 1,869)	168	(155 - 181)	208	(190 - 228)	1,375	(1,267 - 1,485)	1,584	(1,471 - 1,698)
Alaska	227	(210 - 245)	23	(21 - 26)	26	(23 - 30)	177	(161 - 194)	204	(187 - 221)
Arizona	2,322	(2,147 - 2,502)	230	(209 - 252)	251	(226 - 278)	1,841	(1,677 - 2,007)	2,092	(1,923 - 2,265)
Arkansas	1,063	(987 - 1,140)	103	(95 - 112)	113	(103 - 124)	847	(775 - 919)	960	(887 - 1,034)
California	14,244	(13,739 - 14,749)	1,306	(1,241 - 1,372)	1,647	(1,559 - 1,737)	11,291	(10,811 - 11,772)	12,938	(12,444 - 13,432)
Colorado	1,558	(1,430 - 1,690)	141	(127 - 156)	155	(137 - 174)	1,262	(1,142 - 1,385)	1,417	(1,293 - 1,545)
Connecticut	1,248	(1,157 - 1,340)	108	(98 - 119)	107	(95 - 120)	1,033	(948 - 1,119)	1,140	(1,052 - 1,229)
Delaware	303	(280 - 327)	27	(25 - 30)	31	(28 - 35)	245	(224 - 267)	276	(254 - 299)
District of Columbia	216	(200 - 233)	14	(13 - 16)	27	(24 - 31)	175	(159 - 190)	202	(186 - 218)
Florida	7,552	(7,272 - 7,833)	567	(541 - 593)	762	(724 - 802)	6,223	(5,950 - 6,496)	6,986	(6,708 - 7,263)
Georgia	3,479	(3,247 - 3,715)	354	(324 - 384)	395	(358 - 433)	2,731	(2,516 - 2,949)	3,126	(2,901 - 3,354)
Hawaii	453	(418 - 488)	39	(36 - 43)	45	(40 - 50)	368	(336 - 401)	413	(380 - 447)
Idaho	475	(439 - 512)	46	(42 - 51)	54	(49 - 60)	374	(341 - 409)	429	(394 - 464)
Illinois	4,437	(4,264 - 4,612)	440	(419 - 462)	466	(440 - 493)	3,531	(3,365 - 3,699)	3,997	(3,827 - 4,168)
Indiana	2,099	(1,943 - 2,260)	207	(189 - 225)	218	(196 - 242)	1,674	(1,528 - 1,825)	1,892	(1,741 - 2,048)
Iowa	960	(885 - 1,039)	88	(80 - 97)	99	(87 - 111)	773	(702 - 846)	872	(799 - 948)
Kansas	893	(823 - 964)	85	(77 - 94)	97	(86 - 108)	710	(646 - 777)	807	(740 - 876)
Kentucky	1,523	(1,408 - 1,639)	146	(133 - 159)	169	(153 - 186)	1,207	(1,100 - 1,317)	1,377	(1,266 - 1,490)
Louisiana	1,667	(1,564 - 1,771)	166	(154 - 179)	211	(194 - 228)	1,290	(1,193 - 1,387)	1,501	(1,400 - 1,602)
Maine	430	(395 - 466)	34	(30 - 37)	37	(33 - 42)	359	(326 - 393)	397	(362 - 432)
Maryland	2,131	(1,988 - 2,277)	200	(183 - 217)	221	(199 - 243)	1,711	(1,578 - 1,846)	1,932	(1,793 - 2,072)
Massachusetts	2,254	(2,090 - 2,422)	178	(161 - 194)	222	(198 - 247)	1,855	(1,700 - 2,013)	2,076	(1,917 - 2,240)
Michigan	3,221	(3,090 - 3,353)	306	(291 - 322)	350	(331 - 370)	2,564	(2,438 - 2,692)	2,914	(2,786 - 3,044)
Minnesota	1,625	(1,501 - 1,755)	145	(131 - 159)	154	(137 - 173)	1,326	(1,209 - 1,447)	1,480	(1,359 - 1,607)
Mississippi	1,130	(1,062 - 1,199)	122	(113 - 130)	140	(129 - 152)	868	(804 - 933)	1,009	(942 - 1,076)
Missouri	1,811	(1,667 - 1,959)	175	(158 - 193)	197	(176 - 220)	1,439	(1,304 - 1,578)	1,636	(1,497 - 1,780)
Montana	300	(278 - 323)	25	(22 - 27)	27	(24 - 31)	248	(227 - 270)	275	(253 - 298)

(continued)

Table 11 *Perceptions of Great Risk of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)*

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Nebraska	548	(504 - 594)	55	(49 - 60)	56	(50 - 63)	437	(396 - 480)	493	(451 - 538)
Nevada	883	(811 - 958)	84	(75 - 94)	96	(86 - 108)	703	(636 - 772)	799	(730 - 870)
New Hampshire	439	(406 - 473)	37	(34 - 41)	38	(34 - 42)	364	(333 - 396)	402	(370 - 435)
New Jersey	3,128	(2,908 - 3,352)	275	(250 - 300)	290	(260 - 321)	2,563	(2,355 - 2,775)	2,853	(2,639 - 3,071)
New Mexico	801	(749 - 855)	70	(64 - 76)	87	(79 - 95)	645	(595 - 694)	732	(680 - 783)
New York	6,983	(6,694 - 7,275)	611	(581 - 641)	745	(704 - 788)	5,626	(5,346 - 5,909)	6,372	(6,086 - 6,661)
North Carolina	3,425	(3,197 - 3,658)	291	(265 - 317)	343	(311 - 378)	2,791	(2,575 - 3,010)	3,134	(2,912 - 3,361)
North Dakota	200	(185 - 216)	17	(15 - 19)	23	(20 - 26)	160	(146 - 175)	183	(169 - 198)
Ohio	3,766	(3,616 - 3,919)	352	(335 - 370)	360	(339 - 380)	3,054	(2,909 - 3,202)	3,414	(3,266 - 3,564)
Oklahoma	1,328	(1,236 - 1,421)	122	(111 - 133)	157	(141 - 172)	1,049	(964 - 1,135)	1,206	(1,117 - 1,296)
Oregon	1,228	(1,134 - 1,326)	102	(92 - 112)	121	(108 - 135)	1,005	(916 - 1,098)	1,127	(1,034 - 1,222)
Pennsylvania	4,122	(3,945 - 4,301)	371	(351 - 391)	414	(387 - 442)	3,336	(3,167 - 3,509)	3,751	(3,577 - 3,927)
Rhode Island	344	(315 - 373)	29	(27 - 32)	39	(34 - 43)	276	(249 - 303)	314	(287 - 342)
South Carolina	1,701	(1,588 - 1,816)	157	(144 - 170)	187	(171 - 204)	1,357	(1,249 - 1,466)	1,544	(1,434 - 1,657)
South Dakota	237	(217 - 258)	22	(19 - 24)	25	(22 - 29)	190	(172 - 209)	215	(197 - 235)
Tennessee	2,291	(2,127 - 2,459)	207	(190 - 225)	237	(214 - 261)	1,847	(1,692 - 2,004)	2,084	(1,925 - 2,247)
Texas	9,154	(8,828 - 9,481)	940	(897 - 985)	1,084	(1,029 - 1,140)	7,130	(6,814 - 7,448)	8,214	(7,894 - 8,535)
Utah	1,102	(1,035 - 1,169)	136	(125 - 147)	154	(138 - 169)	812	(751 - 873)	966	(901 - 1,031)
Vermont	205	(189 - 222)	15	(14 - 17)	18	(16 - 20)	172	(157 - 188)	190	(175 - 207)
Virginia	2,753	(2,559 - 2,950)	247	(226 - 270)	298	(269 - 329)	2,207	(2,025 - 2,393)	2,505	(2,318 - 2,696)
Washington	2,312	(2,147 - 2,481)	199	(181 - 217)	209	(188 - 232)	1,904	(1,747 - 2,064)	2,113	(1,952 - 2,278)
West Virginia	635	(590 - 681)	50	(45 - 55)	61	(55 - 67)	524	(481 - 568)	585	(541 - 630)
Wisconsin	1,690	(1,552 - 1,834)	158	(143 - 174)	178	(158 - 199)	1,355	(1,225 - 1,490)	1,532	(1,398 - 1,672)
Wyoming	170	(157 - 184)	16	(14 - 17)	18	(16 - 20)	136	(124 - 150)	154	(141 - 168)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 12 Alcohol Use and Binge Alcohol Use in the Past Month among Persons Aged 12 to 20, by State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs

State	Alcohol Use in Past Month (Estimate)	Alcohol Use in Past Month (95% Confidence Interval)	Binge Alcohol Use in Past Month (Estimate)	Binge Alcohol Use in Past Month (95% Confidence Interval)
Total U.S.	9,794	(9,547 - 10,045)	6,249	(6,042 - 6,462)
Northeast	1,999	(1,937 - 2,063)	1,272	(1,215 - 1,330)
Midwest	2,192	(2,134 - 2,251)	1,462	(1,412 - 1,514)
South	3,336	(3,240 - 3,434)	2,055	(1,975 - 2,136)
West	2,267	(2,180 - 2,356)	1,461	(1,386 - 1,539)
Alabama	123	(109 - 137)	74	(64 - 85)
Alaska	22	(20 - 24)	15	(14 - 17)
Arizona	185	(164 - 207)	116	(101 - 133)
Arkansas	78	(70 - 86)	49	(42 - 56)
California	1,250	(1,177 - 1,326)	808	(748 - 873)
Colorado	155	(140 - 170)	91	(80 - 103)
Connecticut	139	(127 - 151)	92	(82 - 103)
Delaware	31	(28 - 34)	20	(18 - 23)
District of Columbia	21	(19 - 23)	13	(11 - 15)
Florida	529	(499 - 560)	317	(293 - 343)
Georgia	256	(228 - 286)	151	(131 - 173)
Hawaii	36	(32 - 40)	24	(21 - 27)
Idaho	46	(41 - 51)	30	(26 - 34)
Illinois	433	(411 - 455)	290	(271 - 311)
Indiana	203	(183 - 225)	136	(120 - 153)
Iowa	108	(98 - 118)	80	(71 - 89)
Kansas	87	(79 - 97)	60	(53 - 67)
Kentucky	112	(100 - 125)	76	(67 - 86)
Louisiana	147	(134 - 161)	83	(74 - 94)
Maine	39	(35 - 43)	26	(23 - 29)
Maryland	167	(151 - 184)	97	(85 - 110)
Massachusetts	233	(214 - 253)	151	(135 - 169)
Michigan	326	(309 - 343)	217	(203 - 232)
Minnesota	164	(149 - 180)	116	(103 - 130)
Mississippi	81	(73 - 91)	46	(39 - 53)
Missouri	182	(164 - 201)	109	(96 - 125)
Montana	35	(32 - 39)	23	(21 - 26)

(continued)

Table 12 Alcohol Use and Binge Alcohol Use in the Past Month among Persons Aged 12 to 20, by State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	Alcohol Use in Past Month (Estimate)	Alcohol Use in Past Month (95% Confidence Interval)	Binge Alcohol Use in Past Month (Estimate)	Alcohol Use in Past Month (95% Confidence Interval)
Nebraska	58	(52 - 64)	40	(35 - 45)
Nevada	92	(82 - 103)	62	(54 - 71)
New Hampshire	57	(53 - 62)	37	(33 - 41)
New Jersey	322	(293 - 353)	217	(193 - 244)
New Mexico	59	(53 - 66)	37	(32 - 42)
New York	703	(667 - 741)	430	(399 - 463)
North Carolina	277	(249 - 307)	158	(138 - 180)
North Dakota	26	(24 - 28)	19	(17 - 21)
Ohio	390	(371 - 410)	255	(238 - 272)
Oklahoma	105	(93 - 117)	61	(52 - 71)
Oregon	116	(106 - 127)	67	(58 - 76)
Pennsylvania	442	(418 - 467)	277	(256 - 300)
Rhode Island	40	(36 - 43)	26	(23 - 29)
South Carolina	122	(110 - 136)	74	(64 - 84)
South Dakota	26	(24 - 29)	18	(16 - 20)
Tennessee	164	(146 - 184)	106	(93 - 122)
Texas	802	(757 - 850)	514	(477 - 553)
Utah	57	(48 - 66)	44	(36 - 52)
Vermont	24	(22 - 26)	16	(14 - 18)
Virginia	267	(243 - 293)	179	(159 - 200)
Washington	196	(177 - 217)	131	(116 - 148)
West Virginia	55	(50 - 61)	38	(33 - 42)
Wisconsin	189	(172 - 208)	123	(109 - 138)
Wyoming	18	(16 - 20)	13	(11 - 15)

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on at least 1 day in the past 30 days.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 13 Tobacco Product Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	68,933	(67,728 - 70,174)	2,551	(2,456 - 2,650)	13,732	(13,489 - 13,979)	52,649	(51,508 - 53,828)	66,382	(65,193 - 67,607)
Northeast	12,104	(11,657 - 12,565)	442	(413 - 473)	2,413	(2,342 - 2,486)	9,249	(8,816 - 9,695)	11,662	(11,222 - 12,117)
Midwest	16,405	(15,932 - 16,883)	616	(585 - 648)	3,319	(3,246 - 3,393)	12,470	(12,012 - 12,936)	15,789	(15,323 - 16,258)
South	26,743	(25,990 - 27,529)	957	(908 - 1,010)	5,073	(4,958 - 5,190)	20,713	(19,983 - 21,475)	25,786	(25,046 - 26,556)
West	13,680	(13,104 - 14,276)	536	(494 - 581)	2,927	(2,824 - 3,032)	10,217	(9,669 - 10,793)	13,144	(12,581 - 13,728)
Alabama	1,278	(1,168 - 1,392)	43	(36 - 51)	237	(218 - 257)	997	(895 - 1,105)	1,235	(1,126 - 1,347)
Alaska	185	(168 - 203)	7	(6 - 9)	38	(35 - 41)	139	(123 - 157)	178	(161 - 196)
Arizona	1,433	(1,277 - 1,600)	58	(47 - 71)	261	(235 - 288)	1,114	(971 - 1,270)	1,375	(1,222 - 1,538)
Arkansas	781	(712 - 853)	29	(24 - 34)	140	(129 - 151)	612	(549 - 680)	752	(685 - 822)
California	6,130	(5,710 - 6,569)	236	(204 - 272)	1,406	(1,320 - 1,495)	4,488	(4,093 - 4,911)	5,894	(5,482 - 6,328)
Colorado	1,144	(1,031 - 1,265)	47	(38 - 57)	243	(222 - 264)	855	(751 - 967)	1,098	(986 - 1,217)
Connecticut	761	(680 - 848)	29	(24 - 35)	156	(142 - 171)	575	(501 - 657)	732	(653 - 817)
Delaware	217	(197 - 238)	8	(6 - 9)	40	(36 - 44)	170	(151 - 190)	209	(190 - 230)
District of Columbia	140	(126 - 156)	2	(2 - 3)	31	(28 - 35)	107	(93 - 121)	138	(123 - 153)
Florida	4,122	(3,886 - 4,370)	118	(104 - 133)	696	(658 - 734)	3,309	(3,080 - 3,549)	4,005	(3,770 - 4,250)
Georgia	2,066	(1,861 - 2,285)	95	(78 - 115)	409	(370 - 450)	1,562	(1,374 - 1,766)	1,971	(1,770 - 2,187)
Hawaii	254	(225 - 285)	9	(7 - 11)	47	(42 - 52)	198	(171 - 228)	245	(217 - 276)
Idaho	346	(311 - 383)	15	(13 - 18)	67	(61 - 74)	263	(231 - 297)	330	(296 - 367)
Illinois	2,861	(2,705 - 3,022)	101	(90 - 114)	604	(576 - 632)	2,156	(2,008 - 2,310)	2,759	(2,606 - 2,919)
Indiana	1,594	(1,449 - 1,746)	57	(47 - 68)	319	(293 - 347)	1,218	(1,084 - 1,361)	1,537	(1,394 - 1,687)
Iowa	726	(659 - 797)	30	(25 - 36)	158	(145 - 172)	538	(476 - 604)	696	(630 - 765)
Kansas	713	(649 - 779)	29	(24 - 35)	153	(140 - 166)	531	(473 - 594)	684	(621 - 749)
Kentucky	1,339	(1,231 - 1,453)	50	(42 - 59)	232	(214 - 250)	1,057	(955 - 1,163)	1,289	(1,181 - 1,401)
Louisiana	1,134	(1,041 - 1,231)	40	(34 - 47)	214	(197 - 231)	881	(793 - 973)	1,094	(1,002 - 1,190)
Maine	330	(299 - 363)	11	(9 - 13)	59	(54 - 64)	259	(230 - 291)	319	(288 - 351)
Maryland	1,114	(993 - 1,244)	35	(28 - 43)	222	(200 - 245)	857	(745 - 979)	1,079	(960 - 1,208)
Massachusetts	1,285	(1,146 - 1,435)	54	(45 - 65)	295	(268 - 324)	935	(808 - 1,076)	1,230	(1,094 - 1,378)
Michigan	2,537	(2,409 - 2,669)	91	(82 - 101)	505	(484 - 526)	1,941	(1,819 - 2,068)	2,446	(2,320 - 2,576)
Minnesota	1,158	(1,046 - 1,275)	43	(36 - 52)	264	(244 - 285)	850	(747 - 961)	1,114	(1,004 - 1,230)
Mississippi	825	(758 - 894)	32	(27 - 38)	158	(146 - 170)	635	(573 - 700)	792	(727 - 861)
Missouri	1,674	(1,532 - 1,823)	72	(61 - 84)	308	(284 - 332)	1,295	(1,162 - 1,435)	1,602	(1,462 - 1,749)
Montana	264	(242 - 288)	12	(10 - 14)	54	(50 - 58)	198	(177 - 221)	252	(230 - 276)

(continued)

Table 13 Tobacco Product Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	423	(382 - 466)	17	(14 - 20)	89	(81 - 97)	317	(280 - 358)	406	(366 - 449)
Nevada	581	(515 - 652)	18	(14 - 24)	103	(93 - 115)	459	(397 - 525)	562	(497 - 632)
New Hampshire	308	(277 - 342)	14	(12 - 17)	65	(60 - 71)	229	(200 - 260)	294	(264 - 327)
New Jersey	1,922	(1,726 - 2,133)	72	(59 - 89)	360	(327 - 393)	1,490	(1,306 - 1,691)	1,850	(1,656 - 2,058)
New Mexico	434	(389 - 481)	20	(16 - 24)	96	(88 - 105)	317	(276 - 362)	414	(370 - 461)
New York	3,930	(3,688 - 4,184)	130	(114 - 148)	782	(738 - 827)	3,018	(2,786 - 3,263)	3,800	(3,560 - 4,052)
North Carolina	2,231	(2,021 - 2,452)	84	(69 - 101)	432	(394 - 471)	1,715	(1,524 - 1,920)	2,147	(1,940 - 2,364)
North Dakota	160	(145 - 175)	6	(5 - 7)	41	(38 - 44)	113	(100 - 127)	154	(139 - 169)
Ohio	3,067	(2,920 - 3,218)	118	(106 - 130)	564	(541 - 587)	2,385	(2,244 - 2,532)	2,949	(2,804 - 3,098)
Oklahoma	1,072	(978 - 1,170)	43	(36 - 51)	205	(188 - 222)	824	(737 - 915)	1,029	(937 - 1,125)
Oregon	868	(787 - 956)	29	(24 - 35)	170	(155 - 185)	670	(593 - 752)	839	(759 - 926)
Pennsylvania	3,165	(3,000 - 3,337)	116	(104 - 130)	606	(575 - 637)	2,443	(2,284 - 2,609)	3,049	(2,885 - 3,218)
Rhode Island	251	(227 - 277)	8	(7 - 10)	57	(52 - 61)	186	(164 - 210)	243	(219 - 268)
South Carolina	1,198	(1,086 - 1,316)	44	(37 - 52)	217	(199 - 235)	938	(833 - 1,049)	1,154	(1,044 - 1,270)
South Dakota	211	(191 - 230)	9	(8 - 11)	44	(40 - 48)	158	(141 - 176)	201	(183 - 221)
Tennessee	1,847	(1,689 - 2,013)	65	(55 - 77)	321	(296 - 347)	1,461	(1,313 - 1,615)	1,782	(1,625 - 1,944)
Texas	5,148	(4,854 - 5,452)	193	(171 - 218)	1,078	(1,024 - 1,133)	3,877	(3,601 - 4,167)	4,955	(4,665 - 5,255)
Utah	344	(297 - 396)	18	(14 - 24)	89	(77 - 103)	237	(196 - 285)	326	(280 - 377)
Vermont	151	(137 - 167)	6	(5 - 8)	33	(31 - 36)	112	(98 - 126)	145	(131 - 160)
Virginia	1,632	(1,470 - 1,806)	55	(45 - 67)	346	(315 - 378)	1,231	(1,084 - 1,392)	1,577	(1,418 - 1,748)
Washington	1,552	(1,400 - 1,713)	60	(50 - 73)	322	(295 - 349)	1,170	(1,031 - 1,320)	1,491	(1,343 - 1,650)
West Virginia	599	(553 - 647)	22	(19 - 26)	95	(89 - 102)	482	(437 - 528)	577	(531 - 624)
Wisconsin	1,283	(1,151 - 1,423)	43	(36 - 53)	272	(247 - 296)	968	(847 - 1,099)	1,239	(1,109 - 1,377)
Wyoming	146	(133 - 159)	6	(5 - 7)	30	(28 - 33)	110	(98 - 122)	140	(127 - 153)

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 14 Cigarette Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	57,569	(56,464 - 58,706)	1,990	(1,905 - 2,077)	11,594	(11,359 - 11,833)	43,985	(42,947 - 45,057)	55,579	(54,488 - 56,703)
Northeast	10,278	(9,863 - 10,708)	347	(321 - 374)	2,054	(1,976 - 2,133)	7,877	(7,462 - 8,309)	9,931	(9,520 - 10,358)
Midwest	13,587	(13,156 - 14,023)	476	(449 - 505)	2,801	(2,724 - 2,879)	10,309	(9,878 - 10,749)	13,110	(12,684 - 13,540)
South	22,255	(21,550 - 22,990)	745	(703 - 790)	4,249	(4,124 - 4,376)	17,261	(16,553 - 18,001)	21,509	(20,814 - 22,234)
West	11,450	(10,929 - 11,991)	421	(383 - 463)	2,491	(2,388 - 2,596)	8,538	(8,024 - 9,082)	11,029	(10,512 - 11,566)
Alabama	1,059	(957 - 1,167)	34	(27 - 41)	194	(176 - 214)	831	(735 - 934)	1,025	(924 - 1,132)
Alaska	156	(139 - 174)	5	(4 - 6)	33	(29 - 36)	119	(103 - 135)	151	(134 - 169)
Arizona	1,235	(1,086 - 1,399)	47	(37 - 59)	225	(200 - 252)	963	(824 - 1,118)	1,188	(1,040 - 1,350)
Arkansas	649	(582 - 721)	22	(18 - 27)	116	(105 - 128)	511	(447 - 580)	627	(560 - 699)
California	5,165	(4,783 - 5,567)	196	(167 - 229)	1,194	(1,113 - 1,278)	3,775	(3,409 - 4,173)	4,969	(4,590 - 5,371)
Colorado	927	(823 - 1,041)	33	(27 - 42)	201	(180 - 222)	693	(597 - 801)	894	(791 - 1,006)
Connecticut	642	(569 - 722)	23	(18 - 29)	135	(121 - 149)	485	(417 - 560)	619	(547 - 698)
Delaware	182	(163 - 203)	6	(4 - 7)	34	(31 - 38)	142	(124 - 162)	177	(157 - 198)
District of Columbia	122	(109 - 137)	2	(2 - 3)	26	(23 - 29)	95	(82 - 109)	120	(107 - 135)
Florida	3,488	(3,259 - 3,733)	94	(81 - 107)	587	(550 - 626)	2,808	(2,582 - 3,047)	3,395	(3,166 - 3,638)
Georgia	1,656	(1,461 - 1,870)	77	(62 - 96)	334	(297 - 374)	1,245	(1,066 - 1,445)	1,579	(1,386 - 1,791)
Hawaii	218	(192 - 247)	6	(5 - 8)	42	(37 - 47)	169	(145 - 197)	211	(185 - 240)
Idaho	286	(253 - 321)	12	(9 - 14)	57	(51 - 64)	217	(187 - 251)	274	(242 - 309)
Illinois	2,366	(2,217 - 2,521)	77	(67 - 88)	519	(489 - 549)	1,770	(1,628 - 1,921)	2,289	(2,142 - 2,443)
Indiana	1,349	(1,215 - 1,493)	43	(35 - 53)	273	(247 - 300)	1,033	(907 - 1,171)	1,306	(1,172 - 1,449)
Iowa	602	(540 - 668)	26	(21 - 31)	133	(121 - 147)	443	(386 - 505)	576	(515 - 641)
Kansas	553	(495 - 616)	22	(18 - 27)	122	(110 - 135)	409	(356 - 467)	531	(473 - 593)
Kentucky	1,124	(1,018 - 1,238)	39	(32 - 47)	195	(177 - 213)	890	(790 - 997)	1,085	(979 - 1,196)
Louisiana	996	(905 - 1,093)	33	(28 - 40)	186	(169 - 204)	777	(690 - 870)	963	(872 - 1,059)
Maine	284	(254 - 317)	8	(7 - 11)	51	(46 - 56)	224	(196 - 256)	276	(246 - 308)
Maryland	930	(821 - 1,049)	27	(21 - 34)	186	(164 - 210)	717	(616 - 830)	903	(795 - 1,022)
Massachusetts	1,069	(944 - 1,206)	40	(33 - 49)	250	(223 - 278)	779	(664 - 909)	1,029	(905 - 1,165)
Michigan	2,177	(2,057 - 2,300)	69	(60 - 78)	434	(413 - 455)	1,674	(1,558 - 1,796)	2,108	(1,989 - 2,232)
Minnesota	975	(870 - 1,087)	36	(29 - 45)	230	(209 - 252)	708	(613 - 814)	939	(835 - 1,049)
Mississippi	645	(583 - 710)	23	(19 - 28)	131	(119 - 143)	491	(433 - 553)	622	(560 - 687)
Missouri	1,340	(1,212 - 1,475)	55	(45 - 67)	251	(227 - 276)	1,033	(913 - 1,163)	1,285	(1,158 - 1,420)
Montana	202	(182 - 224)	9	(7 - 10)	42	(38 - 46)	152	(133 - 172)	194	(174 - 215)

(continued)

Table 14 Cigarette Use in the Past Month, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	346	(309 - 385)	13	(10 - 16)	78	(71 - 86)	254	(220 - 292)	333	(296 - 372)
Nevada	499	(439 - 565)	15	(12 - 19)	89	(79 - 100)	395	(339 - 458)	484	(425 - 550)
New Hampshire	250	(222 - 280)	11	(9 - 14)	54	(49 - 59)	185	(159 - 214)	239	(211 - 269)
New Jersey	1,675	(1,482 - 1,886)	59	(47 - 73)	307	(274 - 341)	1,309	(1,126 - 1,513)	1,616	(1,424 - 1,826)
New Mexico	353	(311 - 398)	15	(12 - 19)	81	(72 - 89)	257	(219 - 300)	338	(296 - 383)
New York	3,424	(3,204 - 3,655)	102	(88 - 117)	670	(627 - 715)	2,652	(2,438 - 2,880)	3,322	(3,104 - 3,552)
North Carolina	1,918	(1,713 - 2,140)	65	(53 - 80)	374	(336 - 414)	1,479	(1,290 - 1,686)	1,853	(1,649 - 2,072)
North Dakota	126	(112 - 140)	5	(4 - 6)	34	(30 - 37)	87	(75 - 101)	121	(107 - 135)
Ohio	2,549	(2,411 - 2,693)	89	(79 - 100)	466	(443 - 490)	1,994	(1,858 - 2,136)	2,460	(2,323 - 2,604)
Oklahoma	879	(790 - 974)	30	(24 - 37)	174	(158 - 190)	675	(592 - 765)	849	(760 - 943)
Oregon	723	(643 - 810)	22	(17 - 27)	149	(134 - 165)	552	(478 - 635)	701	(622 - 788)
Pennsylvania	2,595	(2,443 - 2,755)	92	(81 - 104)	512	(481 - 543)	1,991	(1,844 - 2,147)	2,503	(2,352 - 2,662)
Rhode Island	216	(193 - 240)	7	(5 - 8)	47	(42 - 52)	163	(141 - 186)	209	(187 - 234)
South Carolina	1,028	(923 - 1,141)	34	(28 - 42)	186	(169 - 204)	808	(709 - 915)	994	(890 - 1,105)
South Dakota	164	(146 - 183)	6	(5 - 8)	36	(32 - 39)	122	(106 - 140)	158	(140 - 176)
Tennessee	1,457	(1,304 - 1,621)	46	(37 - 56)	255	(230 - 281)	1,156	(1,013 - 1,311)	1,411	(1,258 - 1,573)
Texas	4,329	(4,051 - 4,617)	156	(135 - 179)	913	(860 - 968)	3,260	(2,995 - 3,542)	4,173	(3,898 - 4,460)
Utah	299	(255 - 346)	13	(10 - 18)	84	(72 - 98)	201	(164 - 245)	285	(243 - 332)
Vermont	122	(110 - 136)	5	(4 - 6)	28	(26 - 31)	89	(78 - 103)	118	(105 - 132)
Virginia	1,317	(1,167 - 1,481)	42	(34 - 52)	277	(247 - 309)	997	(860 - 1,151)	1,274	(1,126 - 1,437)
Washington	1,278	(1,141 - 1,427)	45	(36 - 55)	270	(244 - 297)	964	(837 - 1,104)	1,234	(1,097 - 1,381)
West Virginia	474	(431 - 521)	15	(13 - 19)	80	(73 - 86)	379	(337 - 424)	459	(415 - 505)
Wisconsin	1,041	(923 - 1,168)	36	(29 - 45)	224	(201 - 248)	781	(672 - 901)	1,005	(888 - 1,131)
Wyoming	109	(97 - 122)	5	(4 - 6)	25	(22 - 27)	80	(69 - 92)	105	(93 - 118)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 15 Perceptions of Great Risk of Smoking One or More Packs of Cigarettes per Day, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	182,378	(181,203 - 183,516)	16,213	(16,064 - 16,360)	22,466	(22,252 - 22,677)	143,700	(142,554 - 144,805)	166,166	(165,002 - 167,291)
Northeast	34,424	(34,002 - 34,834)	2,873	(2,824 - 2,920)	4,065	(3,989 - 4,140)	27,486	(27,089 - 27,870)	31,551	(31,138 - 31,953)
Midwest	37,539	(37,083 - 37,989)	3,474	(3,423 - 3,525)	4,578	(4,497 - 4,659)	29,486	(29,059 - 29,906)	34,064	(33,616 - 34,507)
South	66,684	(65,996 - 67,349)	5,904	(5,824 - 5,983)	8,191	(8,060 - 8,319)	52,590	(51,952 - 53,201)	60,780	(60,105 - 61,431)
West	43,732	(43,151 - 44,292)	3,962	(3,887 - 4,035)	5,632	(5,520 - 5,741)	34,138	(33,594 - 34,659)	39,770	(39,198 - 40,320)
Alabama	2,649	(2,535 - 2,758)	231	(218 - 243)	333	(315 - 351)	2,085	(1,977 - 2,187)	2,418	(2,306 - 2,525)
Alaska	373	(355 - 390)	39	(36 - 41)	51	(48 - 54)	284	(267 - 299)	334	(317 - 351)
Arizona	3,741	(3,571 - 3,901)	353	(333 - 373)	467	(440 - 492)	2,921	(2,759 - 3,070)	3,388	(3,220 - 3,545)
Arkansas	1,688	(1,619 - 1,753)	148	(140 - 156)	195	(184 - 205)	1,345	(1,279 - 1,406)	1,540	(1,471 - 1,604)
California	23,560	(23,096 - 24,002)	2,121	(2,058 - 2,182)	3,135	(3,047 - 3,220)	18,303	(17,862 - 18,717)	21,439	(20,980 - 21,872)
Colorado	2,981	(2,862 - 3,094)	257	(242 - 270)	365	(346 - 383)	2,359	(2,246 - 2,464)	2,725	(2,607 - 2,834)
Connecticut	2,213	(2,132 - 2,290)	194	(184 - 204)	248	(235 - 261)	1,771	(1,694 - 1,841)	2,019	(1,939 - 2,093)
Delaware	556	(536 - 575)	45	(42 - 47)	66	(62 - 69)	445	(427 - 462)	511	(492 - 529)
District of Columbia	397	(383 - 411)	21	(20 - 22)	64	(61 - 67)	313	(299 - 325)	377	(362 - 390)
Florida	12,032	(11,796 - 12,258)	921	(895 - 945)	1,332	(1,293 - 1,370)	9,779	(9,550 - 9,996)	11,111	(10,877 - 11,336)
Georgia	5,528	(5,300 - 5,744)	522	(492 - 550)	711	(674 - 746)	4,296	(4,085 - 4,493)	5,006	(4,784 - 5,216)
Hawaii	756	(723 - 787)	61	(57 - 65)	88	(83 - 93)	606	(575 - 636)	695	(662 - 726)
Idaho	856	(818 - 892)	91	(86 - 95)	110	(103 - 116)	655	(620 - 688)	765	(728 - 800)
Illinois	7,501	(7,333 - 7,664)	692	(671 - 713)	910	(881 - 938)	5,899	(5,739 - 6,052)	6,809	(6,644 - 6,968)
Indiana	3,597	(3,431 - 3,754)	349	(330 - 367)	449	(424 - 473)	2,799	(2,641 - 2,947)	3,248	(3,085 - 3,402)
Iowa	1,700	(1,629 - 1,770)	149	(141 - 157)	218	(206 - 230)	1,333	(1,265 - 1,397)	1,551	(1,481 - 1,619)
Kansas	1,502	(1,433 - 1,569)	139	(130 - 147)	199	(187 - 211)	1,164	(1,099 - 1,226)	1,363	(1,296 - 1,428)
Kentucky	2,267	(2,155 - 2,375)	215	(202 - 227)	272	(255 - 288)	1,780	(1,674 - 1,882)	2,052	(1,942 - 2,159)
Louisiana	2,638	(2,542 - 2,730)	235	(223 - 246)	347	(330 - 363)	2,057	(1,965 - 2,141)	2,404	(2,308 - 2,493)
Maine	781	(748 - 813)	63	(59 - 66)	79	(74 - 83)	640	(607 - 670)	718	(685 - 750)
Maryland	3,575	(3,445 - 3,697)	301	(285 - 316)	428	(406 - 448)	2,847	(2,724 - 2,959)	3,275	(3,146 - 3,393)
Massachusetts	4,221	(4,068 - 4,362)	328	(312 - 343)	520	(495 - 544)	3,373	(3,228 - 3,504)	3,893	(3,742 - 4,032)
Michigan	5,659	(5,527 - 5,787)	532	(516 - 547)	690	(669 - 711)	4,437	(4,311 - 4,559)	5,127	(4,997 - 5,253)
Minnesota	2,988	(2,859 - 3,111)	282	(267 - 296)	347	(324 - 368)	2,359	(2,238 - 2,473)	2,706	(2,579 - 2,827)
Mississippi	1,648	(1,584 - 1,710)	154	(146 - 163)	220	(209 - 230)	1,274	(1,212 - 1,331)	1,494	(1,430 - 1,554)
Missouri	3,226	(3,078 - 3,368)	286	(269 - 302)	396	(373 - 418)	2,544	(2,404 - 2,677)	2,940	(2,794 - 3,079)
Montana	563	(541 - 584)	47	(45 - 50)	70	(66 - 73)	445	(425 - 465)	515	(493 - 536)

(continued)

Table 15 Perceptions of Great Risk of Smoking One or More Packs of Cigarettes per Day, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	1,017	(974 - 1,059)	91	(86 - 96)	134	(127 - 141)	792	(751 - 830)	926	(884 - 967)
Nevada	1,548	(1,478 - 1,613)	143	(134 - 151)	182	(171 - 192)	1,223	(1,159 - 1,283)	1,405	(1,338 - 1,469)
New Hampshire	803	(771 - 833)	66	(62 - 69)	91	(86 - 95)	646	(616 - 674)	737	(706 - 766)
New Jersey	5,519	(5,316 - 5,708)	494	(470 - 517)	596	(564 - 626)	4,429	(4,234 - 4,605)	5,025	(4,824 - 5,210)
New Mexico	1,207	(1,158 - 1,253)	108	(102 - 113)	149	(141 - 157)	950	(904 - 992)	1,099	(1,051 - 1,144)
New York	12,597	(12,349 - 12,833)	1,037	(1,007 - 1,065)	1,552	(1,508 - 1,594)	10,008	(9,771 - 10,231)	11,560	(11,316 - 11,792)
North Carolina	5,353	(5,139 - 5,557)	495	(470 - 519)	632	(596 - 667)	4,225	(4,021 - 4,416)	4,857	(4,647 - 5,058)
North Dakota	378	(362 - 393)	32	(30 - 34)	57	(54 - 61)	289	(274 - 303)	346	(330 - 361)
Ohio	6,242	(6,082 - 6,398)	589	(570 - 607)	735	(712 - 757)	4,919	(4,764 - 5,068)	5,653	(5,496 - 5,807)
Oklahoma	2,049	(1,960 - 2,133)	177	(165 - 188)	261	(246 - 277)	1,611	(1,527 - 1,689)	1,872	(1,785 - 1,954)
Oregon	2,245	(2,151 - 2,333)	191	(181 - 200)	260	(246 - 274)	1,793	(1,704 - 1,876)	2,054	(1,961 - 2,141)
Pennsylvania	7,250	(7,081 - 7,414)	609	(589 - 628)	843	(813 - 873)	5,798	(5,635 - 5,955)	6,641	(6,474 - 6,803)
Rhode Island	658	(633 - 682)	53	(50 - 56)	90	(86 - 94)	516	(491 - 538)	606	(581 - 629)
South Carolina	2,605	(2,496 - 2,710)	218	(206 - 231)	315	(298 - 331)	2,072	(1,968 - 2,170)	2,387	(2,279 - 2,489)
South Dakota	449	(428 - 469)	41	(39 - 43)	57	(54 - 61)	350	(331 - 368)	407	(387 - 427)
Tennessee	3,471	(3,310 - 3,626)	310	(294 - 326)	400	(376 - 422)	2,760	(2,607 - 2,906)	3,160	(3,001 - 3,313)
Texas	14,446	(14,141 - 14,741)	1,416	(1,373 - 1,457)	1,905	(1,848 - 1,961)	11,125	(10,836 - 11,402)	13,030	(12,730 - 13,319)
Utah	1,605	(1,548 - 1,658)	193	(184 - 201)	250	(237 - 263)	1,161	(1,109 - 1,209)	1,412	(1,356 - 1,463)
Vermont	382	(366 - 397)	29	(27 - 31)	48	(45 - 50)	305	(291 - 319)	353	(337 - 367)
Virginia	4,723	(4,531 - 4,904)	413	(393 - 432)	597	(567 - 625)	3,714	(3,531 - 3,882)	4,310	(4,121 - 4,487)
Washington	3,982	(3,820 - 4,135)	331	(313 - 349)	464	(438 - 488)	3,187	(3,032 - 3,330)	3,650	(3,490 - 3,801)
West Virginia	1,059	(1,014 - 1,102)	83	(78 - 87)	114	(107 - 121)	862	(819 - 902)	976	(932 - 1,018)
Wisconsin	3,280	(3,142 - 3,412)	293	(277 - 308)	386	(363 - 408)	2,601	(2,470 - 2,724)	2,987	(2,851 - 3,117)
Wyoming	318	(305 - 330)	28	(26 - 29)	41	(38 - 43)	250	(237 - 261)	290	(277 - 302)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 16 Alcohol Dependence or Abuse in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	17,320	(16,767 - 17,888)	1,028	(965 - 1,096)	5,139	(4,973 - 5,310)	11,152	(10,631 - 11,697)	16,291	(15,744 - 16,856)
Northeast	3,268	(3,066 - 3,485)	180	(163 - 199)	957	(902 - 1,015)	2,131	(1,945 - 2,332)	3,088	(2,889 - 3,302)
Midwest	3,897	(3,696 - 4,107)	217	(200 - 235)	1,177	(1,120 - 1,236)	2,503	(2,317 - 2,704)	3,680	(3,481 - 3,888)
South	5,692	(5,391 - 6,007)	342	(315 - 372)	1,689	(1,601 - 1,780)	3,661	(3,379 - 3,964)	5,349	(5,052 - 5,662)
West	4,463	(4,164 - 4,780)	289	(258 - 323)	1,316	(1,236 - 1,401)	2,858	(2,588 - 3,155)	4,174	(3,880 - 4,488)
Alabama	212	(172 - 261)	14	(10 - 18)	65	(54 - 78)	133	(100 - 178)	198	(159 - 247)
Alaska	44	(36 - 54)	3	(2 - 4)	13	(11 - 15)	29	(22 - 37)	42	(34 - 51)
Arizona	436	(355 - 534)	25	(19 - 33)	114	(96 - 135)	297	(227 - 388)	411	(331 - 507)
Arkansas	127	(101 - 159)	9	(7 - 12)	36	(30 - 44)	81	(59 - 111)	117	(92 - 149)
California	2,275	(2,042 - 2,532)	164	(138 - 194)	694	(629 - 764)	1,418	(1,211 - 1,657)	2,112	(1,882 - 2,367)
Colorado	356	(296 - 426)	18	(14 - 24)	107	(92 - 123)	231	(179 - 296)	338	(280 - 407)
Connecticut	218	(178 - 265)	12	(9 - 16)	64	(55 - 75)	141	(107 - 185)	206	(167 - 252)
Delaware	46	(37 - 57)	3	(2 - 3)	14	(12 - 16)	30	(22 - 39)	44	(35 - 54)
District of Columbia	57	(47 - 68)	1	(1 - 2)	16	(14 - 19)	39	(31 - 49)	56	(46 - 66)
Florida	936	(828 - 1,059)	52	(44 - 62)	254	(228 - 283)	630	(532 - 744)	884	(777 - 1,006)
Georgia	446	(359 - 552)	32	(24 - 41)	137	(114 - 163)	278	(205 - 374)	414	(330 - 519)
Hawaii	85	(69 - 105)	4	(3 - 6)	23	(20 - 27)	58	(44 - 76)	81	(65 - 100)
Idaho	94	(77 - 114)	6	(5 - 8)	27	(23 - 32)	60	(46 - 79)	88	(71 - 108)
Illinois	734	(657 - 818)	41	(35 - 49)	235	(215 - 257)	457	(389 - 537)	692	(617 - 776)
Indiana	370	(300 - 454)	20	(15 - 26)	114	(96 - 135)	235	(177 - 312)	350	(282 - 433)
Iowa	196	(163 - 234)	10	(8 - 13)	66	(56 - 77)	120	(93 - 155)	186	(154 - 224)
Kansas	166	(137 - 201)	10	(8 - 13)	53	(44 - 62)	103	(79 - 134)	155	(127 - 190)
Kentucky	172	(136 - 216)	12	(9 - 16)	52	(42 - 63)	108	(78 - 148)	159	(125 - 203)
Louisiana	233	(192 - 281)	12	(9 - 15)	71	(61 - 83)	150	(115 - 195)	221	(181 - 269)
Maine	61	(49 - 76)	4	(3 - 5)	19	(16 - 23)	38	(28 - 52)	57	(46 - 72)
Maryland	246	(196 - 308)	14	(10 - 18)	78	(65 - 95)	154	(112 - 211)	232	(184 - 294)
Massachusetts	471	(393 - 563)	24	(19 - 31)	145	(125 - 168)	301	(234 - 386)	447	(370 - 538)
Michigan	593	(531 - 663)	33	(28 - 38)	164	(149 - 179)	397	(339 - 464)	561	(499 - 630)
Minnesota	339	(281 - 408)	18	(14 - 23)	104	(89 - 121)	218	(168 - 281)	322	(264 - 390)
Mississippi	133	(108 - 164)	9	(7 - 12)	40	(33 - 48)	85	(63 - 113)	125	(100 - 155)
Missouri	276	(222 - 341)	20	(15 - 26)	89	(75 - 106)	166	(121 - 227)	256	(203 - 321)
Montana	70	(59 - 83)	4	(3 - 5)	21	(18 - 25)	45	(35 - 57)	66	(55 - 79)

(continued)

Table 16 Alcohol Dependence or Abuse in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	94	(77 - 115)	5	(4 - 7)	33	(28 - 38)	56	(42 - 75)	89	(72 - 109)
Nevada	197	(161 - 240)	10	(7 - 13)	48	(41 - 56)	139	(106 - 180)	187	(151 - 230)
New Hampshire	76	(62 - 93)	5	(4 - 6)	24	(21 - 28)	47	(35 - 63)	71	(57 - 88)
New Jersey	476	(389 - 582)	29	(23 - 38)	129	(108 - 153)	318	(243 - 413)	447	(361 - 551)
New Mexico	117	(96 - 142)	8	(6 - 10)	36	(30 - 42)	73	(56 - 96)	109	(89 - 134)
New York	1,091	(970 - 1,227)	61	(51 - 72)	320	(288 - 355)	710	(601 - 837)	1,030	(911 - 1,164)
North Carolina	431	(347 - 534)	30	(23 - 38)	131	(109 - 158)	271	(201 - 364)	402	(319 - 503)
North Dakota	49	(41 - 57)	2	(2 - 3)	19	(17 - 22)	28	(21 - 35)	47	(39 - 55)
Ohio	696	(623 - 778)	37	(31 - 43)	188	(172 - 206)	472	(404 - 549)	660	(587 - 741)
Oklahoma	218	(177 - 267)	11	(9 - 15)	59	(49 - 71)	147	(112 - 192)	206	(166 - 255)
Oregon	252	(207 - 306)	12	(9 - 15)	65	(55 - 77)	175	(135 - 225)	240	(196 - 294)
Pennsylvania	755	(676 - 843)	40	(34 - 47)	217	(195 - 240)	499	(427 - 582)	715	(637 - 803)
Rhode Island	82	(68 - 98)	3	(3 - 4)	25	(21 - 29)	53	(41 - 68)	78	(65 - 94)
South Carolina	249	(203 - 304)	14	(11 - 18)	65	(54 - 77)	170	(130 - 221)	235	(190 - 289)
South Dakota	59	(49 - 71)	4	(3 - 5)	17	(14 - 20)	39	(31 - 49)	56	(46 - 67)
Tennessee	323	(263 - 395)	18	(14 - 24)	91	(76 - 108)	214	(163 - 279)	304	(245 - 376)
Texas	1,328	(1,184 - 1,487)	82	(68 - 97)	408	(368 - 450)	838	(710 - 988)	1,246	(1,105 - 1,404)
Utah	100	(79 - 126)	10	(7 - 13)	38	(30 - 47)	53	(37 - 75)	91	(70 - 116)
Vermont	39	(32 - 47)	2	(2 - 3)	14	(12 - 16)	23	(17 - 31)	37	(30 - 45)
Virginia	454	(375 - 548)	26	(20 - 33)	144	(123 - 167)	284	(218 - 369)	428	(351 - 521)
Washington	403	(333 - 486)	23	(18 - 30)	120	(103 - 139)	260	(200 - 336)	380	(311 - 462)
West Virginia	83	(67 - 102)	5	(4 - 7)	28	(24 - 33)	50	(37 - 68)	78	(62 - 97)
Wisconsin	324	(263 - 398)	16	(12 - 22)	95	(79 - 114)	212	(161 - 280)	308	(248 - 381)
Wyoming	33	(28 - 40)	2	(1 - 2)	10	(9 - 12)	21	(16 - 27)	31	(26 - 38)

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 17 Alcohol Dependence in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	8,049	(7,662 - 8,453)	383	(347 - 424)	2,141	(2,034 - 2,252)	5,525	(5,158 - 5,916)	7,665	(7,282 - 8,067)
Northeast	1,484	(1,354 - 1,627)	65	(56 - 76)	376	(343 - 413)	1,043	(923 - 1,178)	1,419	(1,291 - 1,560)
Midwest	1,792	(1,665 - 1,927)	83	(73 - 94)	494	(460 - 529)	1,215	(1,098 - 1,344)	1,709	(1,583 - 1,843)
South	2,614	(2,409 - 2,837)	127	(110 - 145)	689	(636 - 746)	1,799	(1,610 - 2,010)	2,488	(2,285 - 2,708)
West	2,159	(1,950 - 2,387)	109	(92 - 129)	582	(528 - 642)	1,468	(1,277 - 1,686)	2,050	(1,844 - 2,276)
Alabama	104	(80 - 135)	5	(3 - 7)	27	(20 - 36)	72	(51 - 101)	99	(75 - 130)
Alaska	21	(16 - 26)	1	(1 - 1)	6	(4 - 7)	14	(10 - 19)	20	(15 - 25)
Arizona	205	(155 - 271)	9	(6 - 13)	50	(38 - 65)	146	(102 - 208)	196	(147 - 261)
Arkansas	66	(50 - 87)	4	(3 - 6)	18	(13 - 23)	44	(31 - 63)	62	(47 - 83)
California	1,131	(972 - 1,314)	61	(48 - 78)	311	(267 - 360)	759	(615 - 936)	1,070	(913 - 1,252)
Colorado	170	(132 - 218)	7	(5 - 10)	47	(37 - 60)	115	(83 - 159)	162	(125 - 210)
Connecticut	90	(68 - 117)	4	(3 - 6)	23	(18 - 30)	62	(44 - 87)	85	(65 - 112)
Delaware	21	(15 - 27)	1	(1 - 2)	6	(4 - 8)	14	(9 - 20)	19	(14 - 26)
District of Columbia	31	(24 - 39)	1	(0 - 1)	7	(6 - 9)	23	(17 - 31)	30	(24 - 39)
Florida	450	(378 - 536)	20	(16 - 25)	103	(87 - 122)	327	(261 - 408)	430	(359 - 515)
Georgia	211	(158 - 279)	11	(8 - 16)	53	(39 - 71)	146	(102 - 210)	199	(148 - 267)
Hawaii	40	(31 - 53)	1	(1 - 2)	10	(8 - 13)	29	(20 - 41)	39	(29 - 51)
Idaho	45	(35 - 59)	2	(2 - 3)	12	(10 - 16)	31	(22 - 43)	43	(33 - 57)
Illinois	366	(315 - 424)	17	(13 - 21)	110	(95 - 126)	239	(193 - 296)	349	(299 - 407)
Indiana	164	(128 - 210)	8	(6 - 11)	51	(40 - 65)	105	(75 - 147)	156	(121 - 201)
Iowa	90	(71 - 114)	4	(3 - 6)	29	(23 - 37)	57	(41 - 78)	86	(67 - 109)
Kansas	71	(55 - 92)	4	(3 - 6)	23	(17 - 29)	45	(32 - 63)	67	(52 - 87)
Kentucky	87	(64 - 117)	5	(3 - 7)	23	(17 - 31)	59	(40 - 87)	82	(60 - 112)
Louisiana	106	(83 - 136)	5	(4 - 7)	28	(22 - 35)	74	(53 - 102)	101	(78 - 131)
Maine	29	(21 - 39)	1	(1 - 2)	7	(5 - 9)	21	(14 - 30)	28	(20 - 38)
Maryland	115	(83 - 158)	5	(3 - 8)	31	(23 - 42)	79	(52 - 120)	110	(79 - 152)
Massachusetts	213	(167 - 272)	9	(6 - 13)	64	(50 - 81)	140	(101 - 194)	204	(159 - 262)
Michigan	300	(256 - 350)	12	(10 - 15)	66	(57 - 77)	221	(180 - 270)	287	(244 - 338)
Minnesota	157	(122 - 203)	7	(5 - 10)	45	(35 - 57)	105	(75 - 148)	150	(115 - 195)
Mississippi	66	(51 - 87)	3	(2 - 5)	19	(14 - 24)	45	(32 - 63)	63	(48 - 83)
Missouri	131	(100 - 170)	7	(5 - 10)	37	(28 - 48)	86	(61 - 123)	123	(93 - 163)
Montana	27	(22 - 35)	1	(1 - 2)	8	(7 - 11)	18	(13 - 25)	26	(20 - 33)

(continued)

Table 17 Alcohol Dependence in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	37	(27 - 49)	2	(1 - 3)	11	(9 - 15)	23	(16 - 35)	35	(25 - 47)
Nevada	82	(63 - 108)	4	(3 - 6)	21	(16 - 28)	57	(40 - 81)	79	(59 - 104)
New Hampshire	33	(25 - 43)	2	(1 - 2)	8	(6 - 10)	23	(17 - 33)	31	(24 - 41)
New Jersey	209	(158 - 275)	11	(7 - 15)	46	(34 - 61)	152	(107 - 214)	198	(148 - 264)
New Mexico	58	(46 - 74)	3	(2 - 5)	18	(14 - 23)	38	(27 - 52)	55	(43 - 71)
New York	526	(446 - 620)	22	(18 - 29)	135	(115 - 159)	369	(296 - 459)	504	(424 - 597)
North Carolina	184	(137 - 247)	10	(7 - 14)	47	(35 - 64)	127	(87 - 185)	174	(128 - 237)
North Dakota	23	(18 - 28)	1	(1 - 1)	8	(6 - 10)	14	(10 - 19)	22	(17 - 28)
Ohio	297	(254 - 347)	13	(10 - 16)	72	(62 - 84)	212	(173 - 259)	284	(242 - 334)
Oklahoma	80	(61 - 106)	4	(3 - 6)	25	(19 - 32)	52	(36 - 75)	76	(57 - 101)
Oregon	120	(93 - 154)	5	(3 - 7)	27	(21 - 35)	88	(64 - 120)	115	(89 - 149)
Pennsylvania	336	(284 - 397)	13	(10 - 17)	79	(66 - 95)	243	(196 - 301)	323	(271 - 383)
Rhode Island	32	(24 - 41)	1	(1 - 2)	9	(7 - 12)	21	(15 - 29)	30	(23 - 39)
South Carolina	120	(91 - 159)	5	(3 - 7)	24	(18 - 32)	91	(65 - 128)	115	(86 - 154)
South Dakota	25	(19 - 32)	1	(1 - 2)	7	(5 - 9)	16	(12 - 23)	23	(18 - 30)
Tennessee	147	(111 - 195)	7	(5 - 10)	36	(27 - 47)	104	(73 - 150)	140	(104 - 187)
Texas	586	(496 - 691)	30	(24 - 39)	171	(145 - 201)	384	(306 - 482)	556	(468 - 659)
Utah	54	(41 - 71)	4	(3 - 5)	19	(14 - 25)	32	(22 - 46)	51	(38 - 67)
Vermont	17	(13 - 22)	1	(1 - 1)	5	(4 - 6)	11	(8 - 16)	16	(13 - 21)
Virginia	199	(154 - 257)	9	(7 - 13)	60	(47 - 77)	130	(91 - 184)	190	(146 - 248)
Washington	190	(148 - 243)	9	(7 - 13)	49	(39 - 63)	131	(95 - 181)	180	(139 - 233)
West Virginia	41	(31 - 54)	2	(1 - 3)	12	(9 - 15)	27	(19 - 39)	39	(29 - 52)
Wisconsin	131	(97 - 176)	6	(4 - 9)	34	(25 - 45)	91	(62 - 133)	125	(92 - 169)
Wyoming	15	(12 - 19)	1	(0 - 1)	4	(3 - 5)	10	(7 - 14)	14	(11 - 19)

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 18 *Illicit Drug Dependence or Abuse in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs*

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	6,837	(6,528 - 7,162)	1,155	(1,093 - 1,221)	2,625	(2,510 - 2,746)	3,057	(2,793 - 3,344)	5,682	(5,385 - 5,995)
Northeast	1,253	(1,161 - 1,354)	195	(176 - 216)	489	(450 - 532)	568	(493 - 655)	1,058	(970 - 1,154)
Midwest	1,417	(1,328 - 1,511)	237	(218 - 257)	540	(505 - 576)	641	(566 - 724)	1,180	(1,096 - 1,270)
South	2,425	(2,270 - 2,590)	396	(365 - 429)	893	(834 - 957)	1,135	(1,004 - 1,283)	2,029	(1,880 - 2,189)
West	1,743	(1,604 - 1,892)	327	(293 - 365)	703	(642 - 769)	712	(603 - 842)	1,415	(1,284 - 1,560)
Alabama	94	(76 - 117)	15	(12 - 20)	35	(27 - 45)	44	(31 - 63)	79	(62 - 101)
Alaska	15	(12 - 19)	3	(2 - 4)	6	(4 - 8)	6	(4 - 9)	12	(9 - 15)
Arizona	165	(132 - 206)	32	(24 - 43)	57	(44 - 73)	76	(52 - 111)	133	(102 - 171)
Arkansas	61	(50 - 76)	10	(8 - 14)	21	(17 - 27)	29	(21 - 41)	51	(40 - 64)
California	926	(820 - 1,046)	180	(153 - 211)	386	(337 - 442)	360	(279 - 462)	746	(646 - 862)
Colorado	124	(103 - 149)	21	(16 - 28)	57	(46 - 70)	47	(33 - 67)	103	(84 - 127)
Connecticut	80	(65 - 100)	13	(10 - 18)	29	(23 - 38)	38	(27 - 54)	67	(53 - 86)
Delaware	21	(17 - 25)	3	(2 - 4)	8	(6 - 10)	10	(7 - 14)	17	(14 - 22)
District of Columbia	18	(14 - 23)	1	(1 - 2)	6	(5 - 8)	10	(7 - 15)	17	(13 - 21)
Florida	411	(359 - 472)	59	(50 - 70)	155	(134 - 179)	197	(155 - 252)	353	(303 - 411)
Georgia	189	(151 - 237)	33	(25 - 45)	68	(52 - 88)	88	(61 - 126)	156	(121 - 201)
Hawaii	23	(18 - 28)	4	(3 - 6)	8	(6 - 10)	11	(7 - 16)	18	(14 - 24)
Idaho	33	(27 - 41)	6	(5 - 8)	13	(10 - 16)	15	(10 - 21)	27	(22 - 35)
Illinois	274	(240 - 313)	43	(36 - 52)	115	(100 - 132)	116	(89 - 150)	231	(199 - 268)
Indiana	130	(104 - 162)	20	(15 - 27)	51	(39 - 65)	59	(41 - 86)	110	(85 - 141)
Iowa	56	(45 - 70)	10	(8 - 14)	22	(17 - 29)	24	(17 - 35)	46	(36 - 59)
Kansas	59	(49 - 72)	10	(8 - 14)	24	(19 - 31)	25	(17 - 35)	49	(39 - 61)
Kentucky	91	(74 - 111)	16	(12 - 21)	33	(25 - 42)	42	(30 - 58)	75	(59 - 94)
Louisiana	88	(72 - 109)	14	(11 - 19)	31	(24 - 40)	43	(30 - 61)	74	(58 - 94)
Maine	25	(20 - 31)	4	(3 - 5)	11	(9 - 14)	11	(7 - 15)	22	(17 - 27)
Maryland	125	(102 - 154)	19	(14 - 25)	49	(39 - 63)	57	(40 - 80)	106	(84 - 134)
Massachusetts	155	(126 - 190)	30	(23 - 39)	61	(47 - 77)	64	(44 - 94)	125	(98 - 158)
Michigan	208	(183 - 236)	39	(33 - 46)	77	(67 - 88)	92	(72 - 117)	169	(145 - 196)
Minnesota	109	(90 - 133)	20	(15 - 26)	44	(34 - 56)	46	(32 - 64)	89	(71 - 112)
Mississippi	66	(54 - 81)	10	(7 - 13)	25	(20 - 32)	32	(23 - 44)	57	(45 - 71)
Missouri	139	(114 - 169)	22	(17 - 29)	56	(45 - 70)	61	(44 - 85)	117	(94 - 145)
Montana	23	(19 - 28)	4	(3 - 5)	9	(7 - 11)	10	(7 - 13)	19	(15 - 23)

(continued)

Table 18 Illicit Drug Dependence or Abuse in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Nebraska	34	(27 - 42)	6	(4 - 8)	12	(9 - 16)	16	(11 - 24)	28	(22 - 36)
Nevada	58	(46 - 74)	10	(7 - 14)	19	(14 - 25)	29	(20 - 42)	48	(37 - 63)
New Hampshire	33	(27 - 40)	7	(5 - 9)	13	(10 - 16)	13	(9 - 19)	26	(21 - 33)
New Jersey	191	(156 - 235)	34	(25 - 45)	71	(55 - 91)	86	(61 - 121)	158	(125 - 199)
New Mexico	52	(42 - 64)	12	(9 - 15)	20	(16 - 26)	20	(13 - 30)	40	(31 - 51)
New York	439	(384 - 503)	61	(51 - 73)	177	(154 - 204)	201	(158 - 256)	378	(325 - 440)
North Carolina	210	(171 - 257)	36	(27 - 47)	79	(62 - 100)	95	(67 - 135)	174	(138 - 219)
North Dakota	11	(9 - 14)	2	(1 - 2)	5	(3 - 6)	5	(3 - 7)	9	(7 - 12)
Ohio	267	(235 - 304)	41	(35 - 48)	91	(80 - 104)	135	(107 - 169)	226	(195 - 262)
Oklahoma	76	(61 - 96)	12	(9 - 17)	27	(21 - 36)	37	(26 - 53)	64	(50 - 83)
Oregon	99	(80 - 122)	16	(12 - 21)	38	(30 - 48)	45	(32 - 63)	83	(65 - 104)
Pennsylvania	285	(250 - 326)	41	(34 - 49)	107	(91 - 125)	138	(109 - 175)	244	(210 - 284)
Rhode Island	26	(21 - 31)	4	(3 - 5)	11	(9 - 14)	11	(8 - 16)	22	(18 - 28)
South Carolina	100	(81 - 123)	17	(13 - 22)	36	(28 - 45)	48	(34 - 67)	83	(65 - 106)
South Dakota	15	(12 - 19)	3	(2 - 4)	5	(4 - 7)	7	(5 - 10)	12	(9 - 16)
Tennessee	153	(124 - 188)	23	(18 - 30)	58	(46 - 73)	72	(51 - 103)	130	(103 - 164)
Texas	512	(446 - 589)	95	(80 - 114)	179	(152 - 211)	238	(185 - 305)	417	(354 - 491)
Utah	61	(49 - 74)	12	(9 - 16)	24	(18 - 30)	25	(18 - 35)	49	(39 - 61)
Vermont	18	(15 - 22)	2	(2 - 3)	9	(7 - 11)	7	(5 - 10)	16	(13 - 19)
Virginia	166	(135 - 205)	27	(20 - 36)	65	(50 - 84)	75	(53 - 105)	139	(110 - 176)
Washington	153	(126 - 186)	24	(19 - 32)	64	(51 - 80)	65	(46 - 92)	129	(103 - 160)
West Virginia	41	(33 - 50)	5	(3 - 7)	18	(14 - 22)	19	(13 - 27)	36	(29 - 45)
Wisconsin	114	(91 - 142)	19	(14 - 26)	39	(30 - 51)	55	(39 - 80)	94	(73 - 121)
Wyoming	10	(8 - 12)	2	(1 - 2)	4	(3 - 5)	5	(3 - 7)	8	(6 - 11)

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 19 Illicit Drug Dependence in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	4,707	(4,446 - 4,983)	625	(578 - 676)	1,831	(1,733 - 1,935)	2,250	(2,026 - 2,498)	4,081	(3,829 - 4,350)
Northeast	890	(808 - 980)	105	(92 - 118)	345	(313 - 381)	440	(371 - 522)	785	(706 - 874)
Midwest	976	(897 - 1,061)	129	(116 - 143)	372	(343 - 403)	475	(407 - 553)	847	(771 - 929)
South	1,637	(1,507 - 1,779)	216	(194 - 239)	616	(568 - 669)	805	(693 - 936)	1,422	(1,297 - 1,559)
West	1,204	(1,077 - 1,344)	176	(152 - 204)	498	(447 - 555)	530	(428 - 656)	1,028	(906 - 1,165)
Alabama	63	(49 - 82)	8	(6 - 11)	25	(18 - 33)	31	(20 - 46)	55	(41 - 73)
Alaska	11	(8 - 14)	2	(1 - 2)	4	(3 - 5)	5	(3 - 7)	9	(7 - 12)
Arizona	117	(89 - 154)	19	(14 - 26)	43	(32 - 58)	55	(36 - 86)	99	(72 - 133)
Arkansas	42	(33 - 55)	6	(4 - 8)	14	(11 - 19)	22	(15 - 33)	36	(28 - 48)
California	641	(546 - 750)	95	(77 - 117)	278	(236 - 327)	267	(195 - 366)	545	(455 - 653)
Colorado	76	(59 - 97)	11	(8 - 15)	33	(25 - 43)	32	(21 - 49)	65	(50 - 85)
Connecticut	59	(46 - 76)	7	(5 - 10)	21	(16 - 28)	30	(20 - 45)	52	(39 - 68)
Delaware	15	(12 - 19)	2	(1 - 2)	6	(5 - 8)	7	(5 - 10)	13	(10 - 17)
District of Columbia	13	(10 - 17)	1	(1 - 1)	4	(3 - 6)	8	(6 - 12)	12	(9 - 17)
Florida	260	(220 - 307)	31	(25 - 38)	102	(85 - 121)	127	(96 - 167)	229	(191 - 275)
Georgia	121	(92 - 160)	18	(13 - 25)	44	(31 - 60)	59	(38 - 92)	103	(75 - 140)
Hawaii	15	(11 - 21)	2	(2 - 3)	5	(4 - 7)	8	(5 - 12)	13	(9 - 18)
Idaho	23	(18 - 29)	4	(3 - 5)	9	(6 - 12)	11	(7 - 16)	20	(15 - 26)
Illinois	192	(163 - 226)	26	(20 - 33)	79	(67 - 94)	87	(64 - 117)	166	(139 - 199)
Indiana	89	(69 - 115)	12	(9 - 16)	34	(25 - 46)	43	(29 - 64)	77	(58 - 102)
Iowa	38	(29 - 50)	6	(4 - 8)	16	(11 - 21)	17	(11 - 26)	32	(24 - 44)
Kansas	39	(30 - 50)	6	(4 - 8)	15	(11 - 21)	18	(12 - 27)	33	(25 - 44)
Kentucky	67	(52 - 86)	8	(6 - 11)	27	(20 - 35)	32	(22 - 47)	59	(45 - 77)
Louisiana	62	(48 - 80)	8	(6 - 11)	24	(18 - 31)	31	(20 - 46)	54	(41 - 72)
Maine	18	(14 - 23)	2	(1 - 3)	8	(6 - 10)	8	(5 - 12)	16	(12 - 21)
Maryland	91	(71 - 116)	10	(8 - 14)	37	(28 - 48)	44	(30 - 65)	81	(62 - 105)
Massachusetts	116	(92 - 147)	16	(12 - 21)	47	(36 - 62)	53	(36 - 78)	100	(77 - 130)
Michigan	140	(120 - 164)	20	(17 - 25)	55	(47 - 64)	65	(50 - 86)	120	(101 - 143)
Minnesota	68	(52 - 89)	10	(8 - 14)	26	(19 - 36)	32	(21 - 48)	58	(43 - 78)
Mississippi	47	(37 - 60)	6	(4 - 8)	18	(13 - 24)	24	(16 - 34)	41	(32 - 54)
Missouri	93	(73 - 118)	11	(8 - 16)	37	(28 - 49)	44	(31 - 64)	82	(63 - 105)
Montana	17	(13 - 21)	2	(2 - 3)	7	(5 - 9)	8	(5 - 11)	14	(11 - 19)

(continued)

Table 19 Illicit Drug Dependence in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Nebraska	24	(18 - 31)	3	(2 - 4)	8	(6 - 11)	12	(8 - 19)	21	(15 - 28)
Nevada	44	(33 - 58)	6	(4 - 8)	15	(11 - 20)	23	(15 - 34)	38	(28 - 51)
New Hampshire	22	(17 - 28)	3	(2 - 4)	9	(7 - 12)	10	(7 - 14)	19	(15 - 24)
New Jersey	123	(95 - 158)	16	(12 - 22)	46	(34 - 61)	61	(41 - 90)	106	(81 - 140)
New Mexico	33	(26 - 42)	6	(4 - 8)	14	(10 - 18)	13	(8 - 20)	27	(20 - 35)
New York	327	(277 - 386)	34	(27 - 42)	126	(106 - 149)	167	(126 - 221)	293	(245 - 351)
North Carolina	144	(113 - 184)	19	(14 - 26)	56	(42 - 75)	69	(47 - 100)	125	(96 - 163)
North Dakota	8	(6 - 11)	1	(1 - 1)	4	(3 - 5)	3	(2 - 6)	7	(5 - 10)
Ohio	193	(164 - 227)	23	(19 - 28)	67	(57 - 78)	103	(79 - 134)	170	(142 - 204)
Oklahoma	54	(41 - 71)	6	(5 - 9)	20	(14 - 27)	28	(18 - 41)	47	(35 - 64)
Oregon	72	(56 - 93)	8	(6 - 11)	28	(21 - 36)	36	(24 - 54)	64	(49 - 84)
Pennsylvania	194	(163 - 231)	23	(18 - 29)	74	(62 - 88)	97	(73 - 130)	171	(141 - 207)
Rhode Island	19	(15 - 25)	2	(2 - 3)	8	(6 - 10)	9	(6 - 14)	17	(13 - 22)
South Carolina	62	(47 - 80)	8	(6 - 11)	21	(16 - 29)	32	(22 - 48)	54	(40 - 72)
South Dakota	11	(8 - 14)	2	(1 - 2)	4	(3 - 5)	5	(4 - 8)	9	(7 - 13)
Tennessee	113	(87 - 145)	12	(9 - 17)	44	(33 - 58)	57	(38 - 84)	100	(76 - 132)
Texas	334	(280 - 397)	53	(42 - 66)	115	(95 - 140)	165	(123 - 223)	281	(230 - 343)
Utah	40	(31 - 51)	6	(4 - 8)	16	(12 - 22)	17	(12 - 26)	34	(26 - 44)
Vermont	12	(10 - 15)	1	(1 - 2)	6	(5 - 8)	5	(3 - 7)	11	(9 - 14)
Virginia	116	(90 - 150)	15	(11 - 21)	46	(34 - 61)	55	(37 - 83)	101	(77 - 133)
Washington	110	(86 - 140)	14	(11 - 19)	44	(33 - 58)	51	(34 - 75)	95	(73 - 125)
West Virginia	33	(26 - 41)	3	(2 - 4)	15	(12 - 19)	15	(10 - 22)	29	(23 - 38)
Wisconsin	81	(60 - 109)	10	(7 - 13)	27	(20 - 37)	45	(29 - 70)	72	(52 - 99)
Wyoming	7	(5 - 9)	1	(1 - 1)	2	(2 - 3)	3	(2 - 5)	6	(4 - 8)

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 20 Dependence or Abuse of Illicit Drugs or Alcohol in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	21,413	(20,800 - 22,042)	1,752	(1,673 - 1,835)	6,586	(6,401 - 6,776)	13,075	(12,505 - 13,668)	19,661	(19,059 - 20,279)
Northeast	4,091	(3,862 - 4,335)	311	(287 - 336)	1,220	(1,162 - 1,281)	2,560	(2,346 - 2,791)	3,780	(3,556 - 4,020)
Midwest	4,724	(4,514 - 4,941)	364	(341 - 388)	1,461	(1,402 - 1,522)	2,898	(2,703 - 3,107)	4,360	(4,154 - 4,573)
South	7,108	(6,777 - 7,453)	585	(547 - 626)	2,206	(2,108 - 2,308)	4,317	(4,004 - 4,651)	6,523	(6,199 - 6,861)
West	5,490	(5,176 - 5,820)	492	(451 - 535)	1,699	(1,607 - 1,794)	3,300	(3,012 - 3,614)	4,999	(4,692 - 5,322)
Alabama	263	(218 - 316)	23	(18 - 28)	84	(71 - 98)	156	(119 - 204)	240	(197 - 292)
Alaska	52	(44 - 62)	5	(4 - 6)	16	(13 - 18)	32	(25 - 40)	48	(40 - 57)
Arizona	552	(460 - 658)	46	(36 - 58)	150	(129 - 172)	356	(277 - 455)	506	(417 - 609)
Arkansas	164	(135 - 199)	14	(11 - 18)	49	(41 - 57)	101	(77 - 132)	150	(122 - 183)
California	2,813	(2,561 - 3,087)	275	(242 - 313)	907	(833 - 986)	1,631	(1,406 - 1,888)	2,538	(2,292 - 2,807)
Colorado	414	(353 - 484)	32	(25 - 39)	132	(115 - 150)	250	(198 - 315)	382	(323 - 451)
Connecticut	268	(225 - 319)	21	(17 - 26)	79	(69 - 91)	168	(130 - 215)	247	(205 - 296)
Delaware	59	(49 - 71)	4	(4 - 6)	19	(16 - 22)	36	(28 - 46)	55	(45 - 66)
District of Columbia	67	(57 - 78)	2	(2 - 3)	19	(17 - 22)	45	(36 - 56)	65	(55 - 76)
Florida	1,195	(1,083 - 1,320)	90	(78 - 103)	338	(310 - 369)	767	(665 - 884)	1,106	(996 - 1,228)
Georgia	548	(456 - 657)	52	(41 - 65)	183	(156 - 214)	312	(237 - 410)	496	(408 - 601)
Hawaii	94	(77 - 112)	7	(6 - 9)	27	(23 - 31)	60	(46 - 77)	86	(71 - 105)
Idaho	116	(98 - 137)	10	(8 - 12)	35	(30 - 40)	71	(56 - 90)	106	(89 - 127)
Illinois	891	(811 - 978)	68	(59 - 78)	300	(276 - 324)	524	(452 - 606)	823	(745 - 909)
Indiana	444	(372 - 527)	31	(25 - 39)	140	(121 - 161)	273	(212 - 350)	412	(343 - 495)
Iowa	219	(186 - 256)	17	(14 - 21)	75	(65 - 86)	127	(100 - 161)	202	(170 - 238)
Kansas	194	(165 - 227)	17	(13 - 21)	64	(56 - 74)	113	(89 - 143)	177	(149 - 210)
Kentucky	231	(191 - 279)	21	(16 - 27)	71	(60 - 84)	139	(105 - 182)	210	(172 - 257)
Louisiana	275	(231 - 327)	21	(17 - 26)	88	(76 - 100)	167	(129 - 215)	255	(212 - 305)
Maine	77	(63 - 93)	6	(4 - 7)	24	(21 - 28)	47	(36 - 63)	71	(58 - 87)
Maryland	325	(267 - 394)	26	(20 - 33)	109	(93 - 126)	191	(142 - 254)	299	(244 - 366)
Massachusetts	573	(485 - 674)	44	(36 - 54)	178	(156 - 203)	350	(274 - 444)	528	(443 - 628)
Michigan	714	(649 - 784)	58	(51 - 67)	203	(187 - 221)	452	(393 - 519)	655	(592 - 725)
Minnesota	402	(341 - 473)	32	(25 - 40)	124	(108 - 141)	247	(195 - 311)	371	(312 - 439)
Mississippi	170	(142 - 202)	14	(11 - 18)	55	(48 - 64)	100	(77 - 129)	156	(129 - 187)
Missouri	363	(304 - 433)	33	(26 - 42)	120	(103 - 138)	211	(161 - 274)	330	(273 - 398)
Montana	85	(73 - 99)	7	(6 - 9)	26	(23 - 29)	52	(42 - 65)	78	(66 - 92)

(continued)

Table 20 Dependence or Abuse of Illicit Drugs or Alcohol in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	120	(100 - 142)	9	(7 - 11)	41	(35 - 47)	70	(54 - 90)	111	(92 - 132)
Nevada	230	(191 - 277)	17	(13 - 22)	57	(49 - 66)	157	(122 - 200)	214	(176 - 259)
New Hampshire	98	(83 - 116)	9	(7 - 11)	30	(26 - 34)	59	(45 - 76)	89	(74 - 107)
New Jersey	601	(502 - 719)	53	(42 - 66)	169	(146 - 195)	379	(293 - 489)	548	(453 - 663)
New Mexico	153	(129 - 181)	15	(12 - 19)	47	(41 - 54)	90	(70 - 116)	137	(114 - 165)
New York	1,377	(1,244 - 1,523)	101	(88 - 116)	419	(384 - 457)	857	(735 - 996)	1,276	(1,146 - 1,420)
North Carolina	548	(456 - 657)	49	(39 - 61)	177	(153 - 204)	322	(245 - 423)	499	(410 - 605)
North Dakota	52	(45 - 61)	3	(3 - 4)	21	(18 - 24)	28	(22 - 36)	49	(42 - 57)
Ohio	850	(775 - 931)	62	(54 - 71)	240	(222 - 260)	547	(479 - 625)	788	(715 - 868)
Oklahoma	270	(226 - 321)	19	(15 - 25)	76	(65 - 89)	174	(137 - 222)	251	(208 - 301)
Oregon	317	(267 - 375)	23	(19 - 28)	84	(73 - 97)	209	(166 - 263)	294	(246 - 350)
Pennsylvania	948	(858 - 1,046)	67	(58 - 77)	271	(248 - 296)	610	(529 - 702)	881	(793 - 977)
Rhode Island	96	(81 - 113)	6	(5 - 7)	30	(26 - 34)	60	(47 - 76)	90	(75 - 107)
South Carolina	307	(256 - 368)	24	(19 - 30)	87	(75 - 101)	196	(152 - 252)	283	(233 - 342)
South Dakota	67	(56 - 78)	5	(4 - 7)	19	(16 - 22)	42	(34 - 53)	61	(51 - 73)
Tennessee	433	(362 - 516)	34	(27 - 42)	128	(109 - 149)	272	(212 - 347)	400	(331 - 480)
Texas	1,604	(1,455 - 1,767)	142	(123 - 164)	504	(462 - 548)	958	(823 - 1,113)	1,462	(1,316 - 1,622)
Utah	137	(112 - 167)	15	(11 - 19)	53	(44 - 64)	70	(51 - 95)	123	(99 - 151)
Vermont	54	(46 - 63)	4	(3 - 5)	19	(17 - 21)	31	(24 - 39)	50	(42 - 58)
Virginia	538	(456 - 634)	42	(34 - 53)	182	(158 - 208)	314	(245 - 400)	496	(416 - 589)
Washington	489	(413 - 579)	37	(30 - 46)	153	(133 - 175)	299	(234 - 380)	452	(378 - 539)
West Virginia	111	(94 - 131)	8	(6 - 10)	37	(32 - 42)	66	(51 - 85)	103	(86 - 123)
Wisconsin	409	(341 - 488)	29	(23 - 36)	115	(99 - 133)	265	(205 - 339)	380	(314 - 458)
Wyoming	38	(32 - 45)	3	(2 - 4)	12	(10 - 14)	23	(18 - 30)	35	(29 - 42)

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 21 *Needing But Not Receiving Treatment for Illicit Drug Use in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs*

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Total U.S.	6,125	(5,835 - 6,428)	1,070	(1,011 - 1,132)	2,410	(2,300 - 2,525)	2,645	(2,400 - 2,915)	5,055	(4,777 - 5,349)
Northeast	1,109	(1,025 - 1,200)	180	(162 - 199)	446	(409 - 487)	483	(414 - 563)	929	(849 - 1,017)
Midwest	1,243	(1,166 - 1,324)	217	(199 - 236)	493	(460 - 528)	534	(467 - 610)	1,026	(953 - 1,105)
South	2,177	(2,039 - 2,323)	370	(340 - 403)	819	(762 - 879)	988	(871 - 1,120)	1,807	(1,677 - 1,947)
West	1,596	(1,462 - 1,742)	303	(270 - 341)	652	(592 - 718)	641	(533 - 770)	1,293	(1,166 - 1,432)
Alabama	84	(69 - 103)	14	(10 - 19)	34	(26 - 44)	36	(25 - 52)	70	(56 - 88)
Alaska	16	(13 - 19)	3	(2 - 4)	5	(4 - 7)	7	(5 - 11)	13	(10 - 16)
Arizona	149	(118 - 187)	31	(22 - 42)	53	(40 - 69)	65	(44 - 96)	118	(91 - 153)
Arkansas	53	(43 - 65)	9	(7 - 13)	20	(16 - 26)	23	(16 - 33)	44	(35 - 55)
California	858	(754 - 975)	166	(140 - 197)	362	(312 - 418)	330	(250 - 435)	691	(594 - 804)
Colorado	108	(90 - 129)	20	(14 - 26)	52	(41 - 65)	36	(25 - 53)	88	(72 - 108)
Connecticut	68	(55 - 84)	11	(8 - 15)	26	(20 - 34)	30	(21 - 44)	56	(44 - 72)
Delaware	17	(14 - 21)	3	(2 - 4)	7	(5 - 9)	8	(5 - 12)	15	(12 - 19)
District of Columbia	14	(11 - 18)	1	(1 - 2)	6	(4 - 7)	7	(5 - 11)	13	(10 - 17)
Florida	378	(331 - 434)	56	(47 - 68)	143	(123 - 166)	179	(140 - 230)	322	(276 - 376)
Georgia	170	(134 - 216)	31	(23 - 42)	57	(42 - 77)	82	(55 - 122)	139	(105 - 183)
Hawaii	21	(17 - 27)	4	(3 - 5)	8	(6 - 10)	10	(6 - 15)	17	(13 - 23)
Idaho	30	(25 - 37)	6	(4 - 8)	12	(9 - 15)	13	(9 - 18)	25	(20 - 31)
Illinois	241	(212 - 275)	41	(34 - 49)	109	(95 - 125)	92	(69 - 121)	201	(173 - 233)
Indiana	118	(95 - 146)	18	(13 - 25)	47	(36 - 61)	53	(36 - 78)	100	(78 - 127)
Iowa	50	(40 - 62)	10	(7 - 13)	20	(15 - 27)	20	(13 - 30)	40	(31 - 51)
Kansas	49	(39 - 60)	9	(7 - 13)	20	(15 - 26)	19	(13 - 28)	39	(31 - 50)
Kentucky	81	(66 - 98)	15	(11 - 20)	30	(23 - 39)	35	(25 - 51)	66	(53 - 82)
Louisiana	83	(67 - 104)	13	(10 - 18)	29	(22 - 37)	41	(28 - 60)	70	(54 - 89)
Maine	23	(19 - 29)	3	(2 - 4)	10	(8 - 13)	10	(7 - 15)	20	(16 - 26)
Maryland	110	(90 - 134)	18	(13 - 24)	45	(35 - 59)	46	(33 - 65)	92	(74 - 114)
Massachusetts	137	(112 - 168)	29	(22 - 37)	55	(42 - 71)	54	(37 - 78)	108	(85 - 137)
Michigan	179	(156 - 205)	35	(29 - 42)	68	(59 - 79)	76	(57 - 100)	144	(122 - 170)
Minnesota	97	(79 - 118)	18	(14 - 24)	39	(30 - 51)	39	(27 - 56)	79	(63 - 98)
Mississippi	62	(50 - 76)	9	(7 - 12)	23	(18 - 30)	30	(21 - 42)	53	(42 - 67)
Missouri	126	(104 - 152)	20	(15 - 27)	51	(40 - 64)	55	(39 - 78)	106	(86 - 131)
Montana	20	(17 - 25)	4	(3 - 5)	8	(7 - 11)	8	(6 - 11)	16	(13 - 20)

(continued)

Table 21 *Needing But Not Receiving Treatment for Illicit Drug Use in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)*

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	30	(24 - 38)	5	(4 - 7)	11	(8 - 14)	14	(9 - 21)	25	(19 - 32)
Nevada	53	(41 - 69)	9	(7 - 13)	16	(12 - 21)	28	(19 - 42)	44	(33 - 59)
New Hampshire	28	(23 - 34)	6	(5 - 8)	12	(9 - 15)	10	(7 - 15)	22	(17 - 27)
New Jersey	159	(130 - 195)	31	(23 - 41)	63	(48 - 83)	66	(46 - 94)	129	(103 - 161)
New Mexico	47	(39 - 57)	11	(8 - 15)	19	(15 - 24)	18	(12 - 26)	36	(29 - 45)
New York	411	(358 - 472)	56	(46 - 69)	168	(146 - 194)	186	(144 - 242)	355	(304 - 414)
North Carolina	192	(157 - 234)	33	(25 - 43)	73	(56 - 94)	87	(61 - 123)	160	(127 - 200)
North Dakota	10	(8 - 13)	2	(1 - 2)	4	(3 - 6)	4	(3 - 6)	8	(6 - 11)
Ohio	229	(200 - 263)	38	(32 - 46)	83	(72 - 96)	108	(83 - 139)	191	(163 - 224)
Oklahoma	68	(54 - 86)	11	(8 - 15)	25	(19 - 33)	32	(22 - 47)	57	(44 - 74)
Oregon	91	(74 - 110)	15	(12 - 20)	36	(28 - 46)	39	(27 - 56)	75	(60 - 94)
Pennsylvania	245	(213 - 283)	38	(31 - 46)	94	(79 - 111)	113	(87 - 147)	207	(176 - 244)
Rhode Island	23	(18 - 28)	3	(2 - 5)	10	(8 - 13)	9	(6 - 14)	19	(15 - 24)
South Carolina	89	(72 - 108)	16	(12 - 21)	33	(25 - 42)	40	(28 - 57)	73	(58 - 91)
South Dakota	13	(10 - 16)	3	(2 - 4)	4	(3 - 6)	6	(4 - 9)	10	(8 - 13)
Tennessee	139	(112 - 170)	21	(15 - 27)	52	(40 - 67)	67	(47 - 95)	118	(94 - 148)
Texas	456	(398 - 524)	90	(74 - 108)	167	(142 - 196)	200	(153 - 260)	367	(312 - 432)
Utah	55	(44 - 67)	11	(8 - 15)	21	(15 - 28)	23	(16 - 32)	43	(34 - 55)
Vermont	15	(12 - 18)	2	(2 - 3)	8	(6 - 10)	5	(3 - 8)	13	(10 - 16)
Virginia	143	(118 - 174)	26	(19 - 34)	59	(46 - 77)	58	(41 - 82)	117	(94 - 146)
Washington	140	(116 - 169)	22	(16 - 29)	58	(46 - 74)	60	(43 - 84)	118	(96 - 145)
West Virginia	36	(29 - 45)	4	(3 - 6)	16	(13 - 20)	16	(11 - 23)	32	(25 - 40)
Wisconsin	103	(83 - 127)	18	(14 - 25)	36	(28 - 48)	48	(33 - 70)	85	(67 - 107)
Wyoming	9	(7 - 11)	2	(1 - 2)	3	(2 - 4)	4	(3 - 6)	7	(6 - 10)

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs, but not receiving treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 22 *Needing But Not Receiving Treatment for Alcohol Use in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs*

State	12+		12-17		18-25		26+		18+	
	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)	(Estimate)	(95% Confidence Interval)
Total U.S.	16,480	(15,945 - 17,031)	994	(931 - 1,062)	4,945	(4,781 - 5,114)	10,541	(10,037 - 11,067)	15,485	(14,955 - 16,032)
Northeast	3,133	(2,933 - 3,347)	174	(158 - 191)	920	(871 - 972)	2,039	(1,854 - 2,241)	2,959	(2,762 - 3,171)
Midwest	3,686	(3,491 - 3,890)	209	(192 - 227)	1,128	(1,078 - 1,179)	2,349	(2,170 - 2,543)	3,477	(3,285 - 3,679)
South	5,403	(5,112 - 5,709)	333	(306 - 363)	1,613	(1,539 - 1,690)	3,456	(3,189 - 3,744)	5,069	(4,783 - 5,371)
West	4,258	(3,965 - 4,571)	279	(249 - 312)	1,283	(1,205 - 1,366)	2,696	(2,431 - 2,988)	3,980	(3,691 - 4,288)
Alabama	199	(159 - 248)	13	(10 - 17)	60	(50 - 72)	126	(93 - 170)	186	(147 - 234)
Alaska	42	(34 - 51)	2	(2 - 3)	12	(10 - 15)	27	(20 - 36)	39	(32 - 49)
Arizona	410	(330 - 507)	24	(18 - 31)	113	(95 - 133)	273	(205 - 362)	386	(307 - 482)
Arkansas	118	(94 - 148)	9	(7 - 12)	34	(29 - 41)	75	(55 - 102)	109	(86 - 139)
California	2,183	(1,953 - 2,437)	159	(134 - 188)	677	(612 - 747)	1,347	(1,143 - 1,584)	2,024	(1,799 - 2,274)
Colorado	337	(280 - 405)	17	(13 - 23)	105	(91 - 121)	215	(166 - 277)	320	(264 - 386)
Connecticut	208	(168 - 256)	12	(9 - 15)	62	(52 - 73)	134	(101 - 178)	196	(157 - 244)
Delaware	44	(35 - 55)	2	(2 - 3)	13	(11 - 16)	28	(21 - 38)	42	(33 - 52)
District of Columbia	56	(46 - 67)	1	(1 - 1)	16	(14 - 19)	39	(30 - 49)	55	(45 - 66)
Florida	885	(784 - 999)	52	(44 - 62)	240	(216 - 266)	592	(501 - 699)	832	(733 - 946)
Georgia	414	(334 - 512)	30	(23 - 39)	130	(108 - 155)	254	(188 - 342)	384	(306 - 481)
Hawaii	83	(67 - 101)	4	(3 - 6)	23	(20 - 28)	55	(42 - 72)	78	(63 - 97)
Idaho	88	(72 - 107)	6	(4 - 7)	27	(22 - 31)	56	(42 - 73)	82	(67 - 101)
Illinois	699	(627 - 780)	39	(33 - 47)	227	(206 - 249)	433	(368 - 508)	660	(588 - 739)
Indiana	343	(279 - 420)	19	(14 - 24)	109	(92 - 128)	215	(162 - 285)	324	(262 - 400)
Iowa	189	(157 - 227)	10	(8 - 13)	63	(54 - 73)	116	(89 - 150)	179	(147 - 216)
Kansas	153	(125 - 185)	10	(8 - 13)	50	(42 - 59)	93	(70 - 122)	143	(116 - 175)
Kentucky	161	(126 - 204)	12	(9 - 16)	48	(39 - 59)	100	(72 - 139)	148	(115 - 191)
Louisiana	220	(181 - 267)	11	(9 - 14)	68	(58 - 80)	141	(108 - 184)	209	(171 - 255)
Maine	59	(47 - 73)	4	(3 - 5)	18	(15 - 21)	37	(27 - 50)	55	(44 - 69)
Maryland	231	(181 - 293)	13	(10 - 18)	74	(60 - 90)	144	(104 - 198)	217	(169 - 279)
Massachusetts	447	(372 - 536)	23	(18 - 30)	139	(119 - 161)	285	(221 - 366)	424	(350 - 512)
Michigan	563	(504 - 629)	32	(27 - 38)	156	(142 - 171)	375	(320 - 438)	531	(472 - 597)
Minnesota	318	(261 - 386)	17	(14 - 23)	99	(84 - 116)	202	(154 - 264)	301	(245 - 368)
Mississippi	126	(102 - 156)	8	(6 - 11)	38	(31 - 45)	81	(61 - 107)	118	(95 - 147)
Missouri	261	(210 - 323)	19	(15 - 24)	85	(71 - 101)	157	(115 - 214)	242	(193 - 304)
Montana	66	(56 - 79)	4	(3 - 5)	20	(17 - 23)	42	(33 - 54)	62	(52 - 75)

(continued)

Table 22 Needing But Not Receiving Treatment for Alcohol Use in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	12+ (Estimate)	12+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)	18+ (Estimate)	18+ (95% Confidence Interval)
Nebraska	91	(73 - 111)	5	(4 - 7)	32	(28 - 38)	53	(39 - 71)	85	(68 - 105)
Nevada	191	(154 - 236)	10	(7 - 13)	46	(39 - 55)	135	(103 - 176)	181	(145 - 226)
New Hampshire	74	(60 - 90)	5	(4 - 6)	23	(19 - 26)	46	(35 - 61)	69	(56 - 85)
New Jersey	453	(368 - 557)	29	(22 - 38)	124	(104 - 147)	300	(227 - 396)	424	(341 - 527)
New Mexico	111	(91 - 135)	7	(6 - 9)	34	(29 - 40)	70	(53 - 92)	104	(84 - 128)
New York	1,055	(933 - 1,193)	58	(48 - 69)	309	(279 - 341)	688	(577 - 819)	997	(876 - 1,134)
North Carolina	404	(321 - 505)	29	(22 - 38)	123	(102 - 148)	251	(184 - 342)	375	(295 - 474)
North Dakota	47	(40 - 56)	2	(1 - 2)	19	(17 - 22)	26	(20 - 34)	45	(38 - 54)
Ohio	651	(584 - 725)	35	(30 - 42)	178	(163 - 195)	437	(375 - 509)	615	(549 - 689)
Oklahoma	207	(166 - 257)	11	(8 - 15)	58	(48 - 69)	138	(103 - 184)	196	(156 - 246)
Oregon	239	(195 - 291)	12	(9 - 15)	63	(53 - 73)	164	(126 - 214)	227	(184 - 280)
Pennsylvania	721	(644 - 807)	39	(32 - 46)	209	(188 - 232)	473	(404 - 554)	682	(606 - 768)
Rhode Island	80	(66 - 97)	3	(3 - 4)	24	(21 - 28)	52	(40 - 68)	77	(63 - 93)
South Carolina	241	(196 - 295)	14	(11 - 18)	62	(52 - 74)	165	(126 - 215)	227	(183 - 281)
South Dakota	56	(47 - 67)	3	(3 - 5)	16	(14 - 19)	37	(29 - 47)	53	(43 - 63)
Tennessee	314	(251 - 391)	18	(13 - 24)	87	(73 - 104)	209	(155 - 279)	296	(234 - 372)
Texas	1,284	(1,148 - 1,434)	80	(67 - 95)	400	(362 - 440)	804	(683 - 944)	1,204	(1,070 - 1,353)
Utah	92	(72 - 116)	9	(7 - 13)	37	(30 - 45)	46	(31 - 66)	83	(64 - 106)
Vermont	37	(30 - 45)	2	(2 - 3)	13	(11 - 15)	22	(16 - 29)	34	(28 - 42)
Virginia	421	(346 - 512)	25	(20 - 32)	135	(115 - 158)	261	(197 - 344)	396	(322 - 486)
Washington	386	(316 - 470)	22	(17 - 29)	117	(99 - 137)	247	(188 - 323)	364	(295 - 447)
West Virginia	79	(64 - 98)	5	(4 - 6)	26	(22 - 30)	49	(36 - 66)	74	(59 - 93)
Wisconsin	315	(257 - 385)	16	(12 - 21)	93	(78 - 109)	206	(157 - 270)	299	(241 - 368)
Wyoming	31	(25 - 37)	2	(1 - 2)	10	(8 - 12)	19	(15 - 25)	29	(24 - 35)

NOTE: Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but not receiving treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 23 Serious Mental Illness in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs

State	18+ (Estimate)	18+ (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)
Total U.S.	9,135	(8,663 - 9,636)	1,317	(1,238 - 1,401)	7,818	(7,356 - 8,313)
Northeast	1,565	(1,422 - 1,721)	236	(214 - 260)	1,329	(1,191 - 1,482)
Midwest	2,140	(1,991 - 2,301)	293	(271 - 316)	1,848	(1,702 - 2,005)
South	3,298	(3,053 - 3,563)	465	(427 - 507)	2,832	(2,596 - 3,091)
West	2,133	(1,929 - 2,359)	323	(289 - 361)	1,810	(1,612 - 2,033)
Alabama	205	(161 - 262)	21	(16 - 28)	184	(141 - 240)
Alaska	21	(16 - 26)	3	(2 - 4)	18	(14 - 23)
Arizona	233	(178 - 303)	31	(23 - 40)	202	(150 - 271)
Arkansas	117	(91 - 150)	14	(11 - 17)	104	(78 - 137)
California	919	(775 - 1,088)	163	(136 - 195)	756	(618 - 924)
Colorado	159	(123 - 204)	24	(19 - 31)	134	(100 - 180)
Connecticut	94	(72 - 121)	13	(10 - 17)	81	(60 - 108)
Delaware	26	(20 - 33)	4	(3 - 5)	22	(16 - 29)
District of Columbia	18	(14 - 23)	3	(3 - 4)	14	(10 - 19)
Florida	499	(419 - 594)	69	(57 - 82)	430	(353 - 524)
Georgia	250	(192 - 324)	39	(30 - 50)	211	(156 - 284)
Hawaii	34	(26 - 46)	5	(4 - 6)	30	(21 - 41)
Idaho	60	(47 - 76)	8	(6 - 10)	52	(39 - 68)
Illinois	329	(278 - 390)	50	(42 - 59)	280	(230 - 339)
Indiana	235	(184 - 300)	29	(23 - 37)	206	(157 - 271)
Iowa	105	(83 - 133)	15	(11 - 19)	91	(69 - 118)
Kansas	86	(67 - 110)	12	(9 - 15)	75	(56 - 98)
Kentucky	150	(117 - 191)	19	(15 - 24)	131	(99 - 172)
Louisiana	126	(98 - 162)	18	(14 - 23)	108	(81 - 144)
Maine	46	(36 - 59)	5	(4 - 7)	40	(30 - 54)
Maryland	138	(106 - 179)	22	(17 - 29)	116	(86 - 156)
Massachusetts	200	(155 - 256)	30	(24 - 38)	170	(127 - 226)
Michigan	319	(274 - 372)	46	(40 - 54)	273	(229 - 326)
Minnesota	155	(121 - 198)	24	(19 - 30)	131	(98 - 174)
Mississippi	105	(83 - 132)	12	(9 - 16)	92	(71 - 120)
Missouri	212	(166 - 269)	24	(19 - 31)	188	(144 - 245)
Montana	38	(31 - 48)	5	(4 - 6)	34	(26 - 43)

(continued)

Table 23 Serious Mental Illness in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	18+ (Estimate)	18+ (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)
Nebraska	61	(48 - 77)	8	(6 - 10)	52	(40 - 68)
Nevada	80	(62 - 104)	10	(8 - 14)	70	(52 - 93)
New Hampshire	45	(35 - 58)	6	(4 - 7)	39	(29 - 52)
New Jersey	204	(156 - 268)	33	(26 - 42)	171	(125 - 235)
New Mexico	68	(54 - 87)	9	(7 - 11)	60	(46 - 78)
New York	524	(441 - 622)	82	(69 - 98)	442	(362 - 538)
North Carolina	250	(189 - 329)	34	(26 - 45)	216	(158 - 294)
North Dakota	20	(16 - 26)	3	(3 - 4)	17	(12 - 22)
Ohio	421	(363 - 488)	55	(47 - 64)	366	(310 - 432)
Oklahoma	139	(110 - 177)	16	(13 - 21)	123	(94 - 160)
Oregon	144	(114 - 182)	18	(14 - 22)	127	(98 - 164)
Pennsylvania	397	(337 - 467)	59	(49 - 70)	338	(280 - 407)
Rhode Island	32	(25 - 41)	5	(4 - 7)	27	(20 - 36)
South Carolina	148	(114 - 191)	18	(14 - 23)	130	(97 - 172)
South Dakota	25	(19 - 32)	4	(3 - 5)	21	(16 - 29)
Tennessee	208	(160 - 269)	25	(19 - 32)	183	(137 - 244)
Texas	602	(504 - 717)	108	(91 - 127)	494	(401 - 609)
Utah	95	(76 - 118)	18	(14 - 23)	77	(59 - 100)
Vermont	24	(19 - 30)	3	(2 - 4)	21	(16 - 27)
Virginia	242	(189 - 308)	37	(29 - 48)	204	(154 - 270)
Washington	264	(203 - 341)	28	(22 - 36)	236	(177 - 313)
West Virginia	76	(60 - 96)	7	(5 - 9)	69	(53 - 89)
Wisconsin	172	(135 - 220)	24	(19 - 31)	148	(112 - 195)
Wyoming	18	(14 - 23)	2	(2 - 3)	16	(12 - 20)

NOTE: Serious mental illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in serious functional impairment. Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 (Revised March 2012 and October 2013) and 2011 (Revised October 2013).

Table 24 Any Mental Illness in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs

State	18+ (Estimate)	18+ (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)
Total U.S.	41,423	(40,480 - 42,392)	6,261	(6,091 - 6,434)	35,162	(34,239 - 36,114)
Northeast	7,539	(7,216 - 7,873)	1,121	(1,071 - 1,174)	6,417	(6,100 - 6,747)
Midwest	9,233	(8,917 - 9,557)	1,391	(1,342 - 1,443)	7,842	(7,531 - 8,161)
South	15,042	(14,519 - 15,587)	2,165	(2,081 - 2,251)	12,878	(12,366 - 13,411)
West	9,608	(9,157 - 10,080)	1,583	(1,504 - 1,665)	8,025	(7,591 - 8,482)
Alabama	779	(694 - 872)	103	(90 - 118)	676	(594 - 767)
Alaska	97	(85 - 110)	17	(15 - 19)	80	(68 - 92)
Arizona	932	(811 - 1,066)	151	(132 - 171)	781	(665 - 912)
Arkansas	447	(392 - 508)	59	(52 - 67)	388	(334 - 448)
California	4,527	(4,171 - 4,907)	797	(732 - 867)	3,730	(3,389 - 4,099)
Colorado	661	(579 - 753)	115	(101 - 130)	547	(468 - 636)
Connecticut	450	(390 - 517)	66	(57 - 75)	384	(327 - 450)
Delaware	124	(109 - 141)	20	(17 - 23)	104	(90 - 121)
District of Columbia	96	(85 - 109)	19	(16 - 21)	77	(67 - 89)
Florida	2,365	(2,178 - 2,565)	338	(311 - 367)	2,027	(1,845 - 2,223)
Georgia	1,208	(1,051 - 1,383)	183	(159 - 210)	1,025	(873 - 1,196)
Hawaii	174	(152 - 199)	26	(22 - 29)	148	(127 - 172)
Idaho	257	(227 - 290)	36	(31 - 40)	222	(193 - 253)
Illinois	1,535	(1,413 - 1,665)	248	(229 - 268)	1,287	(1,169 - 1,414)
Indiana	937	(820 - 1,066)	149	(131 - 169)	788	(675 - 914)
Iowa	431	(381 - 486)	65	(57 - 74)	366	(318 - 419)
Kansas	361	(317 - 410)	58	(51 - 66)	303	(260 - 351)
Kentucky	634	(554 - 722)	80	(70 - 92)	554	(475 - 641)
Louisiana	619	(545 - 701)	91	(80 - 103)	529	(457 - 609)
Maine	177	(154 - 204)	26	(23 - 30)	151	(128 - 178)
Maryland	764	(665 - 875)	113	(98 - 128)	652	(556 - 760)
Massachusetts	908	(798 - 1,030)	144	(127 - 164)	764	(658 - 883)
Michigan	1,471	(1,368 - 1,580)	212	(196 - 228)	1,259	(1,159 - 1,365)
Minnesota	691	(608 - 784)	112	(98 - 127)	579	(499 - 670)
Mississippi	420	(371 - 474)	57	(50 - 65)	363	(315 - 416)
Missouri	868	(764 - 981)	121	(106 - 138)	746	(646 - 857)
Montana	148	(131 - 166)	22	(19 - 24)	126	(110 - 144)

(continued)

Table 24 Any Mental Illness in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	18+ (Estimate)	18+ (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)
Nebraska	239	(209 - 271)	38	(34 - 43)	200	(172 - 232)
Nevada	332	(285 - 384)	48	(41 - 55)	284	(239 - 335)
New Hampshire	196	(171 - 223)	26	(23 - 30)	169	(146 - 196)
New Jersey	1,066	(922 - 1,228)	150	(131 - 172)	916	(776 - 1,075)
New Mexico	280	(245 - 319)	44	(38 - 49)	237	(203 - 275)
New York	2,787	(2,576 - 3,010)	419	(387 - 453)	2,367	(2,162 - 2,588)
North Carolina	1,168	(1,010 - 1,344)	161	(139 - 185)	1,007	(856 - 1,179)
North Dakota	86	(75 - 98)	17	(15 - 19)	69	(59 - 81)
Ohio	1,754	(1,634 - 1,882)	241	(224 - 260)	1,513	(1,394 - 1,640)
Oklahoma	559	(491 - 633)	78	(68 - 89)	481	(416 - 553)
Oregon	609	(537 - 687)	85	(75 - 96)	524	(454 - 601)
Pennsylvania	1,712	(1,585 - 1,846)	250	(229 - 274)	1,462	(1,339 - 1,593)
Rhode Island	148	(129 - 169)	24	(21 - 28)	124	(106 - 145)
South Carolina	641	(560 - 730)	83	(72 - 95)	558	(480 - 645)
South Dakota	107	(94 - 122)	17	(15 - 20)	90	(77 - 105)
Tennessee	1,024	(898 - 1,163)	116	(101 - 133)	908	(785 - 1,045)
Texas	2,832	(2,605 - 3,075)	473	(435 - 514)	2,359	(2,138 - 2,597)
Utah	417	(371 - 467)	81	(72 - 92)	336	(293 - 384)
Vermont	94	(83 - 107)	15	(13 - 17)	80	(69 - 92)
Virginia	1,067	(929 - 1,219)	158	(139 - 180)	908	(776 - 1,058)
Washington	1,099	(964 - 1,248)	151	(133 - 171)	948	(818 - 1,093)
West Virginia	296	(260 - 335)	34	(30 - 38)	262	(227 - 301)
Wisconsin	753	(655 - 862)	112	(98 - 129)	641	(547 - 747)
Wyoming	76	(67 - 86)	13	(11 - 14)	63	(55 - 73)

NOTE: Any mental illness (AMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for AMI presented in this table may differ from previously published AMI estimates because of revised estimation procedures. For details, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 (Revised March 2012 and October 2013) and 2011 (Revised October 2013).

Table 25 *Had Serious Thoughts of Suicide in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs*

State	18+ (Estimate)	18+ (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)
Total U.S.	8,662	(8,248 - 9,098)	2,303	(2,194 - 2,418)	6,359	(5,966 - 6,780)
Northeast	1,620	(1,481 - 1,773)	414	(381 - 450)	1,206	(1,077 - 1,350)
Midwest	2,001	(1,861 - 2,149)	517	(486 - 549)	1,484	(1,353 - 1,627)
South	3,033	(2,813 - 3,270)	816	(763 - 872)	2,217	(2,011 - 2,444)
West	2,009	(1,823 - 2,214)	556	(510 - 607)	1,452	(1,279 - 1,649)
Alabama	148	(116 - 188)	42	(34 - 51)	106	(78 - 145)
Alaska	22	(18 - 28)	6	(5 - 8)	16	(12 - 22)
Arizona	199	(155 - 253)	53	(42 - 66)	146	(107 - 198)
Arkansas	91	(72 - 116)	22	(18 - 27)	69	(51 - 93)
California	961	(821 - 1,125)	284	(247 - 327)	677	(549 - 834)
Colorado	136	(105 - 174)	38	(31 - 46)	98	(70 - 136)
Connecticut	98	(76 - 126)	22	(18 - 28)	75	(55 - 102)
Delaware	29	(23 - 37)	7	(6 - 9)	22	(17 - 30)
District of Columbia	20	(16 - 25)	6	(5 - 7)	14	(10 - 19)
Florida	487	(414 - 573)	124	(108 - 142)	363	(294 - 448)
Georgia	242	(189 - 308)	67	(53 - 84)	174	(128 - 238)
Hawaii	35	(27 - 45)	9	(7 - 12)	26	(19 - 35)
Idaho	50	(40 - 63)	13	(10 - 15)	37	(28 - 50)
Illinois	319	(273 - 372)	92	(80 - 105)	227	(184 - 279)
Indiana	205	(162 - 259)	51	(42 - 63)	154	(114 - 207)
Iowa	86	(69 - 106)	23	(19 - 29)	62	(47 - 82)
Kansas	82	(64 - 103)	20	(16 - 25)	61	(46 - 82)
Kentucky	134	(106 - 169)	30	(24 - 37)	105	(79 - 139)
Louisiana	123	(98 - 154)	36	(29 - 43)	88	(65 - 117)
Maine	42	(32 - 54)	8	(7 - 10)	33	(25 - 46)
Maryland	156	(123 - 199)	40	(32 - 50)	116	(86 - 158)
Massachusetts	202	(159 - 256)	49	(39 - 61)	153	(114 - 204)
Michigan	336	(290 - 388)	83	(73 - 94)	253	(210 - 304)
Minnesota	159	(126 - 199)	41	(33 - 51)	118	(88 - 157)
Mississippi	83	(66 - 104)	21	(17 - 26)	62	(46 - 83)
Missouri	185	(147 - 233)	43	(34 - 54)	142	(107 - 189)
Montana	32	(26 - 39)	9	(7 - 11)	23	(17 - 30)

(continued)

Table 25 *Had Serious Thoughts of Suicide in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)*

State	18+ (Estimate)	18+ (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)
Nebraska	52	(41 - 65)	13	(11 - 17)	38	(29 - 52)
Nevada	71	(55 - 91)	17	(13 - 21)	54	(39 - 74)
New Hampshire	49	(38 - 62)	10	(8 - 12)	39	(29 - 52)
New Jersey	228	(178 - 290)	58	(46 - 72)	170	(125 - 230)
New Mexico	57	(45 - 73)	16	(13 - 20)	41	(30 - 56)
New York	556	(476 - 648)	155	(135 - 178)	400	(327 - 489)
North Carolina	257	(203 - 324)	64	(51 - 80)	192	(144 - 256)
North Dakota	18	(14 - 23)	6	(5 - 8)	12	(9 - 17)
Ohio	357	(310 - 411)	95	(84 - 106)	262	(218 - 315)
Oklahoma	126	(100 - 159)	28	(22 - 35)	98	(74 - 130)
Oregon	120	(95 - 152)	29	(24 - 36)	91	(68 - 121)
Pennsylvania	389	(336 - 451)	97	(84 - 112)	292	(243 - 351)
Rhode Island	34	(26 - 43)	9	(7 - 11)	25	(18 - 34)
South Carolina	131	(103 - 166)	32	(25 - 39)	100	(74 - 134)
South Dakota	25	(20 - 32)	6	(5 - 8)	19	(14 - 25)
Tennessee	179	(140 - 228)	41	(33 - 52)	138	(102 - 186)
Texas	529	(449 - 622)	181	(157 - 208)	348	(277 - 436)
Utah	87	(70 - 109)	29	(23 - 35)	59	(44 - 79)
Vermont	24	(19 - 31)	5	(4 - 7)	19	(14 - 25)
Virginia	238	(189 - 298)	64	(52 - 79)	174	(131 - 231)
Washington	223	(176 - 282)	50	(40 - 63)	173	(129 - 230)
West Virginia	59	(47 - 75)	12	(10 - 15)	47	(35 - 62)
Wisconsin	178	(139 - 226)	43	(34 - 53)	135	(100 - 182)
Wyoming	16	(13 - 20)	4	(3 - 5)	12	(9 - 17)

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

Table 26 *Had at Least One Major Depressive Episode in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs*

State	18+ (Estimate)	18+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)
Total U.S.	15,464	(14,889 - 16,062)	2,010	(1,928 - 2,094)	2,834	(2,721 - 2,952)	12,630	(12,068 - 13,219)
Northeast	2,802	(2,615 - 3,002)	334	(310 - 360)	504	(469 - 540)	2,299	(2,115 - 2,498)
Midwest	3,521	(3,327 - 3,724)	448	(424 - 474)	641	(608 - 677)	2,880	(2,689 - 3,082)
South	5,588	(5,281 - 5,913)	731	(691 - 774)	990	(934 - 1,049)	4,597	(4,299 - 4,917)
West	3,553	(3,298 - 3,828)	496	(458 - 537)	699	(648 - 755)	2,854	(2,606 - 3,124)
Alabama	272	(224 - 329)	29	(24 - 35)	45	(37 - 54)	227	(181 - 284)
Alaska	35	(28 - 42)	4	(4 - 5)	7	(6 - 9)	27	(21 - 35)
Arizona	337	(273 - 415)	47	(39 - 58)	63	(51 - 76)	274	(213 - 352)
Arkansas	166	(137 - 202)	20	(16 - 24)	27	(23 - 33)	139	(110 - 175)
California	1,655	(1,459 - 1,876)	258	(227 - 292)	355	(314 - 402)	1,300	(1,111 - 1,519)
Colorado	255	(211 - 307)	32	(26 - 39)	50	(42 - 60)	205	(163 - 256)
Connecticut	173	(141 - 211)	22	(18 - 27)	30	(24 - 36)	143	(112 - 182)
Delaware	46	(38 - 56)	6	(5 - 8)	9	(7 - 11)	37	(29 - 47)
District of Columbia	35	(29 - 41)	2	(2 - 3)	8	(6 - 9)	27	(22 - 34)
Florida	862	(759 - 978)	110	(97 - 124)	152	(134 - 173)	709	(610 - 824)
Georgia	482	(399 - 580)	70	(58 - 86)	90	(74 - 108)	392	(313 - 490)
Hawaii	55	(45 - 69)	8	(6 - 9)	11	(9 - 13)	45	(35 - 58)
Idaho	92	(76 - 112)	11	(9 - 14)	16	(14 - 20)	76	(60 - 95)
Illinois	614	(545 - 691)	87	(77 - 98)	120	(107 - 136)	493	(426 - 570)
Indiana	362	(297 - 440)	38	(31 - 47)	63	(52 - 76)	299	(236 - 377)
Iowa	176	(146 - 212)	18	(15 - 22)	30	(25 - 36)	147	(118 - 183)
Kansas	143	(120 - 171)	20	(16 - 24)	28	(23 - 33)	115	(93 - 143)
Kentucky	247	(202 - 300)	30	(25 - 36)	36	(29 - 44)	211	(167 - 264)
Louisiana	212	(175 - 257)	26	(22 - 32)	39	(33 - 47)	173	(137 - 218)
Maine	71	(58 - 86)	8	(7 - 10)	11	(9 - 13)	60	(47 - 75)
Maryland	289	(237 - 352)	34	(28 - 42)	49	(40 - 59)	240	(190 - 303)
Massachusetts	353	(294 - 423)	44	(36 - 53)	65	(54 - 78)	288	(230 - 358)
Michigan	516	(457 - 583)	78	(70 - 88)	101	(91 - 113)	415	(357 - 482)
Minnesota	246	(201 - 299)	37	(30 - 45)	50	(42 - 60)	196	(153 - 249)
Mississippi	150	(124 - 181)	18	(15 - 22)	24	(19 - 29)	126	(101 - 157)
Missouri	326	(269 - 394)	41	(34 - 49)	50	(41 - 60)	276	(221 - 343)
Montana	53	(45 - 63)	6	(5 - 7)	11	(9 - 13)	43	(34 - 52)

(continued)

Table 26 Had at Least One Major Depressive Episode in the Past Year, by Age Group and State: Estimated Numbers (in Thousands), Annual Averages Based on 2010 and 2011 NSDUHs (continued)

State	18+ (Estimate)	18+ (95% Confidence Interval)	12-17 (Estimate)	12-17 (95% Confidence Interval)	18-25 (Estimate)	18-25 (95% Confidence Interval)	26+ (Estimate)	26+ (95% Confidence Interval)
Nebraska	96	(80 - 115)	12	(10 - 15)	18	(15 - 21)	78	(63 - 97)
Nevada	125	(102 - 153)	18	(15 - 23)	21	(17 - 26)	104	(81 - 133)
New Hampshire	73	(60 - 89)	10	(8 - 12)	13	(11 - 16)	60	(47 - 76)
New Jersey	376	(305 - 463)	54	(44 - 66)	67	(55 - 82)	309	(240 - 396)
New Mexico	101	(82 - 123)	15	(12 - 18)	17	(14 - 21)	83	(65 - 106)
New York	983	(870 - 1,109)	108	(93 - 124)	173	(153 - 196)	810	(700 - 935)
North Carolina	445	(362 - 545)	54	(44 - 67)	77	(63 - 94)	368	(288 - 467)
North Dakota	32	(26 - 39)	4	(3 - 4)	8	(6 - 9)	24	(19 - 31)
Ohio	681	(608 - 761)	75	(66 - 85)	110	(99 - 123)	570	(500 - 650)
Oklahoma	201	(166 - 242)	24	(20 - 30)	36	(30 - 44)	165	(132 - 206)
Oregon	221	(182 - 267)	25	(21 - 30)	40	(33 - 48)	181	(144 - 226)
Pennsylvania	679	(604 - 764)	79	(69 - 90)	126	(111 - 144)	553	(480 - 636)
Rhode Island	56	(46 - 69)	6	(5 - 8)	11	(9 - 13)	45	(35 - 58)
South Carolina	258	(209 - 316)	29	(23 - 35)	39	(32 - 47)	219	(172 - 278)
South Dakota	37	(30 - 45)	6	(5 - 7)	8	(6 - 10)	29	(23 - 37)
Tennessee	340	(282 - 409)	41	(34 - 50)	52	(43 - 63)	287	(231 - 356)
Texas	1,059	(933 - 1,200)	174	(154 - 197)	217	(190 - 246)	842	(722 - 982)
Utah	160	(134 - 190)	22	(18 - 27)	36	(30 - 43)	124	(100 - 154)
Vermont	37	(31 - 45)	3	(3 - 4)	6	(5 - 8)	31	(25 - 39)
Virginia	413	(341 - 500)	52	(43 - 63)	77	(64 - 92)	336	(266 - 423)
Washington	433	(358 - 522)	46	(38 - 56)	65	(54 - 79)	367	(294 - 457)
West Virginia	112	(92 - 136)	11	(9 - 13)	15	(12 - 18)	97	(78 - 121)
Wisconsin	292	(240 - 356)	33	(27 - 41)	56	(46 - 68)	236	(185 - 299)
Wyoming	31	(26 - 37)	4	(3 - 5)	6	(5 - 7)	25	(20 - 31)

NOTE: Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. For details, see Section A.11 in Appendix A of the report on *State Estimates of Substance Use and Mental Disorders from the 2009-2010 NSDUHs*.

NOTE: There are minor wording differences in the questions in the adult and adolescent major depressive episode modules. Therefore, data from youths aged 12 to 17 were not combined with data from persons aged 18 or older to produce an estimate for those aged 12 or older.

NOTE: State and census region estimates, along with the 95 percent Bayesian confidence (credible) intervals, are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques. For the "Total U.S." row, design-based (direct) estimates and corresponding 95 percent confidence intervals are given.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011 (2010 Data – Revised March 2012).

