

WISCONSIN

Table 109 Selected Drug Use, Perceptions of Great Risk, Average Annual Marijuana Initiates, Past Year Substance Dependence or Abuse, Needing But Not Receiving Treatment, and Past Year Mental Health Measures in Wisconsin, by Age Group: Estimated Numbers (in Thousands), Annual Averages Based on 2010-2011 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLICIT DRUGS					
Past Month Illicit Drug Use ¹	369	42	120	208	327
Past Year Marijuana Use	454	54	170	230	400
Past Month Marijuana Use	274	28	99	146	245
Past Month Use of Illicit Drugs Other Than Marijuana ¹	168	19	53	96	149
Past Year Cocaine Use	70	4	29	38	67
Past Year Nonmedical Pain Reliever Use	214	27	67	120	187
Perception of Great Risk of Smoking Marijuana Once a Month	1,390	139	100	1,151	1,251
Average Annual Number of Marijuana Initiates ²	44	21	20	3	23
ALCOHOL					
Past Month Alcohol Use	2,965	62	438	2,465	2,903
Past Month Binge Alcohol Use ³	1,289	34	294	960	1,255
Perception of Great Risk of Drinking Five or More Drinks Once or Twice a Week	1,690	158	178	1,355	1,532
Past Month Alcohol Use (Persons Aged 12 to 20)	189 ⁴	--	--	--	--
Past Month Binge Alcohol Use (Persons Aged 12 to 20) ³	123 ⁴	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁵	1,283	43	272	968	1,239
Past Month Cigarette Use	1,041	36	224	781	1,005
Perception of Great Risk of Smoking One or More Packs of Cigarettes per Day	3,280	293	386	2,601	2,987
PAST YEAR DEPENDENCE, ABUSE, AND TREATMENT⁶					
Illicit Drug Dependence ¹	81	10	27	45	72
Illicit Drug Dependence or Abuse ¹	114	19	39	55	94
Alcohol Dependence	131	6	34	91	125
Alcohol Dependence or Abuse	324	16	95	212	308
Alcohol or Illicit Drug Dependence or Abuse ¹	409	29	115	265	380
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,7}	103	18	36	48	85
Needing But Not Receiving Treatment for Alcohol Use ⁷	315	16	93	206	299
PAST YEAR MENTAL HEALTH					
Had at Least One Major Depressive Episode ⁸	--	33	56	236	292
Serious Mental Illness ⁹	--	--	24	148	172
Any Mental Illness ¹⁰	--	--	112	641	753
Had Serious Thoughts of Suicide	--	--	43	135	178

-- Not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach.

NOTE: Estimates for serious mental illness (SMI) and any mental illness (AMI) presented in this table may differ from previously published SMI and AMI estimates because of revised estimation procedures. For details, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates include data from original methamphetamine questions but do not include new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Average annual number of marijuana initiates = $X_1 \div 2$, where X_1 is the number of marijuana initiates in the past 24 months.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

⁴ Underage drinking is defined for persons aged 12 to 20; therefore, the "Total" estimate reflects that age group and not persons aged 12 or older.

⁵ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁶ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁷ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs (or alcohol), but not receiving treatment for an illicit drug (or alcohol) problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

⁸ Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from persons aged 18 or older to produce an estimate for those aged 12 or older. For more details, see Section A.11 in Appendix A of the report on *State Estimates of Substance Use and Mental Disorders from the 2009-2010 NSDUHs*.

⁹ Serious mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in serious functional impairment.

¹⁰ Any mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 (Revised March 2012; SMI and AMI Estimates Revised October 2013) and 2011 (SMI and AMI Estimates Revised October 2013).

WISCONSIN

Table 110 Selected Drug Use, Perceptions of Great Risk, Average Annual Rates of First Use of Marijuana, Past Year Substance Dependence or Abuse, Needing But Not Receiving Treatment, and Past Year Mental Health Measures in Wisconsin, by Age Group: Percentages, Annual Averages Based on 2010-2011 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLCIT DRUGS					
Past Month Illicit Drug Use ¹	7.77	9.34	18.85	5.67	7.61
Past Year Marijuana Use	9.57	12.18	26.81	6.27	9.30
Past Month Marijuana Use	5.76	6.36	15.57	3.99	5.70
Past Month Use of Illicit Drugs Other Than Marijuana ¹	3.55	4.28	8.40	2.62	3.47
Past Year Cocaine Use	1.48	0.86	4.52	1.04	1.55
Past Year Nonmedical Pain Reliever Use	4.51	6.09	10.55	3.27	4.35
Perception of Great Risk of Smoking Marijuana Once a Month	29.28	31.14	15.68	31.41	29.09
Average Annual Rate of First Use of Marijuana ²	1.53	5.06	5.89	0.14	0.95
ALCOHOL					
Past Month Alcohol Use	62.48	13.94	68.92	67.25	67.49
Past Month Binge Alcohol Use ³	27.16	7.69	46.36	26.19	29.17
Perception of Great Risk of Drinking Five or More Drinks Once or Twice a Week	35.62	35.55	27.95	36.95	35.62
Past Month Alcohol Use (Persons Aged 12 to 20)	27.84 ⁴	--	--	--	--
Past Month Binge Alcohol Use (Persons Aged 12 to 20) ³	18.09 ⁴	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁵	27.03	9.76	42.75	26.40	28.81
Past Month Cigarette Use	21.93	8.10	35.25	21.30	23.35
Perception of Great Risk of Smoking One or More Packs of Cigarettes per Day	69.12	65.85	60.80	70.96	69.46
PAST YEAR DEPENDENCE, ABUSE, AND TREATMENT⁶					
Illicit Drug Dependence ¹	1.71	2.15	4.25	1.22	1.67
Illicit Drug Dependence or Abuse ¹	2.40	4.36	6.12	1.51	2.19
Alcohol Dependence	2.76	1.35	5.34	2.49	2.91
Alcohol Dependence or Abuse	6.83	3.67	15.01	5.80	7.16
Alcohol or Illicit Drug Dependence or Abuse ¹	8.61	6.48	18.13	7.22	8.83
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,7}	2.17	4.15	5.72	1.32	1.97
Needing But Not Receiving Treatment for Alcohol Use ⁷	6.63	3.63	14.57	5.62	6.94
PAST YEAR MENTAL HEALTH					
Had at Least One Major Depressive Episode ⁸	--	7.49	8.86	6.44	6.80
Serious Mental Illness ⁹	--	--	3.91	4.02	4.01
Any Mental Illness ¹⁰	--	--	18.20	17.42	17.53
Had Serious Thoughts of Suicide	--	--	6.72	3.68	4.13

-- Not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach.

NOTE: Estimates for serious mental illness (SMI) and any mental illness (AMI) presented in this table may differ from previously published SMI and AMI estimates because of revised estimation procedures. For details, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates include data from original methamphetamine questions but do not include new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.² Average annual marijuana initiation rate = $100 * \{ [X_1 + (0.5 * X_1 + X_2)] / 2 \}$, where X_1 is the number of marijuana initiates in the past 24 months and X_2 is the number of persons who never used marijuana. Both of the computation components, X_1 and X_2 , are based on a survey-weighted hierarchical Bayes estimation approach. Note that the age group is based on a respondent's age at the time of the interview, not his or her age at first use.³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.⁴ Underage drinking is defined for persons aged 12 to 20; therefore, the "Total" estimate reflects that age group and not persons aged 18 or older.⁵ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.⁶ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).⁷ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs (or alcohol), but not receiving treatment for an illicit drug (or alcohol) problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).⁸ Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from persons aged 18 or older to produce an estimate for those aged 12 or older. For more details, see Section A.11 in Appendix A of the report on *State Estimates of Substance Use and Mental Disorders from the 2009-2010 NSDUHs*.⁹ Serious mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in serious functional impairment.¹⁰ Any mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 (Revised March 2012; SMI and AMI Estimates Revised October 2013) and 2011 (SMI and AMI Estimates Revised October 2013).