

UNITED STATES

Table 1 Selected Drug Use, Perceptions of Great Risk, Average Annual Marijuana Initiates, Past Year Substance Dependence or Abuse, Needing But Not Receiving Treatment, and Past Year Mental Health Measures in the *United States*, by Age Group: Estimated Numbers (in Thousands), Annual Averages Based on 2011-2012 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLICIT DRUGS					
Past Month Illicit Drug Use ¹	23,159	2,449	7,367	13,343	20,710
Past Year Marijuana Use	30,627	3,458	10,721	16,448	27,169
Past Month Marijuana Use	18,463	1,884	6,505	10,074	16,580
Past Month Use of Illicit Drugs Other Than Marijuana ¹	8,452	972	2,410	5,070	7,480
Past Year Cocaine Use	4,264	204	1,582	2,478	4,059
Past Year Nonmedical Pain Reliever Use	11,816	1,408	3,430	6,978	10,408
Perception of Great Risk of Smoking Marijuana Once a Month	81,246	6,743	5,984	68,520	74,503
Average Annual Number of Marijuana Initiates ²	2,882	1,309	1,359	214	1,573
ALCOHOL					
Past Month Alcohol Use	134,447	3,272	20,821	110,354	131,175
Past Month Binge Alcohol Use ³	59,015	1,824	13,657	43,533	57,191
Perception of Great Risk of Drinking Five or More Drinks Once or Twice a Week	109,348	10,034	11,885	87,428	99,313
Past Month Alcohol Use (Persons Aged 12 to 20)	9,475 ⁴	--	--	--	--
Past Month Binge Alcohol Use (Persons Aged 12 to 20) ³	5,964 ⁴	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁵	68,853	2,321	13,349	53,183	66,532
Past Month Cigarette Use	57,165	1,786	11,247	44,131	55,379
Perception of Great Risk of Smoking One or More Packs of Cigarettes per Day	184,883	16,450	22,824	145,609	168,433
PAST YEAR DEPENDENCE, ABUSE, AND TREATMENT⁶					
Illicit Drug Dependence ¹	4,811	565	1,870	2,376	4,246
Illicit Drug Dependence or Abuse ¹	6,921	1,076	2,640	3,205	5,845
Alcohol Dependence	8,192	335	2,069	5,788	7,857
Alcohol Dependence or Abuse	17,193	901	4,946	11,346	16,292
Alcohol or Illicit Drug Dependence or Abuse ¹	21,395	1,618	6,452	13,325	19,777
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,7}	6,169	990	2,422	2,758	5,180
Needing But Not Receiving Treatment for Alcohol Use ⁷	16,359	864	4,802	10,692	15,495
PAST YEAR MENTAL HEALTH					
Had at Least One Major Depressive Episode ^{8,9}	--	2,161	2,965	12,752	15,716
Serious Mental Illness ^{9,10}	--	--	1,360	7,929	9,289
Any Mental Illness ^{9,11}	--	--	6,565	35,981	42,546
Had Serious Thoughts of Suicide	--	--	2,422	6,396	8,818

-- Not available.

NOTE: Numbers in this table are based on national design-based estimates.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates include data from original methamphetamine questions but do not include new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Average annual number of marijuana initiates = $X_1 \div 2$, where X_1 is the number of marijuana initiates in the past 24 months.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

⁴ Underage drinking is defined for persons aged 12 to 20; therefore, the "12+" estimate reflects that age group and not persons aged 12 or older.

⁵ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁶ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

⁷ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs (or alcohol), but not receiving treatment for an illicit drug (or alcohol) problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

⁸ Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*, which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from persons aged 18 or older to produce an estimate for those aged 12 or older.

⁹ For details, see Section B of the "2011-2012 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at <http://www.samhsa.gov/data/NSDUH/2k12State/NSDUHsae2012/Index.aspx>.

¹⁰ Serious mental illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in serious functional impairment.

¹¹ Any mental illness (AMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 (SMI and AMI Estimates Revised October 2013) and 2012.

UNITED STATES

Table 2 Selected Drug Use, Perceptions of Great Risk, Average Annual Rates of First Use of Marijuana, Past Year Substance Dependence or Abuse, Needing But Not Receiving Treatment, and Past Year Mental Health Measures in the *United States*, by Age Group: Percentages, Annual Averages Based on 2011-2012 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLICIT DRUGS					
Past Month Illicit Drug Use ¹	8.95	9.82	21.39	6.69	8.85
Past Year Marijuana Use	11.83	13.86	31.12	8.25	11.62
Past Month Marijuana Use	7.13	7.55	18.89	5.05	7.09
Past Month Use of Illicit Drugs Other Than Marijuana ¹	3.27	3.90	7.00	2.54	3.20
Past Year Cocaine Use	1.65	0.82	4.59	1.24	1.74
Past Year Nonmedical Pain Reliever Use	4.57	5.64	9.96	3.50	4.45
Perception of Great Risk of Smoking Marijuana Once a Month	31.37	27.02	17.37	34.36	31.83
Average Annual Rate of First Use of Marijuana ²	1.89	5.96	7.60	0.19	1.21
ALCOHOL					
Past Month Alcohol Use	51.94	13.11	60.45	55.33	56.09
Past Month Binge Alcohol Use ³	22.80	7.31	39.65	21.83	24.45
Perception of Great Risk of Drinking Five or More Drinks Once or Twice a Week	42.24	40.21	34.50	43.84	42.46
Past Month Alcohol Use (Persons Aged 12 to 20)	24.70 ⁴	--	--	--	--
Past Month Binge Alcohol Use (Persons Aged 12 to 20) ³	15.55 ⁴	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁵	26.60	9.30	38.75	26.67	28.45
Past Month Cigarette Use	22.09	7.16	32.65	22.13	23.68
Perception of Great Risk of Smoking One or More Packs of Cigarettes per Day	71.43	65.92	66.26	73.01	72.01
PAST YEAR DEPENDENCE, ABUSE, AND TREATMENT⁶					
Illicit Drug Dependence ¹	1.86	2.27	5.43	1.19	1.82
Illicit Drug Dependence or Abuse ¹	2.67	4.31	7.66	1.61	2.50
Alcohol Dependence	3.16	1.34	6.01	2.90	3.36
Alcohol Dependence or Abuse	6.64	3.61	14.36	5.69	6.97
Alcohol or Illicit Drug Dependence or Abuse ¹	8.27	6.48	18.73	6.68	8.46
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,7}	2.38	3.97	7.03	1.38	2.21
Needing But Not Receiving Treatment for Alcohol Use ⁷	6.32	3.46	13.94	5.36	6.62
PAST YEAR MENTAL HEALTH					
Had at Least One Major Depressive Episode ^{8,9}	--	8.66	8.61	6.39	6.72
Serious Mental Illness ^{9,10}	--	--	3.95	3.98	3.97
Any Mental Illness ^{9,11}	--	--	19.06	18.04	18.19
Had Serious Thoughts of Suicide	--	--	7.03	3.21	3.77

-- Not available.

NOTE: The percentages in this table are based on national design-based estimates.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates include data from original methamphetamine questions but do not include new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Average annual marijuana initiation rate = $100 * \{ [X_1 \div (0.5 * X_1 + X_2)] \div 2 \}$, where X_1 is the number of marijuana initiates in the past 24 months and X_2 is the number of persons who never used marijuana. Both of the computation components, X_1 and X_2 , are based on a survey-weighted hierarchical Bayes estimation approach. Note that the age group is based on a respondent's age at the time of the interview, not his or her age at first use.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

⁴ Underage drinking is defined for persons aged 12 to 20; therefore, the "12+" estimate reflects that age group and not persons aged 12 or older.

⁵ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁶ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁷ Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs (or alcohol), but not receiving treatment for an illicit drug (or alcohol) problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

⁸ Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from persons aged 18 or older to produce an estimate for those aged 12 or older.

⁹ For details, see Section B of the "2011-2012 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at <http://www.samhsa.gov/data/NSDUH/2k12State/NSDUHsae2012/Index.aspx>.

¹⁰ Serious mental illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in serious functional impairment.

¹¹ Any mental illness (AMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2011 (SMI and AMI Estimates Revised October 2013) and 2012.