

States Categorized into Five Groups from Lowest to Highest Estimate, by Age Group: Annual Averages Based on 2012 and 2013 NSDUHs						
Table	Age Group	Group 1	Group 2	Group 3	Group 4	Group 5
Table 1. Illicit Drug Use in the Past Month	12 to 17	6.08 - 7.73	7.74 - 8.23	8.24 - 9.63	9.64 - 11.05	11.06 - 15.12
	18 to 25	13.79 - 17.33	17.34 - 19.76	19.77 - 22.21	22.22 - 24.50	24.51 - 31.89
	26 or Older	4.36 - 5.31	5.32 - 6.04	6.05 - 7.48	7.49 - 9.03	9.04 - 12.75
	18 or Older	5.95 - 7.10	7.11 - 7.89	7.90 - 9.54	9.55 - 11.32	11.33 - 15.82
	12 or Older	6.02 - 7.17	7.18 - 7.94	7.95 - 9.59	9.60 - 11.28	11.29 - 15.76
Table 2. Marijuana Use in the Past Year	12 to 17	9.79 - 11.39	11.40 - 12.62	12.63 - 14.25	14.26 - 16.02	16.03 - 20.28
	18 to 25	20.52 - 26.23	26.24 - 30.82	30.83 - 32.06	32.07 - 35.88	35.89 - 46.77
	26 or Older	5.08 - 6.81	6.82 - 7.57	7.58 - 9.47	9.48 - 11.72	11.73 - 17.24
	18 or Older	8.01 - 9.66	9.67 - 10.91	10.92 - 12.71	12.72 - 15.12	15.13 - 21.20
	12 or Older	8.21 - 9.92	9.93 - 11.05	11.06 - 12.81	12.82 - 15.39	15.40 - 21.02
Table 3. Marijuana Use in the Past Month	12 to 17	4.81 - 5.35	5.36 - 6.25	6.26 - 7.52	7.53 - 9.15	9.16 - 12.95
	18 to 25	10.91 - 14.43	14.44 - 17.03	17.04 - 19.09	19.10 - 23.04	23.05 - 29.79
	26 or Older	2.90 - 3.67	3.68 - 4.36	4.37 - 5.79	5.80 - 7.23	7.24 - 11.18
	18 or Older	4.33 - 5.25	5.26 - 6.16	6.17 - 7.77	7.78 - 9.80	9.81 - 14.18
	12 or Older	4.41 - 5.31	5.32 - 6.21	6.22 - 7.71	7.72 - 9.79	9.80 - 14.08
Table 4. Perceptions of Great Risk of Smoking Marijuana Once a Month	12 to 17	19.06 - 21.52	21.53 - 23.57	23.58 - 26.39	26.40 - 29.01	29.02 - 35.29
	18 to 25	7.87 - 11.53	11.54 - 14.24	14.25 - 15.64	15.65 - 17.30	17.31 - 24.43
	26 or Older	19.90 - 24.85	24.86 - 29.08	29.09 - 31.89	31.90 - 36.13	36.14 - 43.53
	18 or Older	18.66 - 22.54	22.55 - 26.90	26.91 - 29.40	29.41 - 33.41	33.42 - 40.01
	12 or Older	18.70 - 22.42	22.43 - 26.62	26.63 - 29.18	29.19 - 32.94	32.95 - 39.26
Table 5. First Use of Marijuana	12 to 17	3.53 - 4.86	4.87 - 5.50	5.51 - 6.12	6.13 - 6.83	6.84 - 9.07
	18 to 25	4.14 - 6.10	6.11 - 6.97	6.98 - 7.76	7.77 - 8.75	8.76 - 13.17
	26 or Older ¹	--	--	--	--	--
	18 or Older	0.88 - 1.04	1.05 - 1.14	1.15 - 1.27	1.28 - 1.45	1.46 - 2.35
	12 or Older	1.37 - 1.61	1.62 - 1.74	1.75 - 1.98	1.99 - 2.27	2.28 - 3.05
Table 6. Illicit Drug Use Other Than Marijuana in the Past Month	12 to 17	2.39 - 2.73	2.74 - 3.00	3.01 - 3.45	3.46 - 3.86	3.87 - 5.40
	18 to 25	5.04 - 6.02	6.03 - 6.77	6.78 - 7.04	7.05 - 7.58	7.59 - 10.04
	26 or Older	1.73 - 2.12	2.13 - 2.44	2.45 - 2.65	2.66 - 3.27	3.28 - 5.46
	18 or Older	2.22 - 2.78	2.79 - 3.10	3.11 - 3.42	3.43 - 3.85	3.86 - 5.79
	12 or Older	2.25 - 2.79	2.80 - 3.10	3.11 - 3.40	3.41 - 3.86	3.87 - 5.77
Table 7. Cocaine Use in the Past Year	12 to 17	0.33 - 0.49	0.50 - 0.54	0.55 - 0.61	0.62 - 0.69	0.70 - 1.13
	18 to 25	2.59 - 3.34	3.35 - 3.72	3.73 - 4.24	4.25 - 5.32	5.33 - 8.14
	26 or Older	0.83 - 1.04	1.05 - 1.16	1.17 - 1.29	1.30 - 1.50	1.51 - 3.08
	18 or Older	1.12 - 1.40	1.41 - 1.55	1.56 - 1.73	1.74 - 2.07	2.08 - 3.46
	12 or Older	1.06 - 1.31	1.32 - 1.46	1.47 - 1.63	1.64 - 1.96	1.97 - 3.28

States Categorized into Five Groups from Lowest to Highest Estimate, by Age Group: Annual Averages Based on 2012 and 2013 NSDUHs						
Table	Age Group	Group 1	Group 2	Group 3	Group 4	Group 5
Table 8. Nonmedical Use of Pain Relievers in the Past Year	12 to 17	3.54 - 4.42	4.43 - 4.64	4.65 - 5.31	5.32 - 5.71	5.72 - 7.15
	18 to 25	7.59 - 8.22	8.23 - 9.06	9.07 - 9.81	9.82 - 10.65	10.66 - 12.23
	26 or Older	2.71 - 2.90	2.91 - 3.24	3.25 - 3.76	3.77 - 4.11	4.12 - 4.75
	18 or Older	3.51 - 3.75	3.76 - 4.16	4.17 - 4.60	4.61 - 5.05	5.06 - 5.68
	12 or Older	3.56 - 3.80	3.81 - 4.18	4.19 - 4.66	4.67 - 5.13	5.14 - 5.82
Table 9. Alcohol Use in the Past Month	12 to 17	7.82 - 11.22	11.23 - 11.69	11.70 - 12.37	12.38 - 13.10	13.11 - 15.12
	18 to 25	34.46 - 56.25	56.26 - 58.95	58.96 - 63.46	63.47 - 66.54	66.55 - 73.46
	26 or Older	34.65 - 50.92	50.93 - 54.97	54.98 - 59.15	59.16 - 62.32	62.33 - 69.63
	18 or Older	34.62 - 51.68	51.69 - 55.81	55.82 - 59.51	59.52 - 62.98	62.99 - 70.11
	12 or Older	31.30 - 47.57	47.58 - 51.70	51.71 - 55.15	55.16 - 58.82	58.83 - 65.19
Table 10. Binge Alcohol Use in the Past Month	12 to 17	5.47 - 6.35	6.36 - 6.58	6.59 - 7.00	7.01 - 7.37	7.38 - 8.67
	18 to 25	26.16 - 35.68	35.69 - 38.88	38.89 - 41.17	41.18 - 45.13	45.14 - 53.75
	26 or Older	15.78 - 20.04	20.05 - 21.50	21.51 - 23.06	23.07 - 24.69	24.70 - 32.16
	18 or Older	17.73 - 22.29	22.30 - 23.98	23.99 - 26.14	26.15 - 27.36	27.37 - 35.32
	12 or Older	16.25 - 21.02	21.03 - 22.29	22.30 - 24.37	24.38 - 25.41	25.42 - 33.65
Table 11. Perceptions of Great Risk of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week	12 to 17	30.01 - 35.63	35.64 - 37.45	37.46 - 38.59	38.60 - 41.49	41.50 - 47.67
	18 to 25	21.29 - 27.28	27.29 - 30.32	30.33 - 32.82	32.83 - 36.69	36.70 - 39.87
	26 or Older	35.56 - 38.03	38.04 - 40.76	40.77 - 43.07	43.08 - 45.44	45.45 - 48.44
	18 or Older	33.94 - 36.58	36.59 - 38.90	38.91 - 41.60	41.61 - 43.71	43.72 - 46.76
	12 or Older	34.09 - 36.50	36.51 - 38.67	38.68 - 40.98	40.99 - 43.58	43.59 - 46.25
Table 12a. Underage Alcohol Use in the Past Month	12 to 20	14.57 - 20.84	20.85 - 22.94	22.95 - 24.04	24.05 - 26.38	26.39 - 35.72
Table 12b. Underage Binge Alcohol Use in the Past Month	12 to 20	11.21 - 13.24	13.25 - 14.42	14.43 - 15.54	15.55 - 17.57	17.58 - 23.21
Table 13. Tobacco Product Use in the Past Month	12 to 17	5.66 - 7.27	7.28 - 8.59	8.60 - 9.59	9.60 - 10.63	10.64 - 12.84
	18 to 25	25.94 - 35.13	35.14 - 38.69	38.70 - 42.18	42.19 - 45.00	45.01 - 48.94
	26 or Older	18.90 - 24.01	24.02 - 26.87	26.88 - 28.46	28.47 - 32.17	32.18 - 38.66
	18 or Older	20.51 - 25.55	25.56 - 28.68	28.69 - 30.09	30.10 - 33.53	33.54 - 40.01
	12 or Older	19.05 - 23.93	23.94 - 26.94	26.95 - 28.20	28.21 - 31.30	31.31 - 37.76
Table 14. Cigarette Use in the Past Month	12 to 17	4.31 - 5.40	5.41 - 6.25	6.26 - 7.14	7.15 - 7.72	7.73 - 9.50
	18 to 25	23.60 - 29.28	29.29 - 31.53	31.54 - 34.10	34.11 - 35.68	35.69 - 40.59
	26 or Older	16.08 - 19.96	19.97 - 22.25	22.26 - 23.59	23.60 - 26.25	26.26 - 32.06
	18 or Older	17.61 - 21.35	21.36 - 23.44	23.45 - 24.80	24.81 - 27.74	27.75 - 33.17
	12 or Older	16.29 - 19.92	19.93 - 21.65	21.66 - 23.21	23.22 - 25.74	25.75 - 30.62
Table 15. Perceptions of Great Risk of Smoking One or More Packs of Cigarettes per Day	12 to 17	59.08 - 61.69	61.70 - 63.41	63.42 - 64.51	64.52 - 65.93	65.94 - 70.43
	18 to 25	56.64 - 61.44	61.45 - 63.12	63.13 - 65.31	65.32 - 67.96	67.97 - 72.60
	26 or Older	63.78 - 67.79	67.80 - 69.93	69.94 - 72.49	72.50 - 74.69	74.70 - 78.20
	18 or Older	62.87 - 66.45	66.46 - 68.82	68.83 - 71.59	71.60 - 73.62	73.63 - 77.35
	12 or Older	62.60 - 65.90	65.91 - 68.41	68.42 - 70.87	70.88 - 72.82	72.83 - 76.19

States Categorized into Five Groups from Lowest to Highest Estimate, by Age Group: Annual Averages Based on 2012 and 2013 NSDUHs						
Table	Age Group	Group 1	Group 2	Group 3	Group 4	Group 5
Table 16. Alcohol Dependence or Abuse in the Past Year	12 to 17	2.34 - 2.85	2.86 - 3.10	3.11 - 3.28	3.29 - 3.56	3.57 - 4.91
	18 to 25	11.04 - 12.42	12.43 - 13.46	13.47 - 14.50	14.51 - 15.99	16.00 - 19.17
	26 or Older	4.43 - 5.35	5.36 - 5.85	5.86 - 6.33	6.34 - 6.71	6.72 - 11.27
	18 or Older	5.74 - 6.36	6.37 - 6.95	6.96 - 7.48	7.49 - 7.97	7.98 - 12.52
	12 or Older	5.36 - 6.06	6.07 - 6.59	6.60 - 7.09	7.10 - 7.60	7.61 - 11.96
Table 17. Alcohol Dependence in the Past Year	12 to 17	0.89 - 1.03	1.04 - 1.08	1.09 - 1.19	1.20 - 1.26	1.27 - 1.71
	18 to 25	4.32 - 5.07	5.08 - 5.64	5.65 - 5.93	5.94 - 6.65	6.66 - 8.95
	26 or Older	2.54 - 2.79	2.80 - 2.95	2.96 - 3.09	3.10 - 3.28	3.29 - 4.56
	18 or Older	2.88 - 3.19	3.20 - 3.30	3.31 - 3.53	3.54 - 3.79	3.80 - 5.36
	12 or Older	2.63 - 2.99	3.00 - 3.09	3.10 - 3.29	3.30 - 3.54	3.55 - 5.12
Table 18. Illicit Drug Dependence or Abuse in the Past Year	12 to 17	2.44 - 3.15	3.16 - 3.50	3.51 - 3.82	3.83 - 4.13	4.14 - 5.12
	18 to 25	5.67 - 6.73	6.74 - 7.08	7.09 - 7.79	7.80 - 8.37	8.38 - 9.59
	26 or Older	1.26 - 1.45	1.46 - 1.67	1.68 - 1.86	1.87 - 2.08	2.09 - 2.84
	18 or Older	2.02 - 2.18	2.19 - 2.54	2.55 - 2.75	2.76 - 2.91	2.92 - 3.85
	12 or Older	2.14 - 2.34	2.35 - 2.64	2.65 - 2.83	2.84 - 3.02	3.03 - 3.87
Table 19. Illicit Drug Dependence in the Past Year	12 to 17	1.46 - 1.74	1.75 - 1.86	1.87 - 2.01	2.02 - 2.11	2.12 - 2.47
	18 to 25	3.67 - 4.66	4.67 - 5.08	5.09 - 5.63	5.64 - 6.04	6.05 - 7.35
	26 or Older	0.93 - 1.11	1.12 - 1.26	1.27 - 1.35	1.36 - 1.47	1.48 - 2.39
	18 or Older	1.34 - 1.64	1.65 - 1.83	1.84 - 1.97	1.98 - 2.10	2.11 - 3.07
	12 or Older	1.35 - 1.65	1.66 - 1.85	1.86 - 1.96	1.97 - 2.10	2.11 - 3.03
Table 20. Dependence or Abuse of Illicit Drugs or Alcohol in the Past Year	12 to 17	4.56 - 5.13	5.14 - 5.67	5.68 - 5.83	5.84 - 6.38	6.39 - 7.20
	18 to 25	14.67 - 16.52	16.53 - 18.07	18.08 - 19.25	19.26 - 20.15	20.16 - 23.66
	26 or Older	5.66 - 6.67	6.68 - 7.06	7.07 - 7.34	7.35 - 7.93	7.94 - 12.60
	18 or Older	7.43 - 8.20	8.21 - 8.65	8.66 - 9.09	9.10 - 9.65	9.66 - 14.29
	12 or Older	7.07 - 7.89	7.90 - 8.33	8.34 - 8.77	8.78 - 9.34	9.35 - 13.84
Table 21. Needing But Not Receiving Treatment for Illicit Drug Use in the Past Year	12 to 17	2.35 - 2.94	2.95 - 3.17	3.18 - 3.50	3.51 - 3.81	3.82 - 4.77
	18 to 25	5.19 - 6.05	6.06 - 6.61	6.62 - 7.10	7.11 - 7.46	7.47 - 8.61
	26 or Older	1.10 - 1.26	1.27 - 1.45	1.46 - 1.56	1.57 - 1.79	1.80 - 2.34
	18 or Older	1.81 - 1.95	1.96 - 2.23	2.24 - 2.42	2.43 - 2.51	2.52 - 3.34
	12 or Older	1.87 - 2.09	2.10 - 2.31	2.32 - 2.49	2.50 - 2.60	2.61 - 3.38
Table 22. Needing But Not Receiving Treatment for Alcohol Use in the Past Year	12 to 17	2.29 - 2.74	2.75 - 2.96	2.97 - 3.15	3.16 - 3.32	3.33 - 4.35
	18 to 25	10.29 - 12.38	12.39 - 13.18	13.19 - 14.05	14.06 - 15.44	15.45 - 18.66
	26 or Older	4.27 - 5.22	5.23 - 5.47	5.48 - 5.94	5.95 - 6.25	6.26 - 10.65
	18 or Older	5.38 - 6.17	6.18 - 6.59	6.60 - 7.08	7.09 - 7.63	7.64 - 11.92
	12 or Older	5.05 - 5.89	5.90 - 6.24	6.25 - 6.67	6.68 - 7.22	7.23 - 11.38

States Categorized into Five Groups from Lowest to Highest Estimate, by Age Group: Annual Averages Based on 2012 and 2013 NSDUHs

Table	Age Group	Group 1	Group 2	Group 3	Group 4	Group 5
Table 23. Serious Mental Illness (SMI) in the Past Year	18 to 25	3.46 - 3.92	3.93 - 4.15	4.16 - 4.45	4.46 - 4.76	4.77 - 5.77
	26 or Older	3.16 - 3.83	3.84 - 4.16	4.17 - 4.40	4.41 - 4.84	4.85 - 5.56
	18 or Older	3.26 - 3.88	3.89 - 4.16	4.17 - 4.36	4.37 - 4.76	4.77 - 5.48
Table 24. Any Mental Illness (AMI) in the Past Year	18 to 25	16.43 - 18.30	18.31 - 19.17	19.18 - 20.00	20.01 - 20.78	20.79 - 26.90
	26 or Older	15.13 - 17.47	17.48 - 18.12	18.13 - 19.24	19.25 - 19.89	19.90 - 22.05
	18 or Older	15.62 - 17.64	17.65 - 18.38	18.39 - 19.33	19.34 - 20.00	20.01 - 22.31
Table 25. Had Serious Thoughts of Suicide in the Past Year	18 to 25	5.83 - 6.81	6.82 - 7.32	7.33 - 7.68	7.69 - 7.88	7.89 - 9.39
	26 or Older	2.86 - 3.15	3.16 - 3.32	3.33 - 3.44	3.45 - 3.75	3.76 - 4.26
	18 or Older	3.51 - 3.76	3.77 - 3.93	3.94 - 4.07	4.08 - 4.33	4.34 - 4.84
Table 26. Had at Least One Major Depressive Episode (MDE) in the Past Year	12 to 17	7.38 - 9.06	9.07 - 9.44	9.45 - 9.82	9.83 - 10.48	10.49 - 12.65
	18 to 25	7.44 - 8.19	8.20 - 8.56	8.57 - 9.15	9.16 - 9.81	9.82 - 12.90
	26 or Older	5.15 - 5.96	5.97 - 6.44	6.45 - 6.72	6.73 - 7.01	7.02 - 8.66
	18 or Older	5.57 - 6.34	6.35 - 6.75	6.76 - 7.08	7.09 - 7.35	7.36 - 9.45

-- Not available.

NOTE: This table shows the boundaries used to create the maps shown in "2012-2013 NSDUH National Maps of Prevalence Estimates, by State" at <http://www.samhsa.gov/data/population-data-nsduh/reports?tab=33>. Maps for the 18 or older age group are available only for the following measures: serious mental illness (SMI), any mental illness (AMI), serious thoughts of suicide, and depression (i.e., major depressive episode or MDE).

NOTE: State estimates (which are based on a survey-weighted hierarchical Bayes estimation approach) fall into one of five groups according to their ranking by quintiles. Because 51 areas are to be ranked for each measure, the middle quintile was assigned 11 areas and the remaining groups 10 each. In some cases, a "quintile" could have more or fewer States than desired because two (or more) States have the same estimate (to two decimal places). When such ties occurred at the "boundary" between two quintiles, all States with the same estimate were assigned to the lower quintile. Those States with the highest estimates for a given measure are in red, with the exception of the perceptions of risk measures, for which the lowest perceptions of great risk are in red. Those States with the lowest estimates are in white, with the exception of the perceptions of risk measures, for which the highest perceptions of great risk are in white. The upper and lower limits of each quintile shown in the map legend collectively define a continuum and are not necessarily the actual values of a particular area. Tables containing specific estimates for these States are available on the 2012-2013 NSDUH State Web page at <http://www.samhsa.gov/data/population-data-nsduh/reports?tab=33>.

¹The average annual incidence estimates for first use of marijuana were low for the 26 or older age group and had an abbreviated range; therefore, no State grouping has been made for this age group.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.