


2015-2016 National Survey on Drug Use and Health National Maps of Prevalence Estimates, by State

The 112 national maps presented here show the 2015-2016 National Survey on Drug Use and Health (NSDUH) estimates for 29 substance use and mental health outcomes, by age group, for 50 states and the District of Columbia. Each web link in the table of contents can be used to navigate to all of the maps (for all of the age groups) for the outcome listed in that row. The color of each state on the U.S. maps indicates how the state ranks relative to other states for each measure. States could fall into one of five groups according to their ranking by quintiles. Because 51 states were ranked for each measure, the middle quintile was assigned to 11 states, and the remaining quintiles were assigned 10 states each. In some cases, a "quintile" could have more or fewer states than desired because two (or more) states had the same estimate (to two decimal places). When such ties occurred at the "boundary" between two quintiles, all of the states with the same estimate were conservatively assigned to the lower quintile. Those states with the highest rates for a given measure are in red. Those states with the lowest estimates are in white. The upper and lower limits of each quintile shown in the map legend collectively define a continuum and are not necessarily the actual values of a particular state. For example, in [Figure 1a](#), the values on the boundary in the lowest quintile correspond to Iowa (6.76 percent) and New Jersey (8.13 percent) and are displayed in the legend. In the next to lowest quintile, Virginia (8.28 percent) and South Dakota (9.06 percent) are the states with the lowest and highest values; however, in the continuum of the legend, the lower limit was assigned a value of 8.14 percent because the upper limit of the quintile below it is 8.13 percent.


Tables containing specific estimates for these states are available on the 2015-2016 NSDUH web page at <https://www.samhsa.gov/data/>.

Figure 1a *Illicit Drug Use in the Past Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 1b *Illicit Drug Use in the Past Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 1c *Illicit Drug Use in the Past Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 1d *Illicit Drug Use in the Past Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 2a *Marijuana Use in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 2b *Marijuana Use in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 2c *Marijuana Use in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 2d *Marijuana Use in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 3a *Marijuana Use in the Past Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 3b *Marijuana Use in the Past Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 3c *Marijuana Use in the Past Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 3d *Marijuana Use in the Past Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 4a *Perceptions of Great Risk from Smoking Marijuana Once a Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 4b *Perceptions of Great Risk from Smoking Marijuana Once a Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 4c *Perceptions of Great Risk from Smoking Marijuana Once a Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 4d *Perceptions of Great Risk from Smoking Marijuana Once a Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 5a *First Use of Marijuana among Individuals Aged 12 or Older, by State: Average Annual Incidence Estimates (Expressed as Percentages of the At-Risk Population) Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 5b *First Use of Marijuana among Youths Aged 12 to 17, by State: Average Annual Incidence Estimates (Expressed as Percentages of the At-Risk Population) Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 5c *First Use of Marijuana among Adults Aged 18 to 25, by State: Average Annual Incidence Estimates (Expressed as Percentages of the At-Risk Population) Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 5d *First Use of Marijuana among Adults Aged 26 or Older, by State: Average Annual Incidence Estimates (Expressed as Percentages of the At-Risk Population) Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 6a *Illicit Drug Use Other Than Marijuana in the Past Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 6b *Illicit Drug Use Other Than Marijuana in the Past Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 6c *Illicit Drug Use Other Than Marijuana in the Past Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 6d *Illicit Drug Use Other Than Marijuana in the Past Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 7a *Cocaine Use in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 7b *Cocaine Use in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 7c *Cocaine Use in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 7d *Cocaine Use in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 8a *Perceptions of Great Risk from Using Cocaine Once a Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 8b *Perceptions of Great Risk from Using Cocaine Once a Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 8c *Perceptions of Great Risk from Using Cocaine Once a Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 8d *Perceptions of Great Risk from Using Cocaine Once a Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 9a *Heroin Use in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 9b *Heroin Use in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 10a *Perceptions of Great Risk from Trying Heroin Once or Twice among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 10b *Perceptions of Great Risk from Trying Heroin Once or Twice among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 10c *Perceptions of Great Risk from Trying Heroin Once or Twice among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 10d *Perceptions of Great Risk from Trying Heroin Once or Twice among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 11a *Pain Reliever Misuse in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 11b *Pain Reliever Misuse in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 11c *Pain Reliever Misuse in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 11d *Pain Reliever Misuse in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 12a *Alcohol Use in the Past Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 12b *Alcohol Use in the Past Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 12c *Alcohol Use in the Past Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 12d *Alcohol Use in the Past Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 13a *Binge Alcohol Use in the Past Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 13b *Binge Alcohol Use in the Past Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 13c *Binge Alcohol Use in the Past Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 13d *Binge Alcohol Use in the Past Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 14a *Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 14b *Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 14c *Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 14d *Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 15a *Alcohol Use in the Past Month among Individuals Aged 12 to 20, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 15b *Binge Alcohol Use in the Past Month among Individuals Aged 12 to 20, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 16a *Tobacco Product Use in the Past Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 16b *Tobacco Product Use in the Past Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 16c *Tobacco Product Use in the Past Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 16d *Tobacco Product Use in the Past Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 17a *Cigarette Use in the Past Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 17b *Cigarette Use in the Past Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 17c *Cigarette Use in the Past Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 17d *Cigarette Use in the Past Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 18a *Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 18b *Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 18c *Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 18d *Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 19a *Illicit Drug Use Disorder in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 19b *Illicit Drug Use Disorder in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 19c *Illicit Drug Use Disorder in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 19d *Illicit Drug Use Disorder in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 20c *Pain Reliever Use Disorder in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 20d *Pain Reliever Use Disorder in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 21a *Alcohol Use Disorder in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 21b *Alcohol Use Disorder in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 21c *Alcohol Use Disorder in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 21d *Alcohol Use Disorder in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 22a *Substance Use Disorder in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 22b *Substance Use Disorder in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 22c *Substance Use Disorder in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 22d *Substance Use Disorder in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 23a *Needing But Not Receiving Treatment at a Specialty Facility for Illicit Drug Use in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 23b *Needing But Not Receiving Treatment at a Specialty Facility for Illicit Drug Use in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 23c *Needing But Not Receiving Treatment at a Specialty Facility for Illicit Drug Use in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 23d *Needing But Not Receiving Treatment at a Specialty Facility for Illicit Drug Use in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 24a *Needing But Not Receiving Treatment at a Specialty Facility for Alcohol Use in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 24b *Needing But Not Receiving Treatment at a Specialty Facility for Alcohol Use in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 24c *Needing But Not Receiving Treatment at a Specialty Facility for Alcohol Use in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 24d *Needing But Not Receiving Treatment at a Specialty Facility for Alcohol Use in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 25a *Needing But Not Receiving Treatment at a Specialty Facility for Substance Use in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 25b *Needing But Not Receiving Treatment at a Specialty Facility for Substance Use in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 25c *Needing But Not Receiving Treatment at a Specialty Facility for Substance Use in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 25d *Needing But Not Receiving Treatment at a Specialty Facility for Substance Use in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 26a *Serious Mental Illness in the Past Year among Adults Aged 18 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 26b *Serious Mental Illness in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 26c *Serious Mental Illness in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 27a *Any Mental Illness in the Past Year among Adults Aged 18 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 27b *Any Mental Illness in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 27c *Any Mental Illness in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 28a *Received Mental Health Services in the Past Year among Adults Aged 18 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 28b *Received Mental Health Services in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 28c *Received Mental Health Services in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 29a *Had Serious Thoughts of Suicide in the Past Year among Adults Aged 18 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 29b *Had Serious Thoughts of Suicide in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 29c *Had Serious Thoughts of Suicide in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 30a *Major Depressive Episode in the Past Year among Adults Aged 18 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 30b *Major Depressive Episode in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 30c *Major Depressive Episode in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*



Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.

Figure 30d *Major Depressive Episode in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2015 and 2016 NSDUHs*



Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2015 and 2016.