

ARIZONA

Table 15 Selected Drug Use, Perceptions of Great Risk, Past Year Substance Use Disorder and Treatment, and Past Year Mental Health Measures in Arizona, by Age Group: Estimated Numbers (in Thousands), Annual Averages Based on 2015-2016 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLICIT DRUGS					
Past Month Illicit Drug Use ^{1,2}	568	48	157	363	520
Past Year Marijuana Use	696	65	214	417	631
Past Month Marijuana Use	422	37	133	252	385
Perceptions of Great Risk from Smoking Marijuana Once a Month	1,638	149	109	1,380	1,489
First Use of Marijuana ^{3,4}	64	25	27	12	39
Past Month Illicit Drug Use Other Than Marijuana ^{1,2}	223	15	57	151	208
Past Year Cocaine Use	118	4	46	67	113
Perceptions of Great Risk from Using Cocaine Once a Month	3,977	305	474	3,198	3,673
Past Year Heroin Use	26	0	7	19	26
Perceptions of Great Risk from Trying Heroin Once or Twice	4,818	358	616	3,844	4,460
Past Year Misuse of Pain Relievers ^{2,5}	267	23	59	185	244
ALCOHOL					
Past Month Alcohol Use	2,901	44	403	2,454	2,857
Past Month Binge Alcohol Use ⁶	1,285	24	256	1,005	1,260
Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week	2,724	255	289	2,180	2,469
Past Month Alcohol Use ⁷ (Individuals Aged 12 to 20)	132	--	--	--	--
Past Month Binge Alcohol Use ^{6,7} (Individuals Aged 12 to 20)	82	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁸	1,230	24	225	981	1,206
Past Month Cigarette Use	1,024	16	178	830	1,008
Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day	4,202	379	512	3,310	3,822
PAST YEAR SUBSTANCE USE DISORDER AND TREATMENT					
Illicit Drug Use Disorder ^{1,2,9}	198	26	65	108	172
Pain Reliever Use Disorder ^{2,5,9}	43	4	8	31	39
Alcohol Use Disorder ⁹	304	13	85	206	292
Substance Use Disorder ^{1,2,9}	431	31	124	276	400
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,2,10}	165	26	59	80	139
Needing But Not Receiving Treatment for Alcohol Use ¹⁰	290	12	86	191	278
Needing But Not Receiving Treatment for Substance Use ^{1,2,10}	416	29	116	271	387
PAST YEAR MENTAL HEALTH ISSUES					
Serious Mental Illness ^{4,11}	--	--	41	167	208
Any Mental Illness ^{4,11}	--	--	154	748	902
Received Mental Health Services ¹²	--	--	78	533	611
Had Serious Thoughts of Suicide ¹³	--	--	67	140	206
Major Depressive Episode ^{4,14}	--	64	64	246	310

-- Not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach.

NOTE: Estimated numbers appearing as 0 in this table mean that the estimate is greater than 0 but less than 500 because estimated numbers are shown in thousands.

¹ Illicit Drug Use includes the misuse of prescription psychotherapeutics or the use of marijuana, cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine. Illicit Drug Use Other Than Marijuana includes the misuse of prescription psychotherapeutics or the use of cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine.² Misuse of prescription psychotherapeutics is defined as use in any way not directed by a doctor, including use without a prescription of one's own; use in greater amounts, more often, or longer than told; or use in any other way not directed by a doctor. Prescription psychotherapeutics do not include over-the-counter drugs.³ First Use of Marijuana is defined as follows: *Average annual number of marijuana initiates* = $X_1 \div 2$, where X_1 is the number of marijuana initiates in the past 24 months.⁴ For details, see Section B of the "2015-2016 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at <https://www.samhsa.gov/data/>.⁵ Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.⁶ Binge Alcohol Use is defined as drinking five or more drinks (for males) or four or more drinks (for females) on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. In 2015, the definition for females changed from five to four drinks.⁷ Underage drinking is defined for individuals aged 12 to 20; therefore, the "12+" estimate reflects that age group and not individuals aged 12 or older.⁸ Tobacco Products include cigarettes, smokeless tobacco (i.e., snuff, dip, chewing tobacco, or "snus"), cigars, or pipe tobacco.⁹ Substance Use Disorder is defined as meeting criteria for illicit drug or alcohol dependence or abuse. Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).¹⁰ Respondents were classified as needing treatment for a substance use problem if they met the criteria for a substance use disorder as defined in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) or received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).¹¹ Mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, assessed by the Mental Health Surveillance Study (MHSS) *Structured Clinical Interview for the Diagnostic and Statistical Manual of Mental Disorders—Fourth Edition—Research Version—Axis I Disorders* (MHSS-SCID), which is based on the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Any mental illness (AMI) includes adults with any mental disorder regardless of whether their disorder resulted in functional impairment. Serious mental illness (SMI) includes adults with any mental disorder that resulted in serious functional impairment.¹² Mental health services are defined as having received inpatient treatment/counseling or outpatient treatment/counseling or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use.¹³ Respondents were asked, "At any time in the past 12 months, did you seriously think about trying to kill yourself?" If they answered "Yes," they were categorized as having serious thoughts of suicide in the past year.¹⁴ Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when an individual experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from adults aged 18 or older to produce an estimate for those aged 12 or older.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015 and 2016.

ARIZONA

Table 16 Selected Drug Use, Perceptions of Great Risk, Past Year Substance Use Disorder and Treatment, and Past Year Mental Health Measures in Arizona, by Age Group: Percentages, Annual Averages Based on 2015-2016 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLCIT DRUGS					
Past Month Illicit Drug Use ^{1,2}	9.97	8.79	21.05	8.24	10.10
Past Year Marijuana Use	12.22	11.86	28.68	9.48	12.26
Past Month Marijuana Use	7.42	6.75	17.81	5.74	7.49
Perceptions of Great Risk from Smoking Marijuana Once a Month	28.75	27.10	14.64	31.37	28.92
First Use of Marijuana ^{3,4}	2.01	5.12	7.31	0.50	1.46
Past Month Illicit Drug Use Other Than Marijuana ^{1,2}	3.92	2.72	7.62	3.44	4.05
Past Year Cocaine Use	2.07	0.79	6.22	1.52	2.20
Perceptions of Great Risk from Using Cocaine Once a Month	69.86	55.56	63.55	72.70	71.37
Past Year Heroin Use	0.45	0.03	0.88	0.43	0.50
Perceptions of Great Risk from Trying Heroin Once or Twice	84.64	65.29	82.60	87.36	86.67
Past Year Misuse of Pain Relievers ^{2,5}	4.69	4.21	7.92	4.20	4.74
ALCOHOL					
Past Month Alcohol Use	50.94	7.95	54.01	55.78	55.53
Past Month Binge Alcohol Use ⁶	22.56	4.43	34.25	22.84	24.49
Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week	47.83	46.50	38.70	49.56	47.98
Past Month Alcohol Use ⁷ (Individuals Aged 12 to 20)	16.74	--	--	--	--
Past Month Binge Alcohol Use ^{6,7} (Individuals Aged 12 to 20)	10.44	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁸	21.60	4.35	30.19	22.29	23.44
Past Month Cigarette Use	17.98	2.94	23.81	18.87	19.59
Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day	73.79	69.14	68.63	75.24	74.28
PAST YEAR SUBSTANCE USE DISORDER AND TREATMENT					
Illicit Drug Use Disorder ^{1,2,9}	3.48	4.69	8.68	2.44	3.35
Pain Reliever Use Disorder ^{2,5,9}	0.75	0.65	1.14	0.70	0.76
Alcohol Use Disorder ⁹	5.34	2.28	11.46	4.68	5.66
Substance Use Disorder ^{1,2,9}	7.57	5.62	16.66	6.28	7.78
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,2,10}	2.89	4.75	7.91	1.81	2.69
Needing But Not Receiving Treatment for Alcohol Use ¹⁰	5.09	2.24	11.54	4.35	5.39
Needing But Not Receiving Treatment for Substance Use ^{1,2,10}	7.30	5.23	15.59	6.15	7.52
PAST YEAR MENTAL HEALTH ISSUES					
Serious Mental Illness ^{4,11}	--	--	5.52	3.79	4.04
Any Mental Illness ^{4,11}	--	--	20.65	16.99	17.52
Received Mental Health Services ¹²	--	--	10.44	12.12	11.87
Had Serious Thoughts of Suicide ¹³	--	--	8.92	3.18	4.01
Major Depressive Episode ^{4,14}	--	11.68	8.61	5.60	6.03

-- Not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach.

¹ Illicit Drug Use includes the misuse of prescription psychotherapeutics or the use of marijuana, cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine. Illicit Drug Use Other Than Marijuana includes the misuse of prescription psychotherapeutics or the use of cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine.

² Misuse of prescription psychotherapeutics is defined as use in any way not directed by a doctor, including use without a prescription of one's own; use in greater amounts, more often, or longer than told; or use in any other way not directed by a doctor. Prescription psychotherapeutics do not include over-the-counter drugs.

³ First Use of Marijuana is defined as follows: $Average\ annual\ initiation\ of\ marijuana\ (\%) = 100 * \{ [X_1 + (0.5 * X_1 + X_2)] \div 2 \}$, where X_1 is the number of marijuana initiates in the past 24 months and X_2 is the number of individuals who never used marijuana (with the at-risk population defined as $0.5 * X_1 + X_2$). Both of the computation components, X_1 and X_2 , are based on a survey-weighted hierarchical Bayes estimation approach. The age group shown is based on a respondent's age at the time of the interview, not his or her age at first use.

⁴ For details, see Section B of the "2015-2016 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at <https://www.samhsa.gov/data/>.

⁵ Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

⁶ Binge Alcohol Use is defined as drinking five or more drinks (for males) or four or more drinks (for females) on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. In 2015, the definition for females changed from five to four drinks.

⁷ Underage drinking is defined for individuals aged 12 to 20; therefore, the "12+" estimate reflects that age group and not individuals aged 12 or older.

⁸ Tobacco Products include cigarettes, smokeless tobacco (i.e., snuff, dip, chewing tobacco, or "snus"), cigars, or pipe tobacco.

⁹ Substance Use Disorder is defined as meeting criteria for illicit drug or alcohol dependence or abuse. Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹⁰ Respondents were classified as needing treatment for a substance use problem if they met the criteria for a substance use disorder as defined in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) or received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹¹ Mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, assessed by the Mental Health Surveillance Study (MHSS) *Structured Clinical Interview for the Diagnostic and Statistical Manual of Mental Disorders—Fourth Edition—Research Version—Axis I Disorders* (MHSS-SCID), which is based on the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Any mental illness (AMI) includes adults with any mental disorder regardless of whether their disorder resulted in functional impairment. Serious mental illness (SMI) includes adults with any mental disorder that resulted in serious functional impairment.

¹² Mental health services are defined as having received inpatient treatment/counseling or outpatient treatment/counseling or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use.

¹³ Respondents were asked, "At any time in the past 12 months, did you seriously think about trying to kill yourself?" If they answered "Yes," they were categorized as having serious thoughts of suicide in the past year.

¹⁴ Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when an individual experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from adults aged 18 or older to produce an estimate for those aged 12 or older.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015 and 2016.