		licit Drug Use he Past Month		Illicit Drug Use Other Than Marijuana in the Past Month		
State/Substate Region	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval		
Arkansas	8.37	(7.22 - 9.68)	3.82	(3.07 - 4.74)		
Catchment Area 1	8.20	(6.41 - 10.43)	3.72	(2.65 - 5.21)		
Catchment Area 2	7.45	(5.59 - 9.86)	3.62	(2.51 - 5.18)		
Catchment Area 3	8.65	(6.66 - 11.16)	3.91	(2.77 - 5.48)		
Catchment Area 4	8.06	(5.96 - 10.81)	3.66	(2.53 - 5.27)		
Catchment Area 5	9.66	(7.52 - 12.33)	4.29	(3.06 - 5.98)		
Catchment Area 6	8.80	(6.59 - 11.65)	3.65	(2.48 - 5.33)		
Catchment Area 7	7.84	(5.79 - 10.55)	3.77	(2.58 - 5.49)		
Catchment Area 8	8.12	(6.18 - 10.60)	3.76	(2.63 - 5.35)		

Table 4.1 Illicit Drug Use in the Past Month and Illicit Drug Use Other Than Marijuanain the Past Month in Arkansas among Individuals Aged 12 or Older, by SubstateRegion: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs

NOTE: For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a surveyweighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates include data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

Table 4.2Marijuana Use in the Past Month, Average Annual Incidence Estimates
(Expressed as Percentages of the At-Risk Population) of First Use of Marijuana,
and Perceptions of Great Risk from Smoking Marijuana Once a Month
in Arkansas among Individuals Aged 12 or Older, by Substate Region:
Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs

	Marijuana Use in the Past Month		Incider (Ex Percent Risk Pop	rage Annual nce Estimates pressed as ages of the At- ulation) of First f Marijuana ¹	Perceptions of Great Risk from Smoking Marijuana Once a Month	
State/Substate Region	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval
Arkansas	6.24	(5.25 - 7.41)	1.73	(1.52 - 1.97)	34.99	(32.53 - 37.54)
Catchment Area 1	6.46	(4.87 - 8.53)	2.01	(1.61 - 2.51)	33.02	(28.65 - 37.71)
Catchment Area 2	5.48	(3.91 - 7.61)	1.50	(1.15 - 1.97)	33.08	(28.28 - 38.26)
Catchment Area 3	5.76	(4.21 - 7.83)	1.22	(0.94 - 1.57)	39.74	(35.22 - 44.45)
Catchment Area 4	6.10	(4.36 - 8.47)	1.44	(1.11 - 1.87)	31.39	(26.72 - 36.48)
Catchment Area 5	7.42	(5.56 - 9.84)	1.92	(1.52 - 2.44)	32.94	(28.46 - 37.75)
Catchment Area 6	6.96	(5.00 - 9.61)	1.90	(1.43 - 2.52)	41.33	(36.20 - 46.65)
Catchment Area 7	5.49	(3.91 - 7.66)	1.66	(1.25 - 2.19)	39.01	(33.71 - 44.59)
Catchment Area 8	6.07	(4.51 - 8.13)	2.00	(1.56 - 2.56)	33.85	(29.40 - 38.60)

NOTE: For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a surveyweighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹*Average annual initiation of marijuana* (%) = 100 * { $[X_1 \div (0.5 * X_1 + X_2)] \div 2$ }, where X_1 is the number of marijuana initiates in the past 24 months and X_2 is the number of individuals who never used marijuana (with the at-risk population defined as $0.5 * X_1 + X_2$). Both of the computation components, X_1 and X_2 , are based on a survey-weighted hierarchical Bayes estimation approach. The age group shown is based on a respondent's age at the time of the interview, not his or her age at first use. The Bayesian confidence (credible) intervals presented for Average Annual Estimates of First Use of Marijuana use a simultaneous solution for First Use of Marijuana and Never Used Marijuana outcomes, which had age group-specific random effects that were allowed to be correlated across the two outcomes. The associated Markov Chain Monte Carlo chains were used to calculate the posterior variance. For details, see Section B of the "2011-2012 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at http://www.samhsa.gov/data/.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

Table 4.3 Marijuana Use in the Past Year, Cocaine Use in the Past Year, and Nonmedical
Use of Pain Relievers in the Past Year in Arkansas among Individuals Aged 12 or
Older, by Substate Region: Percentages, Annual Averages Based on 2012, 2013,
and 2014 NSDUHs

		Marijuana Use in the Past Year		Cocaine Use in the Past Year		Nonmedical Use of Pain Relievers in the Past Year	
State/Substate Region	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval	
Arkansas	10.41	(9.12 - 11.86)	1.26	(0.90 - 1.77)	5.21	(4.37 - 6.20)	
Catchment Area 1	11.25	(9.05 - 13.91)	1.26	(0.78 - 2.04)	5.06	(3.94 - 6.47)	
Catchment Area 2	8.62	(6.65 - 11.12)	1.04	(0.60 - 1.79)	4.83	(3.71 - 6.28)	
Catchment Area 3	9.58	(7.40 - 12.32)	1.09	(0.63 - 1.88)	5.64	(4.36 - 7.28)	
Catchment Area 4	9.38	(7.08 - 12.34)	1.19	(0.70 - 2.02)	5.25	(4.00 - 6.85)	
Catchment Area 5	12.17	(9.76 - 15.09)	1.58	(0.96 - 2.58)	5.56	(4.34 - 7.11)	
Catchment Area 6	11.24	(8.71 - 14.40)	1.50	(0.84 - 2.65)	5.18	(3.93 - 6.79)	
Catchment Area 7	9.14	(6.84 - 12.11)	1.05	(0.59 - 1.86)	5.05	(3.80 - 6.67)	
Catchment Area 8	10.71	(8.50 - 13.41)	1.32	(0.76 - 2.27)	5.08	(3.93 - 6.54)	

NOTE: For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a surveyweighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

Table 4.4Alcohol Use in the Past Month, Binge Alcohol Use in the Past Month, and
Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic
Beverage Once or Twice a Week in Arkansas among Individuals Aged 12 or
Older, by Substate Region: Percentages, Annual Averages Based on 2012, 2013,
and 2014 NSDUHs

	Alcohol Use in the Past Month		Binge Alcohol Use in the Past Month ¹		Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week	
State/Substate Region	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval
Arkansas	41.43	(38.98 - 43.92)	20.64	(18.68 - 22.75)	40.80	(38.28 - 43.38)
Catchment Area 1	44.62	(39.61 - 49.75)	21.30	(18.17 - 24.81)	40.09	(35.92 - 44.41)
Catchment Area 2	39.60	(34.50 - 44.94)	21.01	(17.59 - 24.88)	38.95	(34.68 - 43.40)
Catchment Area 3	34.94	(30.07 - 40.14)	19.08	(15.93 - 22.67)	42.29	(37.95 - 46.75)
Catchment Area 4	37.39	(32.12 - 42.98)	20.52	(17.00 - 24.56)	39.80	(35.45 - 44.31)
Catchment Area 5	46.11	(40.94 - 51.36)	21.12	(17.87 - 24.78)	39.14	(34.80 - 43.67)
Catchment Area 6	39.60	(33.88 - 45.62)	22.55	(18.80 - 26.81)	44.64	(40.13 - 49.23)
Catchment Area 7	36.38	(30.56 - 42.62)	18.60	(15.21 - 22.54)	42.92	(38.50 - 47.46)
Catchment Area 8	45.28	(39.98 - 50.68)	20.75	(17.43 - 24.50)	41.01	(36.65 - 45.52)

NOTE: For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a surveyweighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

		Jse in the Past Month ividuals Aged 12 to 20	Binge Alcohol Use in the Past Month among Individuals Aged 12 to 20 ¹		
State/Substate Region	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval	
Arkansas	19.19	(17.30 - 21.22)	13.26	(11.70 - 15.00)	
Catchment Area 1	20.17	(17.05 - 23.70)	14.04	(11.37 - 17.23)	
Catchment Area 2	20.13	(16.57 - 24.24)	15.30	(12.09 - 19.17)	
Catchment Area 3	16.89	(13.81 - 20.50)	12.00	(9.39 - 15.20)	
Catchment Area 4	18.12	(14.76 - 22.05)	13.72	(10.87 - 17.18)	
Catchment Area 5	22.33	(18.57 - 26.59)	14.69	(11.57 - 18.48)	
Catchment Area 6	19.39	(15.52 - 23.96)	13.84	(10.74 - 17.65)	
Catchment Area 7	17.31	(13.83 - 21.46)	11.22	(8.61 - 14.50)	
Catchment Area 8	18.00	(14.97 - 21.50)	11.49	(9.12 - 14.39)	

Table 4.5Alcohol Use in the Past Month and Binge Alcohol Use in the Past Month among
Individuals Aged 12 to 20 in Arkansas, by Substate Region: Percentages, Annual
Averages Based on 2012, 2013, and 2014 NSDUHs

NOTE: For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a surveyweighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

Table 4.6 Cigarette Use in the Past Month, Tobacco Product Use in the Past Month, and
Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day
in Arkansas among Individuals Aged 12 or Older, by Substate Region:
Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs

	U	arette Use Past Month) Product Use Past Month ¹	from Sm More Pacl	s of Great Risk oking One or xs of Cigarettes er Day
State/Substate Region	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval
Arkansas	27.05	(25.05 - 29.15)	33.86	(31.59 - 36.20)	65.34	(63.16 - 67.46)
Catchment Area 1	20.64	(17.42 - 24.27)	27.33	(23.55 - 31.47)	68.73	(64.83 - 72.38)
Catchment Area 2	31.37	(27.20 - 35.88)	39.41	(34.64 - 44.39)	61.31	(56.84 - 65.60)
Catchment Area 3	29.46	(25.61 - 33.64)	35.99	(31.59 - 40.64)	62.97	(58.76 - 66.99)
Catchment Area 4	27.34	(23.30 - 31.80)	34.62	(30.08 - 39.46)	63.15	(58.69 - 67.39)
Catchment Area 5	27.82	(24.08 - 31.90)	35.00	(30.63 - 39.64)	66.09	(61.93 - 70.00)
Catchment Area 6	31.02	(26.55 - 35.87)	37.73	(32.71 - 43.02)	65.41	(60.98 - 69.60)
Catchment Area 7	28.55	(24.21 - 33.32)	34.60	(29.77 - 39.76)	64.93	(60.33 - 69.27)
Catchment Area 8	25.27	(21.58 - 29.35)	31.43	(27.36 - 35.80)	67.22	(63.00 - 71.19)

NOTE: For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a surveyweighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

		hol Dependence the Past Year		Drug Dependence he Past Year ¹
State/Substate Region	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval
Arkansas	2.85	(2.30 - 3.52)	1.82	(1.47 - 2.25)
Catchment Area 1	2.85	(2.03 - 3.98)	1.62	(1.17 - 2.23)
Catchment Area 2	2.88	(2.00 - 4.12)	1.49	(1.05 - 2.11)
Catchment Area 3	2.73	(1.91 - 3.89)	1.89	(1.35 - 2.64)
Catchment Area 4	2.84	(1.98 - 4.05)	1.54	(1.11 - 2.14)
Catchment Area 5	2.86	(2.03 - 4.02)	1.97	(1.41 - 2.73)
Catchment Area 6	2.96	(2.03 - 4.29)	2.25	(1.59 - 3.18)
Catchment Area 7	2.64	(1.82 - 3.81)	1.89	(1.33 - 2.67)
Catchment Area 8	2.95	(2.08 - 4.15)	1.97	(1.41 - 2.76)

Table 4.7Alcohol Dependence in the Past Year and Illicit Drug Dependence in the Past Yearin Arkansas among Individuals Aged 12 or Older, by Substate Region:
Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs

NOTE: For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a surveyweighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

NOTE: Dependence is based on the definition found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescriptiontype psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

Table 4.8Alcohol Dependence or Abuse in the Past Year, Illicit Drug Dependence or Abuse
in the Past Year, and Dependence or Abuse of Illicit Drugs or Alcohol in the
Past Year in Arkansas among Individuals Aged 12 or Older, by Substate Region:
Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs

		Alcohol Dependence or Abuse in the Past Year		Illicit Drug Dependence or Abuse in the Past Year ¹		Dependence or Abuse of Illicit Drugs or Alcohol in the Past Year ¹	
State/Substate Region	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval	
Arkansas	5.50	(4.71 - 6.41)	2.67	(2.19 - 3.26)	7.53	(6.66 - 8.51)	
Catchment Area 1	6.08	(4.85 - 7.58)	2.60	(1.96 - 3.44)	7.79	(6.44 - 9.39)	
Catchment Area 2	5.26	(4.05 - 6.80)	2.17	(1.57 - 3.00)	6.94	(5.56 - 8.62)	
Catchment Area 3	5.20	(4.11 - 6.57)	2.71	(1.99 - 3.69)	7.67	(6.17 - 9.49)	
Catchment Area 4	5.97	(4.61 - 7.70)	2.36	(1.72 - 3.23)	7.56	(6.08 - 9.36)	
Catchment Area 5	5.68	(4.47 - 7.18)	2.81	(2.09 - 3.77)	7.82	(6.40 - 9.52)	
Catchment Area 6	5.26	(4.05 - 6.80)	3.23	(2.30 - 4.51)	7.57	(6.09 - 9.36)	
Catchment Area 7	4.93	(3.76 - 6.45)	2.85	(2.06 - 3.93)	7.33	(5.89 - 9.09)	
Catchment Area 8	5.33	(4.19 - 6.75)	2.79	(2.07 - 3.76)	7.43	(6.02 - 9.12)	

NOTE: For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a surveyweighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescriptiontype psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

Table 4.9Needing But Not Receiving Treatment for Alcohol Use in the Past Year and
Needing But Not Receiving Treatment for Illicit Drug Use in the Past Year
in Arkansas among Individuals Aged 12 or Older, by Substate Region:
Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs

	C	g But Not Receiving r Alcohol Use in the Pa Year ¹		Needing But Not Receiving Treatment for Illicit Drug Use in the Past Year ²		
State/Substate Region	Estimate	95% Prediction Interval	Estimate	95% Prediction Interval		
Arkansas	5.26	(4.46 - 6.19)	2.34	(1.93 - 2.85)		
Catchment Area 1	5.57	(4.43 - 6.99)	2.20	(1.66 - 2.90)		
Catchment Area 2	5.16	(3.95 - 6.70)	1.92	(1.41 - 2.61)		
Catchment Area 3	5.14	(3.98 - 6.61)	2.52	(1.86 - 3.41)		
Catchment Area 4	5.40	(4.19 - 6.92)	2.05	(1.51 - 2.79)		
Catchment Area 5	5.38	(4.20 - 6.88)	2.43	(1.83 - 3.23)		
Catchment Area 6	5.09	(3.90 - 6.63)	2.68	(1.95 - 3.67)		
Catchment Area 7	4.84	(3.70 - 6.30)	2.56	(1.90 - 3.46)		
Catchment Area 8	5.23	(4.03 - 6.76)	2.47	(1.84 - 3.29)		

NOTE: For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a surveyweighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but not receiving treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], or mental health centers).

²Needing But Not Receiving Treatment refers to respondents classified as needing treatment for illicit drugs, but not receiving treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], or mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

		us Mental Illness the Past Year	•	y Mental Illness the Past Year
State/Substate Region	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Arkansas	5.20	(4.28 - 6.30)	20.13	(18.13 - 22.29)
Catchment Area 1	4.98	(3.73 - 6.62)	18.44	(15.67 - 21.57)
Catchment Area 2	6.16	(4.52 - 8.34)	21.03	(17.91 - 24.54)
Catchment Area 3	5.12	(3.82 - 6.83)	21.06	(17.90 - 24.60)
Catchment Area 4	5.24	(3.86 - 7.08)	20.21	(17.07 - 23.75)
Catchment Area 5	5.16	(3.81 - 6.95)	20.19	(17.21 - 23.54)
Catchment Area 6	5.06	(3.65 - 6.98)	19.57	(16.56 - 22.98)
Catchment Area 7	5.44	(3.92 - 7.51)	20.71	(17.39 - 24.48)
Catchment Area 8	4.78	(3.54 - 6.42)	20.21	(17.18 - 23.62)

Table 4.10Serious Mental Illness in the Past Year and Any Mental Illness in the Past Yearin Arkansas among Adults Aged 18 or Older, by Substate Region: Percentages,
Annual Averages Based on 2012, 2013, and 2014 NSDUHs

NOTE: For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a surveyweighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, assessed by the Mental Health Surveillance Study (MHSS) *Structured Clinical Interview for the Diagnostic and Statistical Manual of Mental Disorders—Fourth Edition—Research Version—Axis I Disorders* (MHSS-SCID), which is based on the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes individuals in any of the three categories. For details, see Section B of the "2011-2012 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at http://www.samhsa.gov/data/.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

		us Thoughts of Suicide the Past Year ¹	Major Depressive Episode in the Past Year ²		
State/Substate Region	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	
Arkansas	4.45	(3.75 - 5.27)	7.17	(6.14 - 8.36)	
Catchment Area 1	4.23	(3.24 - 5.51)	7.10	(5.61 - 8.95)	
Catchment Area 2	4.97	(3.67 - 6.70)	7.63	(5.96 - 9.73)	
Catchment Area 3	4.44	(3.33 - 5.91)	7.22	(5.71 - 9.10)	
Catchment Area 4	4.63	(3.46 - 6.16)	6.89	(5.37 - 8.81)	
Catchment Area 5	4.46	(3.37 - 5.89)	7.00	(5.50 - 8.88)	
Catchment Area 6	4.21	(3.13 - 5.65)	7.04	(5.42 - 9.09)	
Catchment Area 7	4.61	(3.40 - 6.23)	7.05	(5.47 - 9.04)	
Catchment Area 8	4.24	(3.20 - 5.60)	7.26	(5.74 - 9.13)	

Table 4.11Had Serious Thoughts of Suicide in the Past Year and Major Depressive Episodein the Past Year in Arkansas among Adults Aged 18 or Older, by SubstateRegion: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs

NOTE: For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.


NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a surveyweighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹Respondents were asked, "At any time in the past 12 months, did you seriously think about trying to kill yourself?" If they answered "Yes," they were categorized as having serious thoughts of suicide in the past year.

²Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when an individual experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. For details, see Section B in the "2011-2012 National Survey on Drug Use and Health: Guide to State Tables and Summary of Small Area Estimation Methodology" at http://www.samhsa.gov/data/.


Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

Figure 4.1 *Illicit Drug Use in the Past Month* among Individuals Aged 12 or Older in Arkansas, by Substate Region: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs




- NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.
- Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

Figure 4.2 *Marijuana Use in the Past Year* among Individuals Aged 12 or Older in Arkansas, by Substate Region: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs




- NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.
- Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

Figure 4.3 *Marijuana Use in the Past Month* among Individuals Aged 12 or Older in Arkansas, by Substate Region: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs




NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.

Figure 4.4 Perceptions of Great Risk from Smoking Marijuana Once a Month among Individuals Aged 12 or Older in Arkansas, by Substate Region: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs




NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.

Figure 4.5 *First Use of Marijuana* among Individuals Aged 12 or Older in Arkansas, by Substate Region: Average Annual Incidence Estimates (Expressed as Percentages of the At-Risk Population) Based on 2012, 2013, and 2014 NSDUHs





- NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.
- Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

Figure 4.6 Illicit Drug Use Other Than Marijuana in the Past Month among Individuals Aged 12 or Older in Arkansas, by Substate Region: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs



NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.





NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.




Figure 4.8 Nonmedical Use of Pain Relievers in the Past Year among Individuals Aged 12 or Older in Arkansas, by Substate Region: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs

- NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.
- Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.






Figure 4.9 Alcohol Use in the Past Month among Individuals Aged 12 or Older in Arkansas, by Substate Region: Percentages, Annual Averages Based on 2012, 2013, and 2014

NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.


Substate Regions





- NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.
- Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.


Figure 4.11 Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Individuals Aged 12 or Older in Arkansas, by Substate Region: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs



NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.




Figure 4.12 Alcohol Use in the Past Month among Individuals Aged 12 to 20 in Arkansas,





- NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.
- Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

Figure 4.13 Binge Alcohol Use in the Past Month among Individuals Aged 12 to 20 in Arkansas, by Substate Region: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs





NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.






- NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.
- Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.





- NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.
- Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

Figure 4.16 Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day among Individuals Aged 12 or Older in Arkansas, by Substate Region: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs



NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.






Figure 4.17 Alcohol Dependence or Abuse in the Past Year among Individuals Aged 12 or Older in Arkansas, by Substate Region: Percentages, Annual Averages Based on


- NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.
- Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.







- NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.
- Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

Figure 4.19 *Illicit Drug Dependence or Abuse in the Past Year* among Individuals Aged 12 or Older in Arkansas, by Substate Region: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs




NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.






- NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.
- Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

Figure 4.21 Dependence or Abuse of Illicit Drugs or Alcohol in the Past Year among Individuals Aged 12 or Older in Arkansas, by Substate Region: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs




- NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.
- Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

Figure 4.22 Needing But Not Receiving Treatment for Illicit Drug Use in the Past Year among Individuals Aged 12 or Older in Arkansas, by Substate Region: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs





NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.

Figure 4.23 Needing But Not Receiving Treatment for Alcohol Use in the Past Year among Individuals Aged 12 or Older in Arkansas, by Substate Region: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs





- NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.
- Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.






- NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.
- Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.






- NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.
- Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.

Figure 4.26 *Had Serious Thoughts of Suicide in the Past Year* among Adults Aged 18 or Older in Arkansas, by Substate Region: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs



NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.

Figure 4.27 *Major Depressive Episode in the Past Year* among Adults Aged 18 or Older in Arkansas, by Substate Region: Percentages, Annual Averages Based on 2012, 2013, and 2014 NSDUHs



- NOTE: The legend's ranges were created by dividing 362 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2012-2014 National Survey on Drug Use and Health Substate Region Definitions" at http://www.samhsa.gov/data/.
- Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012, 2013, and 2014.