

Table 1.74A (REVISED) Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Numbers in Thousands, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	8,615	8,454	1,452 ^a	1,113	611 ^b	363
GEOGRAPHIC DIVISION						
Northeast	1,176	1,330	184	255	67	98
New England	300	332	55	62	14	34
Middle Atlantic	876	998	128	193	53	64
Midwest	1,776	1,700	361	275	141	74
East North Central	1,211	1,086	281	198	117	60
West North Central	566	614	80	77	25	14
South	3,158	3,089	639 ^a	374	285 ^a	142
South Atlantic	1,795	1,703	394 ^a	201	179	73
East South Central	539	656	103	47	51 ^a	11
West South Central	824	730	141	126	55	58
West	2,506	2,335	268	209	117	49
Mountain	773	627	100	58	43	8
Pacific	1,733	1,708	169	151	74	41
COUNTY TYPE						
Large Metro	4,865	4,333	748	588	298	186
Small Metro	2,360	2,593	445	388	245	145
250K – 1 Mil. Pop.	1,568	1,779	319	256	165	114
< 250K Pop.	792	813	126	132	80	31
Nonmetro	1,390	1,528	259 ^a	137	67	31
Urbanized	697	593	109	56	25	7
Less Urbanized	570	748	133 ^a	58	34	19
Completely Rural	123	187	17	22	9	5

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 1.74B (REVISED) Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2007 and 2008

Geographic Characteristic	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
TOTAL	3.5	3.4	0.6 ^a	0.4	0.2 ^b	0.1
GEOGRAPHIC DIVISION						
Northeast	2.6	2.9	0.4	0.6	0.1	0.2
New England	2.5	2.8	0.5	0.5	0.1	0.3
Middle Atlantic	2.6	2.9	0.4	0.6	0.2	0.2
Midwest	3.2	3.1	0.7	0.5	0.3	0.1
East North Central	3.2	2.8	0.7	0.5	0.3	0.2
West North Central	3.4	3.7	0.5	0.5	0.1	0.1
South	3.5	3.4	0.7 ^a	0.4	0.3 ^a	0.2
South Atlantic	3.8	3.5	0.8 ^a	0.4	0.4	0.2
East South Central	3.7	4.4	0.7	0.3	0.3 ^a	0.1
West South Central	3.0	2.6	0.5	0.4	0.2	0.2
West	4.4	4.0	0.5	0.4	0.2	0.1
Mountain	4.5	3.6	0.6	0.3	0.2	0.0
Pacific	4.3	4.2	0.4	0.4	0.2	0.1
COUNTY TYPE						
Large Metro	3.7	3.3	0.6	0.4	0.2	0.1
Small Metro	3.2	3.4	0.6	0.5	0.3	0.2
250K – 1 Mil. Pop.	3.2	3.6	0.6	0.5	0.3	0.2
< 250K Pop.	3.2	3.2	0.5	0.5	0.3	0.1
Nonmetro	3.4	3.7	0.6 ^b	0.3	0.2	0.1
Urbanized	4.1	3.5	0.7	0.3	0.1	0.0
Less Urbanized	2.9	3.7	0.7 ^a	0.3	0.2	0.1
Completely Rural	2.7	4.6	0.4	0.6	0.2	0.1

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.