

**Table 2.57A (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	45,057	45,958	11,047	11,826	8,059	8,706
<b>GEOGRAPHIC DIVISION</b>						
Northeast	6,325	6,160	1,272	1,415	853	999
New England	1,666	1,594	355	274	221	172
Middle Atlantic	4,659	4,566	917	1,141	632	827
Midwest	11,431	11,384	3,066	2,856	2,190	2,125
East North Central	7,502	7,428	1,923	1,933	1,397	1,455
West North Central	3,930	3,956	1,143	923	792	671
South	17,300	18,176	4,479 <sup>a</sup>	5,099	3,422	3,985
South Atlantic	8,557	8,710	2,009	2,192	1,471	1,657
East South Central	3,478	3,842	974	1,244	775	1,008
West South Central	5,264	5,624	1,496	1,663	1,175	1,321
West	10,002	10,238	2,230	2,456	1,595	1,596
Mountain	3,806	4,110	957	940	710	590
Pacific	6,195	6,127	1,273	1,515	885	1,005
<b>COUNTY TYPE</b>						
Large Metro	19,293	19,038	3,865	4,114	2,714	2,818
Small Metro	14,483	15,634	3,733	4,089	2,579	3,022
250K – 1 Mil. Pop.	8,607 <sup>a</sup>	9,594	2,137	2,456	1,478	1,711
< 250K Pop.	5,876	6,041	1,596	1,633	1,101	1,311
Nonmetro	11,282	11,286	3,449	3,623	2,766	2,866
Urbanized	4,485	4,267	1,377	1,117	1,093 <sup>a</sup>	850
Less Urbanized	5,505	5,880	1,679 <sup>a</sup>	2,089	1,349 <sup>a</sup>	1,693
Completely Rural	1,293	1,139	393	417	323	322

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.57B (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	18.2	18.4	4.5	4.7	3.3	3.5
<b>GEOGRAPHIC DIVISION</b>						
Northeast	13.8	13.4	2.8	3.1	1.9	2.2
New England	13.9	13.2	3.0	2.3	1.8	1.4
Middle Atlantic	13.8	13.4	2.7	3.4	1.9	2.4
Midwest	20.9	20.7	5.6	5.2	4.0	3.9
East North Central	19.6	19.4	5.0	5.0	3.6	3.8
West North Central	23.8	23.9	6.9 <sup>a</sup>	5.6	4.8	4.0
South	19.2	20.0	5.0	5.6	3.8	4.4
South Atlantic	18.0	18.1	4.2	4.6	3.1	3.5
East South Central	23.6	25.9	6.6	8.4	5.3	6.8
West South Central	19.0	20.0	5.4	5.9	4.3	4.7
West	17.5	17.7	3.9	4.2	2.8	2.8
Mountain	22.0	23.4	5.5	5.3	4.1	3.4
Pacific	15.5	15.2	3.2	3.8	2.2	2.5
<b>COUNTY TYPE</b>						
Large Metro	14.5	14.3	2.9	3.1	2.0	2.1
Small Metro	19.5	20.7	5.0	5.4	3.5	4.0
250K – 1 Mil. Pop.	17.4 <sup>a</sup>	19.2	4.3	4.9	3.0	3.4
< 250K Pop.	23.9	23.7	6.5	6.4	4.5	5.1
Nonmetro	27.5	27.4	8.4	8.8	6.7	6.9
Urbanized	26.7	24.9	8.2 <sup>a</sup>	6.5	6.5 <sup>a</sup>	5.0
Less Urbanized	28.0	29.3	8.5 <sup>a</sup>	10.4	6.9 <sup>a</sup>	8.4
Completely Rural	28.1	28.3	8.6	10.4	7.0	8.0

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.58A (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	1,961	1,819	1,306	1,219	620	555
<b>GEOGRAPHIC DIVISION</b>						
Northeast	247	271	175	202	81	91
New England	64	61	43	48	17	18
Middle Atlantic	183	210	132	155	64	73
Midwest	484	436	323	285	149	147
East North Central	321	296	218	198	99	99
West North Central	162	140	105	88	50	49
South	897	797	589	529	303	243
South Atlantic	421	388	271	265	142	122
East South Central	222	190	152	130	84 <sup>a</sup>	54
West South Central	254	219	167	134	77	67
West	333	315	218	202	87	73
Mountain	133	143	84	99	37	37
Pacific	200	172	135	103	50	36
<b>COUNTY TYPE</b>						
Large Metro	714 <sup>a</sup>	596	492	418	196	171
Small Metro	596	625	384	427	166	201
250K – 1 Mil. Pop.	349	393	228	268	91	125
< 250K Pop.	247	232	157	159	76	76
Nonmetro	651	598	430	373	257 <sup>b</sup>	183
Urbanized	244	228	166	139	93	77
Less Urbanized	337	290	218	181	138 <sup>a</sup>	86
Completely Rural	70	79	45	53	26	20

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.58B (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	7.8	7.3	5.2	4.9	2.5	2.2
<b>GEOGRAPHIC DIVISION</b>						
Northeast	5.5	6.2	3.9	4.6	1.8	2.1
New England	5.6	5.4	3.7	4.2	1.5	1.6
Middle Atlantic	5.5	6.5	4.0	4.8	1.9	2.3
Midwest	8.6	7.9	5.8	5.2	2.6	2.7
East North Central	8.1	7.6	5.5	5.1	2.5	2.5
West North Central	9.7	8.5	6.3	5.3	3.0	3.0
South	9.8	8.8	6.5	5.8	3.3	2.7
South Atlantic	9.1	8.5	5.8	5.8	3.1	2.7
East South Central	15.0	13.0	10.2	8.9	5.6 <sup>a</sup>	3.7
West South Central	8.4	7.3	5.5	4.5	2.6	2.2
West	5.5	5.3	3.6	3.4	1.4	1.2
Mountain	7.4	7.9	4.6	5.5	2.0	2.1
Pacific	4.7	4.1	3.2	2.5	1.2	0.9
<b>COUNTY TYPE</b>						
Large Metro	5.2 <sup>a</sup>	4.5	3.6	3.1	1.4	1.3
Small Metro	8.1	8.4	5.2	5.8	2.3	2.7
250K – 1 Mil. Pop.	7.0	8.0	4.6	5.5	1.8	2.6
< 250K Pop.	10.3	9.2	6.5	6.3	3.2	3.0
Nonmetro	15.6	14.3	10.3	8.9	6.2 <sup>b</sup>	4.4
Urbanized	14.2	12.8	9.7	7.8	5.4	4.3
Less Urbanized	16.6	14.4	10.8	9.0	6.8 <sup>b</sup>	4.3
Completely Rural	16.5	21.1	10.6	14.1	6.1	5.3

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.59A (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	6,562	6,717	2,932	2,947	1,718	1,762
<b>GEOGRAPHIC DIVISION</b>						
Northeast	1,012	993	398	395	235	203
New England	291	310	111	124	54	67
Middle Atlantic	721	683	287	271	181	137
Midwest	1,857	1,928	850	857	488	553
East North Central	1,196	1,195	547	538	311	333
West North Central	661	733	303	319	177	220
South	2,302	2,333	1,048	1,081	653	672
South Atlantic	1,101	1,084	477	470	271	272
East South Central	460	459	232	234	161	168
West South Central	741	790	338	377	221	232
West	1,391	1,463	636	614	343	334
Mountain	528	580	228	245	137	146
Pacific	863	882	408	369	205	188
<b>COUNTY TYPE</b>						
Large Metro	2,722	2,784	1,095	1,099	596	595
Small Metro	2,259	2,321	1,043	1,014	608	602
250K – 1 Mil. Pop.	1,408	1,373	646	587	386	342
< 250K Pop.	851	948	397	427	222	261
Nonmetro	1,581	1,613	794	834	514	566
Urbanized	668	667	324	315	202	215
Less Urbanized	773	809	390	454	266	310
Completely Rural	140	137	81	65	46	41

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.59B (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	20.0	20.4	9.0	8.9	5.2	5.3
<b>GEOGRAPHIC DIVISION</b>						
Northeast	17.2	16.6	6.7	6.6	4.0	3.4
New England	19.0	19.9	7.3	8.0	3.5	4.3
Middle Atlantic	16.5	15.4	6.6	6.1	4.1	3.1
Midwest	25.5	26.5	11.7	11.8	6.7	7.6
East North Central	23.8	23.7	10.9	10.7	6.2	6.6
West North Central	29.2	32.7	13.4	14.3	7.8	9.8
South	19.7	19.8	9.0	9.2	5.6	5.7
South Atlantic	18.6	18.2	8.1	7.9	4.6	4.6
East South Central	24.6	24.5	12.4	12.5	8.6	9.0
West South Central	18.9	20.1	8.6	9.6	5.7	5.9
West	17.7	18.5	8.1	7.8	4.4	4.2
Mountain	22.4	24.5	9.7	10.3	5.8	6.2
Pacific	15.7	15.9	7.4	6.7	3.7	3.4
<b>COUNTY TYPE</b>						
Large Metro	15.7	15.6	6.3	6.2	3.4	3.3
Small Metro	21.9	23.0	10.1	10.0	5.9	6.0
250K – 1 Mil. Pop.	20.8	21.0	9.5	9.0	5.7	5.2
< 250K Pop.	24.0	26.6	11.2	12.0	6.3	7.3
Nonmetro	31.0	32.3	15.6	16.7	10.1	11.3
Urbanized	29.3	29.2	14.2	13.8	8.9	9.4
Less Urbanized	31.6	34.4	15.9 <sup>a</sup>	19.3	10.9	13.2
Completely Rural	37.6	38.5	21.7	18.4	12.4	11.5

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.60A (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	36,534	37,421	6,810 <sup>a</sup>	7,661	5,722	6,388
<b>GEOGRAPHIC DIVISION</b>						
Northeast	5,065	4,895	699	817	538	704
New England	1,310	1,223	201	102	151	87
Middle Atlantic	3,755	3,673	498	715	387 <sup>a</sup>	617
Midwest	9,091	9,020	1,893	1,714	1,553	1,425
East North Central	5,985	5,937	1,158	1,198	987	1,024
West North Central	3,106	3,083	735 <sup>a</sup>	516	566	402
South	14,101	15,046	2,842 <sup>a</sup>	3,490	2,466 <sup>a</sup>	3,070
South Atlantic	7,035	7,238	1,261	1,457	1,058	1,262
East South Central	2,796	3,193	590 <sup>a</sup>	880	531 <sup>a</sup>	785
West South Central	4,269	4,615	991	1,152	877	1,023
West	8,277	8,460	1,375	1,640	1,165	1,189
Mountain	3,145	3,387	645	596	536	407
Pacific	5,132	5,073	730	1,044	629	781
<b>COUNTY TYPE</b>						
Large Metro	15,856	15,658	2,279	2,597	1,923	2,052
Small Metro	11,628	12,689	2,305	2,648	1,805	2,219
250K – 1 Mil. Pop.	6,850 <sup>a</sup>	7,828	1,263	1,601	1,002	1,244
< 250K Pop.	4,777	4,861	1,042	1,046	803	975
Nonmetro	9,050	9,075	2,226	2,416	1,994	2,117
Urbanized	3,573	3,372	887	663	799 <sup>a</sup>	558
Less Urbanized	4,394	4,780	1,071 <sup>a</sup>	1,454	945 <sup>a</sup>	1,298
Completely Rural	1,082	922	267	299	251	261

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.60B (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	19.2	19.5	3.6	4.0	3.0	3.3
<b>GEOGRAPHIC DIVISION</b>						
Northeast	14.3	13.7	2.0	2.3	1.5	2.0
New England	14.0	13.0	2.2	1.1	1.6	0.9
Middle Atlantic	14.3	13.9	1.9	2.7	1.5	2.3
Midwest	21.7	21.4	4.5	4.1	3.7	3.4
East North Central	20.4	20.1	3.9	4.1	3.4	3.5
West North Central	24.7	24.3	5.9 <sup>a</sup>	4.1	4.5	3.2
South	20.4	21.4	4.1 <sup>a</sup>	5.0	3.6 <sup>a</sup>	4.4
South Atlantic	19.0	19.3	3.4	3.9	2.9	3.4
East South Central	24.6	27.7	5.2 <sup>a</sup>	7.6	4.7	6.8
West South Central	20.6	21.8	4.8	5.4	4.2	4.8
West	19.1	19.3	3.2	3.7	2.7	2.7
Mountain	24.0	25.3	4.9	4.4	4.1	3.0
Pacific	17.0	16.6	2.4	3.4	2.1	2.6
<b>COUNTY TYPE</b>						
Large Metro	15.6	15.4	2.2	2.5	1.9	2.0
Small Metro	20.6	21.9	4.1	4.6	3.2	3.8
250K – 1 Mil. Pop.	18.1 <sup>a</sup>	20.3	3.3	4.2	2.7	3.2
< 250K Pop.	25.7	25.0	5.6	5.4	4.3	5.0
Nonmetro	28.5	28.3	7.0	7.5	6.3	6.6
Urbanized	27.9	25.7	6.9 <sup>a</sup>	5.1	6.2 <sup>a</sup>	4.3
Less Urbanized	28.9	30.5	7.1 <sup>a</sup>	9.3	6.2 <sup>a</sup>	8.3
Completely Rural	28.5	28.0	7.0	9.1	6.6	7.9

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.



**Table 2.61A (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	43,096	44,139	9,742	10,607	7,440	8,150
<b>GEOGRAPHIC DIVISION</b>						
Northeast	6,078	5,889	1,098	1,212	773	908
New England	1,601	1,533	312	227	204	154
Middle Atlantic	4,476	4,356	785	986	568	754
Midwest	10,948	10,948	2,743	2,571	2,041	1,978
East North Central	7,181	7,132	1,705	1,735	1,298	1,356
West North Central	3,767	3,816	1,038	836	743	622
South	16,403	17,379	3,890 <sup>a</sup>	4,570	3,119 <sup>a</sup>	3,742
South Atlantic	8,136	8,322	1,738	1,927	1,329	1,534
East South Central	3,256	3,652	822 <sup>a</sup>	1,114	692 <sup>a</sup>	954
West South Central	5,010	5,405	1,329	1,529	1,098	1,254
West	9,668	9,923	2,012	2,254	1,508	1,522
Mountain	3,673	3,967	873	841	673	553
Pacific	5,995	5,956	1,138	1,413	834	969
<b>COUNTY TYPE</b>						
Large Metro	18,578	18,442	3,374	3,696	2,518	2,647
Small Metro	13,887	15,009	3,348	3,661	2,413	2,821
250K – 1 Mil. Pop.	8,258 <sup>a</sup>	9,201	1,909	2,188	1,388	1,585
< 250K Pop.	5,629	5,808	1,440	1,473	1,025	1,235
Nonmetro	10,631	10,687	3,020	3,250	2,509	2,683
Urbanized	4,241	4,039	1,211	978	1,001	773
Less Urbanized	5,168	5,589	1,461 <sup>b</sup>	1,908	1,211 <sup>b</sup>	1,608
Completely Rural	1,222	1,059	348	364	297	302

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.61B (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	19.4	19.6	4.4	4.7	3.3	3.6
<b>GEOGRAPHIC DIVISION</b>						
Northeast	14.7	14.1	2.6	2.9	1.9	2.2
New England	14.7	14.0	2.9	2.1	1.9	1.4
Middle Atlantic	14.7	14.1	2.6	3.2	1.9	2.4
Midwest	22.3	22.1	5.6	5.2	4.1	4.0
East North Central	20.9	20.7	5.0	5.0	3.8	3.9
West North Central	25.4	25.6	7.0	5.6	5.0	4.2
South	20.3	21.2	4.8 <sup>a</sup>	5.6	3.9 <sup>a</sup>	4.6
South Atlantic	19.0	19.2	4.0	4.4	3.1	3.5
East South Central	24.6	27.3	6.2 <sup>a</sup>	8.3	5.2	7.1
West South Central	20.3	21.6	5.4	6.1	4.5	5.0
West	18.9	19.1	3.9	4.3	2.9	2.9
Mountain	23.7	25.1	5.6	5.3	4.4	3.5
Pacific	16.8	16.5	3.2	3.9	2.3	2.7
<b>COUNTY TYPE</b>						
Large Metro	15.6	15.4	2.8	3.1	2.1	2.2
Small Metro	20.8	22.1	5.0	5.4	3.6	4.1
250K – 1 Mil. Pop.	18.5 <sup>a</sup>	20.4	4.3	4.9	3.1	3.5
< 250K Pop.	25.4	25.3	6.5	6.4	4.6	5.4
Nonmetro	28.8	28.8	8.2	8.8	6.8	7.2
Urbanized	28.1	26.3	8.0 <sup>a</sup>	6.4	6.6 <sup>a</sup>	5.0
Less Urbanized	29.3	31.0	8.3 <sup>b</sup>	10.6	6.9 <sup>a</sup>	8.9
Completely Rural	29.3	29.0	8.3	10.0	7.1	8.3

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.