

**Table 2.47A (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	172,420	174,000	85,017	84,550	71,073	70,955
<b>GEOGRAPHIC DIVISION</b>						
Northeast	31,305	31,588	14,665	14,542	12,041	12,091
New England	8,594	8,733	3,803	3,826	3,046	3,018
Middle Atlantic	22,711	22,855	10,862	10,716	8,996	9,073
Midwest	40,434	40,139	20,747	20,067	17,541	16,811
East North Central	28,056	27,993	14,323	14,313	12,208	11,973
West North Central	12,377	12,146	6,424 <sup>a</sup>	5,754	5,333	4,837
South	63,153	63,697	31,976	32,015	27,253	27,763
South Atlantic	33,667	34,002	16,654	16,157	14,213	13,924
East South Central	10,690	10,807	5,537	5,943	4,873	5,323
West South Central	18,796	18,887	9,785	9,915	8,167	8,516
West	37,528	38,577	17,630	17,926	14,237	14,290
Mountain	11,959	12,205	5,882	5,752	4,969	4,702
Pacific	25,569	26,372	11,748	12,174	9,268	9,588
<b>COUNTY TYPE</b>						
Large Metro	89,251	89,907	43,191	43,063	35,139	35,245
Small Metro	52,864	53,494	25,912	25,235	21,953	21,327
250K – 1 Mil. Pop.	35,006	34,930	16,989	16,358	14,368	13,865
< 250K Pop.	17,859	18,564	8,923	8,877	7,585	7,462
Nonmetro	30,304	30,600	15,914	16,251	13,980	14,383
Urbanized	12,424	12,879	6,597	6,540	5,693	5,722
Less Urbanized	14,591	14,786	7,708	8,217	6,885	7,286
Completely Rural	3,289	2,935	1,609	1,494	1,402	1,375

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.47B (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	69.6	69.7	34.3	33.8	28.7	28.4
<b>GEOGRAPHIC DIVISION</b>						
Northeast	68.2	68.5	32.0	31.5	26.2	26.2
New England	71.5	72.4	31.6	31.7	25.3	25.0
Middle Atlantic	67.1	67.1	32.1	31.5	26.6	26.7
Midwest	73.8	73.0	37.9	36.5	32.0	30.6
East North Central	73.3	72.9	37.4	37.3	31.9	31.2
West North Central	75.0	73.3	38.9 <sup>a</sup>	34.7	32.3	29.2
South	70.2	70.0	35.6	35.2	30.3	30.5
South Atlantic	70.8	70.8	35.0	33.6	29.9	29.0
East South Central	72.6	72.7	37.6	40.0	33.1	35.8
West South Central	68.0	67.2	35.4	35.3	29.6	30.3
West	65.6	66.7	30.8	31.0	24.9	24.7
Mountain	69.2	69.4	34.1	32.7	28.8	26.7
Pacific	64.0	65.6	29.4	30.3	23.2	23.8
<b>COUNTY TYPE</b>						
Large Metro	67.3	67.6	32.6	32.4	26.5	26.5
Small Metro	71.3	70.9	35.0	33.4	29.6	28.3
250K – 1 Mil. Pop.	70.6	69.9	34.3	32.7	29.0	27.7
< 250K Pop.	72.8	72.9	36.4	34.8	30.9	29.3
Nonmetro	73.8	74.2	38.7	39.4	34.0	34.9
Urbanized	73.9	75.0	39.3	38.1	33.9	33.3
Less Urbanized	74.2	73.7	39.2	41.0	35.0	36.3
Completely Rural	71.5	72.9	35.0	37.1	30.5	34.1

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.48A (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:**  
**Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	7,027	6,821	5,010 <sup>a</sup>	4,747	3,141 <sup>b</sup>	2,870
<b>GEOGRAPHIC DIVISION</b>						
Northeast	1,119	1,134	820	813	517	498
New England	315	299	227	227	146 <sup>a</sup>	112
Middle Atlantic	804	834	593	586	371	385
Midwest	1,683	1,584	1,235 <sup>a</sup>	1,103	783	730
East North Central	1,206	1,119	876 <sup>a</sup>	771	572	513
West North Central	476	465	358	332	211	217
South	2,774	2,610	1,941 <sup>a</sup>	1,752	1,247 <sup>b</sup>	1,049
South Atlantic	1,341	1,319	953	896	629	535
East South Central	547	488	369	331	242 <sup>a</sup>	198
West South Central	886	804	618 <sup>a</sup>	525	376	315
West	1,451	1,493	1,014	1,079	594	594
Mountain	507	511	353	378	222	199
Pacific	944	983	661	701	372	395
<b>COUNTY TYPE</b>						
Large Metro	3,369	3,230	2,406	2,260	1,439	1,332
Small Metro	2,176	2,137	1,527	1,507	949	901
250K – 1 Mil. Pop.	1,415	1,395	981	1,004	612	575
< 250K Pop.	761	742	546	503	337	325
Nonmetro	1,482	1,455	1,076	979	753 <sup>a</sup>	637
Urbanized	612	598	442	409	313	270
Less Urbanized	733	698	540	465	382 <sup>a</sup>	300
Completely Rural	137	159	95	105	58	66

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.48B (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	27.8	27.4	19.8	19.1	12.4 <sup>a</sup>	11.5
<b>GEOGRAPHIC DIVISION</b>						
Northeast	25.1	25.9	18.4	18.6	11.6	11.4
New England	27.3	26.5	19.6	20.1	12.7	9.9
Middle Atlantic	24.3	25.7	18.0	18.1	11.2	11.9
Midwest	30.0	28.8	22.0 <sup>a</sup>	20.0	13.9	13.3
East North Central	30.6	29.0	22.2 <sup>a</sup>	19.9	14.5	13.3
West North Central	28.5	28.3	21.4	20.2	12.6	13.2
South	30.4	28.8	21.3 <sup>a</sup>	19.4	13.7 <sup>b</sup>	11.6
South Atlantic	28.9	28.8	20.6	19.6	13.6	11.7
East South Central	36.9	33.2	24.9	22.5	16.3 <sup>a</sup>	13.5
West South Central	29.5	26.7	20.6 <sup>a</sup>	17.5	12.5	10.5
West	24.0	25.1	16.8	18.1	9.8	10.0
Mountain	28.1	28.3	19.5	20.9	12.3	11.0
Pacific	22.3	23.7	15.6	16.9	8.8	9.5
<b>COUNTY TYPE</b>						
Large Metro	24.6	24.3	17.6	17.0	10.5	10.0
Small Metro	29.5	28.8	20.7	20.3	12.9	12.2
250K – 1 Mil. Pop.	28.4	28.5	19.7	20.5	12.3	11.7
< 250K Pop.	31.8	29.5	22.8	20.0	14.1	13.0
Nonmetro	35.5	34.8	25.8	23.4	18.1 <sup>b</sup>	15.2
Urbanized	35.6	33.5	25.7	22.9	18.2 <sup>a</sup>	15.1
Less Urbanized	36.2	34.6	26.6	23.1	18.9 <sup>b</sup>	14.9
Completely Rural	32.0 <sup>a</sup>	42.2	22.2	27.9	13.6	17.7

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.49A (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:**  
**Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	23,016	23,151	17,156	17,248	13,700	13,638
<b>GEOGRAPHIC DIVISION</b>						
Northeast	4,065 <sup>a</sup>	4,242	3,006	3,141	2,368	2,442
New England	1,138	1,172	853	853	659	647
Middle Atlantic	2,927	3,070	2,153	2,288	1,709	1,795
Midwest	5,501	5,452	4,226	4,223	3,441	3,457
East North Central	3,800	3,718	2,939	2,879	2,400	2,349
West North Central	1,700	1,734	1,287	1,344	1,042	1,108
South	8,167	8,037	6,075	6,002	4,995	4,891
South Atlantic	4,156	4,026	3,077	2,990	2,491	2,422
East South Central	1,331	1,333	1,054	1,049	905	887
West South Central	2,680	2,678	1,945	1,963	1,599	1,582
West	5,283	5,420	3,848	3,882	2,895	2,849
Mountain	1,635	1,675	1,222	1,195	985	918
Pacific	3,648	3,745	2,626	2,687	1,911	1,931
<b>COUNTY TYPE</b>						
Large Metro	11,733	11,991	8,575	8,729	6,758	6,743
Small Metro	7,409	7,288	5,566	5,497	4,443	4,366
250K – 1 Mil. Pop.	4,766	4,660	3,603	3,520	2,879	2,813
< 250K Pop.	2,644	2,628	1,963	1,977	1,564	1,553
Nonmetro	3,874	3,873	3,015	3,022	2,499	2,529
Urbanized	1,698	1,804	1,326	1,377	1,095	1,108
Less Urbanized	1,878	1,794	1,456	1,442	1,206	1,245
Completely Rural	298	275	233	202	197	176

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.49B (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	70.3	70.3	52.4	52.4	41.9	41.4
<b>GEOGRAPHIC DIVISION</b>						
Northeast	68.9	70.9	50.9	52.5	40.1	40.8
New England	74.5	75.3	55.9	54.8	43.2	41.5
Middle Atlantic	66.9	69.3	49.2	51.6	39.1	40.5
Midwest	75.5	74.9	58.0	58.0	47.2	47.5
East North Central	75.6	73.8	58.5	57.2	47.7	46.6
West North Central	75.2	77.4	56.9	60.0	46.1	49.5
South	69.9	68.3	52.0	51.0	42.7	41.6
South Atlantic	70.4	67.5	52.1	50.1	42.2	40.6
East South Central	71.2	71.2	56.4	56.0	48.4	47.3
West South Central	68.5	68.2	49.7	50.0	40.9	40.3
West	67.3	68.5	49.0	49.1	36.9	36.0
Mountain	69.4	70.7	51.9	50.4	41.8	38.7
Pacific	66.4	67.6	47.8	48.5	34.8	34.8
<b>COUNTY TYPE</b>						
Large Metro	67.8	67.2	49.5	48.9	39.0	37.8
Small Metro	71.8	72.2	53.9	54.4	43.0	43.2
250K – 1 Mil. Pop.	70.4	71.2	53.2	53.8	42.5	43.0
< 250K Pop.	74.4	73.9	55.2	55.6	44.0	43.7
Nonmetro	76.0	77.6	59.1	60.5	49.0	50.7
Urbanized	74.5 <sup>a</sup>	79.0	58.2	60.3	48.0	48.5
Less Urbanized	76.7	76.2	59.5	61.3	49.3	52.9
Completely Rural	80.2	77.4	62.7	57.0	53.0	49.6

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.50A (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	142,377	144,028	62,852	62,555	54,232	54,446
<b>GEOGRAPHIC DIVISION</b>						
Northeast	26,121	26,212	10,838	10,589	9,156	9,151
New England	7,141	7,262	2,723	2,747	2,240	2,259
Middle Atlantic	18,980	18,950	8,116	7,842	6,916	6,892
Midwest	33,250	33,103	15,286	14,741	13,317	12,624
East North Central	23,049	23,155	10,507	10,663	9,236	9,112
West North Central	10,201	9,948	4,779 <sup>b</sup>	4,079	4,081 <sup>a</sup>	3,512
South	52,212	53,049	23,960	24,261	21,011	21,824
South Atlantic	28,170	28,657	12,624	12,270	11,094	10,967
East South Central	8,813	8,986	4,113	4,563	3,725 <sup>a</sup>	4,238
West South Central	15,230	15,406	7,222	7,427	6,192	6,619
West	30,794	31,664	12,767	12,964	10,747	10,848
Mountain	9,816	10,020	4,307	4,179	3,762	3,586
Pacific	20,978	21,644	8,460	8,785	6,985	7,262
<b>COUNTY TYPE</b>						
Large Metro	74,149	74,687	32,210	32,074	26,942	27,170
Small Metro	43,279	44,069	18,819	18,231	16,562	16,060
250K – 1 Mil. Pop.	28,825	28,875	12,405	11,835	10,878	10,476
< 250K Pop.	14,454	15,194	6,414	6,397	5,684	5,584
Nonmetro	24,948	25,272	11,823	12,250	10,728	11,217
Urbanized	10,114	10,477	4,829	4,754	4,285	4,344
Less Urbanized	11,980	12,294	5,712	6,309	5,296	5,740
Completely Rural	2,854	2,501	1,281	1,187	1,147	1,133

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.50B (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	75.0	75.0	33.1	32.6	28.6	28.4
<b>GEOGRAPHIC DIVISION</b>						
Northeast	73.5	73.3	30.5	29.6	25.8	25.6
New England	76.5	77.5	29.1	29.3	24.0	24.1
Middle Atlantic	72.5	71.9	31.0	29.7	26.4	26.1
Midwest	79.4	78.5	36.5	35.0	31.8	29.9
East North Central	78.6	78.5	35.8	36.2	31.5	30.9
West North Central	81.2	78.4	38.0 <sup>b</sup>	32.1	32.5 <sup>a</sup>	27.7
South	75.5	75.6	34.7	34.6	30.4	31.1
South Atlantic	76.1	76.5	34.1	32.7	30.0	29.3
East South Central	77.4	78.0	36.1	39.6	32.7	36.8
West South Central	73.5	72.8	34.9	35.1	29.9	31.3
West	71.1	72.1	29.5	29.5	24.8	24.7
Mountain	74.9	74.7	32.9	31.2	28.7	26.7
Pacific	69.4	70.9	28.0	28.8	23.1	23.8
<b>COUNTY TYPE</b>						
Large Metro	72.9	73.3	31.7	31.5	26.5	26.7
Small Metro	76.7	76.0	33.4	31.5	29.4	27.7
250K – 1 Mil. Pop.	76.3	74.9	32.8	30.7	28.8	27.2
< 250K Pop.	77.7	78.3	34.5	33.0	30.6	28.8
Nonmetro	78.4	78.8	37.2	38.2	33.7	35.0
Urbanized	78.9	80.0	37.7	36.3	33.4	33.2
Less Urbanized	78.8	78.4	37.6	40.2	34.9	36.6
Completely Rural	75.1	75.9	33.7	36.0	30.2	34.4

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.



**Table 2.51A (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	165,393	167,179	80,008	79,803	67,932	68,085
<b>GEOGRAPHIC DIVISION</b>						
Northeast	30,186	30,454	13,845	13,730	11,524	11,593
New England	8,279	8,434	3,576	3,600	2,899	2,905
Middle Atlantic	21,907	22,020	10,269	10,130	8,625	8,688
Midwest	38,751	38,555	19,512	18,964	16,759	16,081
East North Central	26,850	26,873	13,446	13,542	11,636	11,461
West North Central	11,901	11,682	6,066 <sup>a</sup>	5,423	5,123	4,620
South	60,379	61,087	30,035	30,263	26,006	26,714
South Atlantic	32,326	32,683	15,701	15,261	13,585	13,388
East South Central	10,143	10,320	5,167	5,612	4,631	5,125
West South Central	17,910	18,084	9,167	9,390	7,791	8,201
West	36,077	37,084	16,616	16,846	13,642	13,697
Mountain	11,452	11,695	5,530	5,373	4,747	4,503
Pacific	24,625	25,389	11,086	11,473	8,895	9,193
<b>COUNTY TYPE</b>						
Large Metro	85,882	86,678	40,785	40,803	33,700	33,913
Small Metro	50,689	51,357	24,385	23,728	21,005	20,426
250K – 1 Mil. Pop.	33,591	33,535	16,008	15,354	13,756	13,290
< 250K Pop.	17,097	17,822	8,376	8,373	7,248	7,137
Nonmetro	28,822	29,145	14,838	15,272	13,227	13,746
Urbanized	11,812	12,280	6,155	6,131	5,380	5,452
Less Urbanized	13,858	14,088	7,168	7,751	6,503	6,985
Completely Rural	3,152	2,776	1,514	1,389	1,344	1,309

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 2.51B (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2007 and 2008**

<b>Geographic Characteristic</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	74.3	74.3	35.9	35.5	30.5	30.3
<b>GEOGRAPHIC DIVISION</b>						
Northeast	72.9	73.0	33.4	32.9	27.8	27.8
New England	76.2	77.2	32.9	32.9	26.7	26.6
Middle Atlantic	71.7	71.5	33.6	32.9	28.2	28.2
Midwest	78.8	78.0	39.7	38.4	34.1	32.5
East North Central	78.1	77.9	39.1	39.2	33.9	33.2
West North Central	80.3	78.2	40.9 <sup>a</sup>	36.3	34.6 <sup>a</sup>	30.9
South	74.7	74.6	37.2	36.9	32.2	32.6
South Atlantic	75.3	75.2	36.6	35.1	31.6	30.8
East South Central	76.5	77.0	39.0	41.9	34.9	38.3
West South Central	72.7	72.1	37.2	37.4	31.6	32.7
West	70.5	71.5	32.5	32.5	26.7	26.4
Mountain	74.0	74.1	35.7	34.1	30.7	28.5
Pacific	68.9	70.4	31.0	31.8	24.9	25.5
<b>COUNTY TYPE</b>						
Large Metro	72.2	72.4	34.3	34.1	28.3	28.3
Small Metro	76.0	75.5	36.5	34.9	31.5	30.0
250K – 1 Mil. Pop.	75.4	74.4	35.9	34.0	30.9	29.5
< 250K Pop.	77.2	77.6	37.8	36.5	32.7	31.1
Nonmetro	78.1	78.6	40.2	41.2	35.8	37.1
Urbanized	78.3	79.9	40.8	39.9	35.7	35.5
Less Urbanized	78.5	78.1	40.6	43.0	36.9	38.7
Completely Rural	75.6	76.1	36.3	38.1	32.2	35.8

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2008 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>a</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.