

Table 7.11A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2007 and 2008

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLICIT DRUGS²	5,330	5,082	6,547	6,703	3,809	4,346	4,171	3,945
Marijuana and Hashish	4,268	4,209	5,071	5,321	2,782 ^a	3,351	2,327	2,322
Cocaine	932	890	611	660	263	188	270 ^a	117
Crack	204	143	135	124	107	50	164 ^a	43
Heroin	29 ^a	82	17 ^a	75	65	28	42	27
Hallucinogens	367	446	363	360	126	123	139	131
LSD	51	68	55	47	20	11	18	28
PCP	4	9	15	9	4	4	18	*
Ecstasy	181	251	180	207	92	51	51	46
Inhalants	149	77	159	222	77	133	231	209
Nonmedical Use of Psychotherapeutics ^{3,4}	1,780	1,555	1,884	1,856	1,324	1,096	1,908	1,717
Pain Relievers	1,452	1,238	1,348	1,432	894	758	1,480	1,319
OxyContin [®]	108	117	118	139	36	64	106	115
Tranquilizers	580	470	500	587	372	360	382	383
Stimulants ⁴	315	302	302	298	191	103	244	200
Methamphetamine ⁴	154	77	152	113	93 ^a	20	130	104
Sedatives	77	56	117	69	57	41	95	67
ILLICIT DRUGS OTHER THAN MARIJUANA²	2,615	2,385	2,657	2,674	1,634	1,445	2,365	2,062

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.11B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2007 and 2008

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	31.3	29.4	16.1	16.4	5.5	6.1	3.4	3.3
Marijuana and Hashish	25.1	24.3	12.4	13.0	4.0	4.7	1.9	1.9
Cocaine	5.5	5.1	1.5	1.6	0.4	0.3	0.2 ^a	0.1
Crack	1.2	0.8	0.3	0.3	0.2	0.1	0.1 ^a	0.0
Heroin	0.2 ^a	0.5	0.0 ^a	0.2	0.1	0.0	0.0	0.0
Hallucinogens	2.2	2.6	0.9	0.9	0.2	0.2	0.1	0.1
LSD	0.3	0.4	0.1	0.1	0.0	0.0	0.0	0.0
PCP	0.0	0.1	0.0	0.0	0.0	0.0	0.0	*
Ecstasy	1.1	1.5	0.4	0.5	0.1	0.1	0.0	0.0
Inhalants	0.9	0.4	0.4	0.5	0.1	0.2	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{3,4}	10.5	9.0	4.6	4.5	1.9	1.5	1.6	1.4
Pain Relievers	8.5	7.2	3.3	3.5	1.3	1.1	1.2	1.1
OxyContin [®]	0.6	0.7	0.3	0.3	0.1	0.1	0.1	0.1
Tranquilizers	3.4	2.7	1.2	1.4	0.5	0.5	0.3	0.3
Stimulants ⁴	1.9	1.7	0.7	0.7	0.3 ^a	0.1	0.2	0.2
Methamphetamine ⁴	0.9	0.4	0.4	0.3	0.1 ^a	0.0	0.1	0.1
Sedatives	0.5	0.3	0.3	0.2	0.1	0.1	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	15.4	13.8	6.5	6.6	2.4	2.0	2.0	1.7

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.12A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2007 and 2008

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	355	336	704	724	296	332	1,053 ^a	913
Marijuana and Hashish	317	284	593	608	226	274	550	490
Cocaine	33	42	37	34	9	8	24	13
Crack	6	2	4	4	1	*	5	*
Heroin	1	4	*	7	1	*	*	3
Hallucinogens	44	59	74	95	18	29	39	59
LSD	6	10	12	18	3	6	8	16
PCP	1	6	4	2	1	4	3	2
Ecstasy	20	30	24	42	8	4	13	28
Inhalants	36	25	60	58	27	28	172	157
Nonmedical Use of Psychotherapeutics ^{3,4}	144	136	194	186	90	60	402	328
Pain Relievers	126	103	154	143	83 ^a	46	306	275
OxyContin [®]	18	13	13	21	7	1	9	14
Tranquilizers	54	33	57	51	12	12	59	53
Stimulants ⁴	26	36	36	37	9	10	56 ^a	30
Methamphetamine ⁴	13	15	6	10	2	*	13	3
Sedatives	3	10	4	7	2	3	26	15
ILLCIT DRUGS OTHER THAN MARIJUANA²	177	187	290	294	122	106	592	504

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.12B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2007 and 2008

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	60.1 ^a	68.5	37.9 ^a	42.6	18.8	22.9	5.0 ^a	4.3
Marijuana and Hashish	53.7	57.9	32.0	35.8	14.3 ^a	18.9	2.6	2.3
Cocaine	5.5	8.6	2.0	2.0	0.6	0.6	0.1	0.1
Crack	1.1	0.5	0.2	0.3	0.1	*	0.0	*
Heroin	0.2	0.9	*	0.4	0.1	*	*	0.0
Hallucinogens	7.5	12.0	4.0	5.6	1.2	2.0	0.2	0.3
LSD	1.0	2.0	0.7	1.0	0.2	0.4	0.0	0.1
PCP	0.1	1.2	0.2	0.1	0.1	0.3	0.0	0.0
Ecstasy	3.4	6.2	1.3	2.5	0.5	0.3	0.1	0.1
Inhalants	6.2	5.0	3.2	3.4	1.7	1.9	0.8	0.7
Nonmedical Use of Psychotherapeutics ^{3,4}	24.3	27.7	10.4	11.0	5.7	4.1	1.9 ^a	1.5
Pain Relievers	21.3	20.9	8.3	8.4	5.3 ^a	3.2	1.4	1.3
OxyContin [®]	3.0	2.7	0.7	1.2	0.4	0.1	0.0	0.1
Tranquilizers	9.1	6.8	3.1	3.0	0.8	0.8	0.3	0.3
Stimulants ⁴	4.4	7.3	2.0	2.2	0.6	0.7	0.3 ^a	0.1
Methamphetamine ⁴	2.2	3.1	0.3	0.6	0.1	*	0.1	0.0
Sedatives	0.5	2.1	0.2	0.4	0.1	0.2	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	29.9 ^a	38.1	15.6	17.3	7.8	7.3	2.8	2.4

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.13A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2007 and 2008

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	2,331	2,227	2,482	2,389	882 ^a	1,047	769	802
Marijuana and Hashish	2,021	1,936	2,079	2,017	729 ^a	891	550	581
Cocaine	309	294	199	154	38	28	26	31
Crack	24	26	14	15	9	1	10	10
Heroin	8 ^b	38	17	20	13	2	9	14
Hallucinogens	235	271	171	188	36	55	33	36
LSD	41	56	19	30	1	5	10	7
PCP	4	4	2	0	*	*	9	*
Ecstasy	91 ^a	156	101	106	21	25	16	17
Inhalants	43	41	49	44	12	13	25	6
Nonmedical Use of Psychotherapeutics ^{3,4}	747	748	698	676	239	249	266	270
Pain Relievers	595	574	538	517	168	192	216	218
OxyContin [®]	58	56	59	35	19	23	20	15
Tranquilizers	269	245	167	185	76	85	59	55
Stimulants ⁴	146	160	120	133	49	22	37	32
Methamphetamine ⁴	44	24	38	28	21	4	12	17
Sedatives	38 ^a	16	21	22	3	14	10	12
ILLCIT DRUGS OTHER THAN MARIJUANA²	1,067	1,051	957	898	295	318	324	318

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.13B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2007 and 2008

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	48.4	46.8	28.0	27.3	13.9	15.8	6.1	6.3
Marijuana and Hashish	41.9	40.7	23.5	23.1	11.5 ^a	13.4	4.3	4.5
Cocaine	6.4	6.2	2.2	1.8	0.6	0.4	0.2	0.2
Crack	0.5	0.5	0.2	0.2	0.1	0.0	0.1	0.1
Heroin	0.2 ^b	0.8	0.2	0.2	0.2	0.0	0.1	0.1
Hallucinogens	4.9	5.7	1.9	2.1	0.6	0.8	0.3	0.3
LSD	0.9	1.2	0.2	0.3	0.0	0.1	0.1	0.1
PCP	0.1	0.1	0.0	0.0	*	*	0.1	*
Ecstasy	1.9 ^b	3.3	1.1	1.2	0.3	0.4	0.1	0.1
Inhalants	0.9	0.9	0.6	0.5	0.2	0.2	0.2 ^a	0.1
Nonmedical Use of Psychotherapeutics ^{3,4}	15.5	15.7	7.9	7.7	3.8	3.7	2.1	2.1
Pain Relievers	12.3	12.1	6.1	5.9	2.6	2.9	1.7	1.7
OxyContin [®]	1.2	1.2	0.7	0.4	0.3	0.3	0.2	0.1
Tranquilizers	5.6	5.1	1.9	2.1	1.2	1.3	0.5	0.4
Stimulants ⁴	3.0	3.4	1.4	1.5	0.8 ^a	0.3	0.3	0.2
Methamphetamine ⁴	0.9	0.5	0.4	0.3	0.3	0.1	0.1	0.1
Sedatives	0.8 ^a	0.3	0.2	0.3	0.1	0.2	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	22.1	22.1	10.8	10.3	4.6	4.8	2.5	2.5

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.14A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2007 and 2008

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	2,643	2,520	3,361	3,591	2,630	2,968	2,349	2,230
Marijuana and Hashish	1,931	1,988	2,399	2,696	1,828	2,186	1,227	1,250
Cocaine	590	554	374	472	215	152	220 ^a	72
Crack	174	115	117	104	97	48	149 ^a	32
Heroin	19	40	*	49	50	26	32	11
Hallucinogens	88	116	118	77	72	39	68	37
LSD	4	2	24	*	16	*	*	5
PCP	*	*	9	6	*	*	6	*
Ecstasy	69	64	55	60	63	22	22	*
Inhalants	69	11	51	120	38	92	34	45
Nonmedical Use of Psychotherapeutics ^{3,4}	889	671	992	993	994	787	1,241	1,119
Pain Relievers	731	561	656	772	643	520	957	825
OxyContin [®]	33	47	45	84	11	40	78	86
Tranquilizers	258	191	276	351	285	264	263	275
Stimulants ⁴	143	106	145	128	132	71	152	138
Methamphetamine ⁴	98	37	108	74	70	16	104	84
Sedatives	36	29	93	40	52	25	59	40
ILLCIT DRUGS OTHER THAN MARIJUANA²	1,371	1,147	1,409	1,482	1,217	1,020	1,449	1,240

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.14B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2007 and 2008

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2007	2008	2007	2008	2007	2008	2007	2008
ILLCIT DRUGS²	22.8	20.9	11.2	11.8	4.3	4.7	2.7	2.6
Marijuana and Hashish	16.6	16.5	8.0	8.9	3.0	3.5	1.4	1.4
Cocaine	5.1	4.6	1.2	1.6	0.4	0.2	0.3 ^a	0.1
Crack	1.5	1.0	0.4	0.3	0.2	0.1	0.2 ^a	0.0
Heroin	0.2	0.3	*	0.2	0.1	0.0	0.0	0.0
Hallucinogens	0.8	1.0	0.4	0.3	0.1	0.1	0.1	0.0
LSD	0.0	0.0	0.1	*	0.0	*	*	0.0
PCP	*	*	0.0	0.0	*	*	0.0	*
Ecstasy	0.6	0.5	0.2	0.2	0.1	0.0	0.0	*
Inhalants	0.6	0.1	0.2	0.4	0.1	0.1	0.0	0.1
Nonmedical Use of Psychotherapeutics ^{3,4}	7.7 ^a	5.6	3.3	3.3	1.6	1.3	1.4	1.3
Pain Relievers	6.3	4.7	2.2	2.5	1.1	0.8	1.1	1.0
OxyContin [®]	0.3	0.4	0.2	0.3	0.0	0.1	0.1	0.1
Tranquilizers	2.2	1.6	0.9	1.2	0.5	0.4	0.3	0.3
Stimulants ⁴	1.2	0.9	0.5	0.4	0.2	0.1	0.2	0.2
Methamphetamine ⁴	0.8	0.3	0.4	0.2	0.1	0.0	0.1	0.1
Sedatives	0.3	0.2	0.3	0.1	0.1	0.0	0.1	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA²	11.8	9.5	4.7	4.9	2.0	1.6	1.7	1.4

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.15A Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Numbers in Thousands, 2007 and 2008

Drug	Total (2007)	Total (2008)	Aged 12-20 (2007)	Aged 12-20 (2008)	Aged 21+ (2007)	Aged 21+ (2008)
ILLCIT DRUGS¹	15,686	16,132	3,751	3,772	11,935	12,360
Marijuana or Hashish	12,122	12,881	3,224	3,280	8,897	9,601
Cocaine (Including Crack)	1,806	1,738	300	278	1,505	1,460
Heroin	111	186	16	33	95	153
Hallucinogens	856	929	360 ^a	460	496	469
Inhalants	385	432	180	174	205	258
Pain Relievers ²	3,694	3,428	914	829	2,780	2,599
Tranquilizers ²	1,453	1,417	315	290	1,138	1,128
Stimulants ^{2,3}	809	704	190	195	619	509
Methamphetamine ³	400 ^a	210	53	35	347 ^a	175
Sedatives ²	251	166	28	25	224	141
ILLCIT DRUGS OTHER THAN MARIJUANA¹	6,905	6,503	1,562	1,530	5,343	4,973

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.15B Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Percentages, 2007 and 2008

Drug	Total (2007)	Total (2008)	Aged 12-20 (2007)	Aged 12-20 (2008)	Aged 21+ (2007)	Aged 21+ (2008)
ILLCIT DRUGS¹	12.4	12.5	35.0 ^a	37.5	10.3	10.4
Marijuana or Hashish	9.6	10.0	30.0 ^a	32.6	7.7	8.1
Cocaine (Including Crack)	1.4	1.3	2.8	2.8	1.3	1.2
Heroin	0.1	0.1	0.1	0.3	0.1	0.1
Hallucinogens	0.7	0.7	3.4 ^b	4.6	0.4	0.4
Inhalants	0.3	0.3	1.7	1.7	0.2	0.2
Pain Relievers ²	2.9	2.7	8.5	8.2	2.4	2.2
Tranquilizers ²	1.1	1.1	2.9	2.9	1.0	0.9
Stimulants ^{2,3}	0.6	0.5	1.8	1.9	0.5	0.4
Methamphetamine ³	0.3 ^a	0.2	0.5	0.3	0.3 ^a	0.1
Sedatives ²	0.2	0.1	0.3	0.2	0.2	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	5.4	5.0	14.6	15.2	4.6	4.2

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.16A Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: Numbers in Thousands, 2007 and 2008

Drug¹	Total (2007)	Total (2008)	Aged 12-20 (2007)	Aged 12-20 (2008)	Aged 21+ (2007)	Aged 21+ (2008)
ILLCIT DRUGS²	6,951	7,205	1,716	1,722	5,235	5,483
Marijuana or Hashish	5,983	6,396	1,616	1,627	4,367	4,769
Cocaine (Including Crack)	614	542	71	82	543	459
Heroin	38	53	6	3	32	50
Hallucinogens	99	64	35	39	63	24
Inhalants	16	32	12	20	*	12
Pain Relievers ³	519	554	84	79	435	475
Tranquilizers ³	89 ^a	171	19	13	69 ^a	158
Stimulants ³	39	58	11	9	29	49
Methamphetamine ⁴	123	46	11	6	111	40
Sedatives ³	49	15	2	2	47	13
ILLCIT DRUGS OTHER THAN MARIJUANA²	1,386	1,302	217	212	1,170	1,090

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.16B Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: Percentages, 2007 and 2008

Drug¹	Total (2007)	Total (2008)	Aged 12-20 (2007)	Aged 12-20 (2008)	Aged 21+ (2007)	Aged 21+ (2008)
ILLCIT DRUGS²	5.5	5.6	16.3	17.4	4.5	4.6
Marijuana or Hashish	4.7	5.0	15.3	16.5	3.8	4.0
Cocaine (Including Crack)	0.5	0.4	0.7	0.8	0.5	0.4
Heroin	0.0	0.0	0.1	0.0	0.0	0.0
Hallucinogens	0.1	0.0	0.3	0.4	0.1	0.0
Inhalants	0.0	0.0	0.1	0.2	*	0.0
Pain Relievers ³	0.4	0.4	0.8	0.8	0.4	0.4
Tranquilizers ³	0.1	0.1	0.2	0.1	0.1 ^a	0.1
Stimulants ³	0.0	0.0	0.1	0.1	0.0	0.0
Methamphetamine ⁴	0.1 ^a	0.0	0.1	0.1	0.1	0.0
Sedatives ³	0.0	0.0	0.0	0.0	0.0	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA²	1.1	1.0	2.0	2.1	1.0	0.9

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.17A Types of Illicit Drugs Used in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2007 and 2008

Drug	Total (2007)	Total (2008)	Aged 12-17 (2007)	Aged 12-17 (2008)	Aged 18-25 (2007)	Aged 18-25 (2008)	Aged 26+ (2007)	Aged 26+ (2008)
ILLCIT DRUGS¹	15,686	16,132	1,356	1,392	5,695	5,663	8,635	9,078
Marijuana or Hashish	12,122	12,881	1,136	1,166	4,829	4,845	6,157	6,870
Cocaine (Including Crack)	1,806	1,738	79	84	547	475	1,180	1,178
Heroin	111	186	3	11	38	60	70	115
Hallucinogens	856	929	136 ^a	183	442	514	278	232
Inhalants	385	432	123	111	105	97	157	223
Pain Relievers ²	3,694	3,428	363 ^a	292	1,301	1,283	2,030	1,853
Tranquilizers ²	1,453	1,417	123	96	512	515	818	806
Stimulants ^{2,3}	809	704	72	82	316	316	421	305
Methamphetamine ³	400 ^a	210	20	25	103 ^a	57	276 ^a	127
Sedatives ²	251	166	9	20	62	52	180	94
ILLCIT DRUGS OTHER THAN MARIJUANA¹	6,905	6,503	589	587	2,319	2,267	3,998	3,649

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.17B Types of Illicit Drugs Used in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Percentages, 2007 and 2008

Drug	Total (2007)	Total (2008)	Aged 12-17 (2007)	Aged 12-17 (2008)	Aged 18-25 (2007)	Aged 18-25 (2008)	Aged 26+ (2007)	Aged 26+ (2008)
ILLCIT DRUGS¹	12.4	12.5	33.7 ^b	38.3	28.4	28.1	8.4	8.6
Marijuana or Hashish	9.6	10.0	28.3 ^a	32.1	24.1	24.0	6.0	6.5
Cocaine (Including Crack)	1.4	1.3	2.0	2.3	2.7	2.4	1.1	1.1
Heroin	0.1	0.1	0.1	0.3	0.2	0.3	0.1	0.1
Hallucinogens	0.7	0.7	3.4 ^b	5.0	2.2	2.5	0.3	0.2
Inhalants	0.3	0.3	3.0	3.1	0.5	0.5	0.2	0.2
Pain Relievers ²	2.9	2.7	9.0	8.0	6.5	6.4	2.0	1.8
Tranquilizers ²	1.1	1.1	3.1	2.6	2.6	2.6	0.8	0.8
Stimulants ^{2,3}	0.6	0.5	1.8	2.3	1.6	1.6	0.4	0.3
Methamphetamine ³	0.3 ^a	0.2	0.5	0.7	0.5 ^a	0.3	0.3 ^a	0.1
Sedatives ²	0.2	0.1	0.2	0.5	0.3	0.3	0.2	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	5.4	5.0	14.6	16.1	11.6	11.3	3.9	3.5

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.18A Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2007 and 2008

Drug¹	Total (2007)	Total (2008)	Aged 12-17 (2007)	Aged 12-17 (2008)	Aged 18-25 (2007)	Aged 18-25 (2008)	Aged 26+ (2007)	Aged 26+ (2008)
ILLICIT DRUGS²	6,951	7,205	580	554	2,739	2,725	3,632	3,926
Marijuana or Hashish	5,983	6,396	539	510	2,535	2,539	2,909	3,348
Cocaine (Including Crack)	614	542	17	23	134	142	463	377
Heroin	38	53	2	2	11	12	25	39
Hallucinogens	99	64	20	16	41	39	38	9
Inhalants	16	32	8	11	6	10	*	11
Pain Relievers ³	519	554	33	35	198	166	288	353
Tranquilizers ³	89 ^a	171	12	3	46	47	31 ^a	121
Stimulants ³	39	58	3	5	22	26	14	27
Methamphetamine ⁴	123	46	3	4	32	14	87	29
Sedatives ³	49	15	2	2	13	*	33	13
ILLICIT DRUGS OTHER THAN MARIJUANA²	1,386	1,302	79	74	420	395	888	833

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.18B Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Percentages, 2007 and 2008

Drug¹	Total (2007)	Total (2008)	Aged 12-17 (2007)	Aged 12-17 (2008)	Aged 18-25 (2007)	Aged 18-25 (2008)	Aged 26+ (2007)	Aged 26+ (2008)
ILLICIT DRUGS²	5.5	5.6	14.8	15.7	13.8	13.6	3.5	3.7
Marijuana or Hashish	4.7	5.0	13.7	14.5	12.7	12.7	2.8	3.2
Cocaine (Including Crack)	0.5	0.4	0.4	0.6	0.7	0.7	0.5	0.4
Heroin	0.0	0.0	0.1	0.1	0.1	0.1	0.0	0.0
Hallucinogens	0.1	0.0	0.5	0.4	0.2	0.2	0.0	0.0
Inhalants	0.0	0.0	0.2	0.3	0.0	0.0	*	0.0
Pain Relievers ³	0.4	0.4	0.8	1.0	1.0	0.8	0.3	0.3
Tranquilizers ³	0.1	0.1	0.3	0.1	0.2	0.2	0.0 ^a	0.1
Stimulants ³	0.0	0.0	0.1	0.1	0.1	0.1	0.0	0.0
Methamphetamine ⁴	0.1 ^a	0.0	0.1	0.1	0.2	0.1	0.1	0.0
Sedatives ³	0.0	0.0	0.1	0.1	0.1	*	0.0	0.0
ILLICIT DRUGS OTHER THAN MARIJUANA²	1.1	1.0	2.0	2.1	2.1	2.0	0.9	0.8

*Low precision; no estimate reported.

^a Difference between estimate and 2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2008 estimate is statistically significant at the 0.01 level.

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.