

Table 8.22C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2008

Gender/Substance	2002	2003	2004	2005	2006	2007	2008
TOTAL							
TOBACCO PRODUCTS¹	832	803	794	845	848	844	866
Cigarettes	795	789	768	777	804	808	806
Smokeless Tobacco	279	284	245	267	284	281	310
Cigars	350	331	323	354	352	337	364
Pipe Tobacco	166	131	147	146	166	170	147
ALCOHOL	989	919	957	975	969	1,018	981
Binge Alcohol Use ²	732	682	698	701	746	767	733
Heavy Alcohol Use ²	392	380	382	379	427	429	412
MALE							
TOBACCO PRODUCTS¹	572	573	570	576	584	595	627
Cigarettes	527	545	537	515	543	574	564
Smokeless Tobacco	264	256	234	241	273	251	294
Cigars	327	305	290	324	319	311	332
Pipe Tobacco	139	127	139	141	160	168	132
ALCOHOL	656	599	626	621	611	667	632
Binge Alcohol Use ²	540	517	513	529	559	585	585
Heavy Alcohol Use ²	346	321	319	321	355	375	355
FEMALE							
TOBACCO PRODUCTS¹	513	507	497	543	511	511	556
Cigarettes	501	488	491	538	502	493	551
Smokeless Tobacco	100	126	65	87	76	113	75
Cigars	114	123	122	112	131	116	114
Pipe Tobacco	90	35	43	42	47	30	55
ALCOHOL	644	600	664	637	644	674	670
Binge Alcohol Use ²	415	382	390	402	399	394	396
Heavy Alcohol Use ²	177	174	174	163	168	171	193

*Low precision; no estimate reported.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, 2007, and 2008.

Table 8.22D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Standard Errors of Percentages, 2002-2008

Gender/Substance	2002	2003	2004	2005	2006	2007	2008
TOTAL							
TOBACCO PRODUCTS¹	0.35	0.34	0.33	0.35	0.34	0.34	0.35
Cigarettes	0.34	0.33	0.32	0.32	0.33	0.33	0.32
Smokeless Tobacco	0.12	0.12	0.10	0.11	0.12	0.11	0.12
Cigars	0.15	0.14	0.13	0.15	0.14	0.14	0.15
Pipe Tobacco	0.07	0.06	0.06	0.06	0.07	0.07	0.06
ALCOHOL	0.42	0.39	0.40	0.40	0.39	0.41	0.39
Binge Alcohol Use ²	0.31	0.29	0.29	0.29	0.30	0.31	0.29
Heavy Alcohol Use ²	0.17	0.16	0.16	0.16	0.17	0.17	0.16
MALE							
TOBACCO PRODUCTS¹	0.50	0.50	0.49	0.49	0.49	0.50	0.52
Cigarettes	0.46	0.47	0.46	0.44	0.45	0.48	0.46
Smokeless Tobacco	0.23	0.22	0.20	0.20	0.23	0.21	0.24
Cigars	0.29	0.27	0.25	0.27	0.27	0.26	0.27
Pipe Tobacco	0.12	0.11	0.12	0.12	0.13	0.14	0.11
ALCOHOL	0.58	0.52	0.54	0.53	0.51	0.55	0.52
Binge Alcohol Use ²	0.48	0.45	0.44	0.45	0.47	0.49	0.48
Heavy Alcohol Use ²	0.30	0.28	0.27	0.27	0.30	0.31	0.29
FEMALE							
TOBACCO PRODUCTS¹	0.42	0.41	0.40	0.43	0.40	0.40	0.43
Cigarettes	0.41	0.40	0.40	0.43	0.40	0.39	0.43
Smokeless Tobacco	0.08	0.10	0.05	0.07	0.06	0.09	0.06
Cigars	0.09	0.10	0.10	0.09	0.10	0.09	0.09
Pipe Tobacco	0.07	0.03	0.03	0.03	0.04	0.02	0.04
ALCOHOL	0.53	0.49	0.54	0.51	0.51	0.53	0.52
Binge Alcohol Use ²	0.34	0.31	0.31	0.32	0.32	0.31	0.31
Heavy Alcohol Use ²	0.15	0.14	0.14	0.13	0.13	0.13	0.15

*Low precision; no estimate reported.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, 2007, and 2008.

Table 8.22P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: *P* Values from Tests of Differences of Percentages, 2008 Versus 2002, 2003, 2004, 2005, 2006, and 2007

Gender/Substance	2002	2003	2004	2005	2006	2007	2008
TOTAL							
TOBACCO PRODUCTS¹	0.0000	0.0039	0.0786	0.0297	0.0131	0.5856	
Cigarettes	0.0000	0.0013	0.0322	0.0311	0.0203	0.4816	
Smokeless Tobacco	0.3540	0.2008	0.0020	0.0592	0.4767	0.1817	
Cigars	0.4194	0.4672	0.0226	0.0836	0.1297	0.6303	
Pipe Tobacco	0.8173	0.3848	0.8892	0.0808	0.0336	0.4128	
ALCOHOL	0.2434	0.0043	0.0162	0.7387	0.2151	0.3948	
Binge Alcohol Use ²	0.3727	0.1233	0.2236	0.1430	0.5322	0.8932	
Heavy Alcohol Use ²	0.4507	0.5728	0.9401	0.1551	0.8882	0.8096	
MALE							
TOBACCO PRODUCTS¹	0.0008	0.0569	0.1071	0.0792	0.0132	0.3231	
Cigarettes	0.0003	0.0098	0.0368	0.0977	0.0238	0.2279	
Smokeless Tobacco	0.2350	0.0676	0.0016	0.0301	0.5558	0.1451	
Cigars	0.3108	0.9354	0.0360	0.0987	0.4367	0.7782	
Pipe Tobacco	0.6111	0.9585	0.4216	0.0261	0.0072	0.1327	
ALCOHOL	0.6851	0.6037	0.2952	0.6189	0.3106	0.1539	
Binge Alcohol Use ²	0.5877	0.3264	0.4398	0.1145	0.6173	0.8556	
Heavy Alcohol Use ²	0.7590	0.5800	0.9757	0.4603	0.8470	0.9857	
FEMALE							
TOBACCO PRODUCTS¹	0.0043	0.0141	0.3241	0.1441	0.2265	0.7876	
Cigarettes	0.0026	0.0259	0.2950	0.1445	0.2249	0.8173	
Smokeless Tobacco	0.3493	0.1799	0.8783	0.5635	0.5601	0.9413	
Cigars	0.8873	0.0384	0.2134	0.4358	0.0139	0.4813	
Pipe Tobacco	0.7035	0.0150	0.0763	0.3149	0.2269	0.0243	
ALCOHOL	0.2004	0.0002	0.0135	0.9654	0.3671	0.8867	
Binge Alcohol Use ²	0.4826	0.1951	0.2905	0.6593	0.6705	0.9945	
Heavy Alcohol Use ²	0.0394	0.9288	0.8561	0.0773	0.5074	0.5533	

*Low precision; no estimate reported.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, 2007, and 2008.

Table 8.23C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Standard Errors of Numbers in Thousands, 2002-2008

Gender/Substance	2002	2003	2004	2005	2006	2007	2008
TOTAL							
TOBACCO PRODUCTS¹	81	80	80	78	72	76	69
Cigarettes	74	73	76	71	65	67	61
Smokeless Tobacco	27	30	32	29	31	35	30
Cigars	47	44	44	46	41	44	40
Pipe Tobacco	14	18	19	17	18	19	17
ALCOHOL	80	82	81	81	82	87	78
Binge Alcohol Use ²	67	67	73	65	69	68	56
Heavy Alcohol Use ²	31	33	36	32	33	37	30
MALE							
TOBACCO PRODUCTS¹	59	56	55	59	54	57	50
Cigarettes	53	47	52	53	46	48	42
Smokeless Tobacco	26	29	30	28	29	34	28
Cigars	37	36	37	38	34	37	34
Pipe Tobacco	13	17	15	14	15	16	12
ALCOHOL	55	57	59	57	57	61	52
Binge Alcohol Use ²	46	46	52	48	47	51	40
Heavy Alcohol Use ²	24	23	27	27	26	28	23
FEMALE							
TOBACCO PRODUCTS¹	51	52	53	50	47	48	46
Cigarettes	49	50	52	48	45	45	44
Smokeless Tobacco	9	6	11	8	10	9	9
Cigars	26	23	25	24	23	22	21
Pipe Tobacco	9	7	13	11	10	11	12
ALCOHOL	58	60	59	58	59	61	56
Binge Alcohol Use ²	46	46	50	43	49	45	42
Heavy Alcohol Use ²	19	23	23	20	21	23	20

*Low precision; no estimate reported.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, 2007, and 2008.

Table 8.23D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Standard Errors of Percentages, 2002-2008

Gender/Substance	2002	2003	2004	2005	2006	2007	2008
TOTAL							
TOBACCO PRODUCTS¹	0.33	0.32	0.32	0.31	0.29	0.30	0.28
Cigarettes	0.30	0.29	0.30	0.28	0.26	0.26	0.24
Smokeless Tobacco	0.11	0.12	0.13	0.12	0.12	0.14	0.12
Cigars	0.19	0.17	0.18	0.18	0.16	0.18	0.16
Pipe Tobacco	0.06	0.07	0.08	0.07	0.07	0.08	0.07
ALCOHOL	0.32	0.33	0.32	0.32	0.32	0.34	0.31
Binge Alcohol Use ²	0.27	0.27	0.29	0.26	0.27	0.27	0.23
Heavy Alcohol Use ²	0.12	0.13	0.14	0.13	0.13	0.15	0.12
MALE							
TOBACCO PRODUCTS¹	0.47	0.44	0.43	0.46	0.41	0.44	0.39
Cigarettes	0.42	0.37	0.40	0.41	0.36	0.37	0.33
Smokeless Tobacco	0.21	0.23	0.23	0.22	0.22	0.26	0.22
Cigars	0.30	0.28	0.28	0.29	0.26	0.29	0.27
Pipe Tobacco	0.10	0.13	0.11	0.11	0.12	0.12	0.10
ALCOHOL	0.43	0.45	0.45	0.44	0.44	0.47	0.41
Binge Alcohol Use ²	0.37	0.36	0.40	0.37	0.36	0.39	0.32
Heavy Alcohol Use ²	0.19	0.18	0.21	0.21	0.20	0.22	0.18
FEMALE							
TOBACCO PRODUCTS¹	0.42	0.42	0.43	0.40	0.38	0.39	0.37
Cigarettes	0.40	0.41	0.42	0.39	0.37	0.37	0.36
Smokeless Tobacco	0.08	0.05	0.09	0.07	0.08	0.07	0.08
Cigars	0.21	0.19	0.20	0.19	0.19	0.18	0.17
Pipe Tobacco	0.07	0.06	0.10	0.09	0.08	0.09	0.10
ALCOHOL	0.48	0.49	0.48	0.47	0.47	0.49	0.46
Binge Alcohol Use ²	0.38	0.38	0.41	0.35	0.39	0.37	0.34
Heavy Alcohol Use ²	0.16	0.19	0.19	0.16	0.17	0.18	0.16

*Low precision; no estimate reported.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, 2007, and 2008.

Table 8.23P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: *P* Values from Tests of Differences of Percentages, 2008 Versus 2002, 2003, 2004, 2005, 2006, and 2007

Gender/Substance	2002	2003	2004	2005	2006	2007	2008
TOTAL							
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0001	0.0002	0.0133	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0002	0.0335	
Smokeless Tobacco	0.1596	0.3574	0.6958	0.4252	0.3259	0.1796	
Cigars	0.0040	0.0051	0.0000	0.0908	0.1465	0.0561	
Pipe Tobacco	0.0864	0.5031	0.9868	0.4016	0.6214	0.9090	
ALCOHOL	0.0000	0.0000	0.0000	0.0000	0.0000	0.0037	
Binge Alcohol Use ²	0.0000	0.0000	0.0000	0.0012	0.0000	0.0090	
Heavy Alcohol Use ²	0.0010	0.0004	0.0002	0.0112	0.0248	0.0501	
MALE							
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0061	0.0144	0.0075	
Cigarettes	0.0000	0.0000	0.0000	0.0008	0.0253	0.0377	
Smokeless Tobacco	0.1630	0.6979	0.5737	0.5345	0.2817	0.1247	
Cigars	0.0206	0.0293	0.0010	0.2541	0.6068	0.0682	
Pipe Tobacco	0.6072	0.2789	0.5628	0.6262	0.4916	0.3808	
ALCOHOL	0.0000	0.0000	0.0000	0.0058	0.0005	0.0099	
Binge Alcohol Use ²	0.0000	0.0000	0.0000	0.0031	0.0001	0.0009	
Heavy Alcohol Use ²	0.0017	0.0136	0.0017	0.0183	0.0442	0.0563	
FEMALE							
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0025	0.0033	0.4568	
Cigarettes	0.0000	0.0000	0.0000	0.0022	0.0025	0.3285	
Smokeless Tobacco	0.7726	0.0332	0.6806	0.4667	0.8856	0.6775	
Cigars	0.0601	0.0592	0.0159	0.1614	0.0666	0.4574	
Pipe Tobacco	0.0537	0.0058	0.5213	0.0704	0.1063	0.3637	
ALCOHOL	0.0000	0.0000	0.0000	0.0008	0.0022	0.1191	
Binge Alcohol Use ²	0.0167	0.0048	0.0006	0.1200	0.0206	0.8722	
Heavy Alcohol Use ²	0.1650	0.0088	0.0408	0.3282	0.2561	0.4249	

*Low precision; no estimate reported.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, 2007, and 2008.

Table 8.24C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Standard Errors of Numbers in Thousands, 2002-2008

Gender/Substance	2002	2003	2004	2005	2006	2007	2008
TOTAL							
TOBACCO PRODUCTS¹	150	151	161	155	159	162	153
Cigarettes	150	151	159	152	155	159	148
Smokeless Tobacco	59	56	60	64	65	63	68
Cigars	83	84	97	92	94	92	95
Pipe Tobacco	24	25	30	37	33	32	33
ALCOHOL	164	160	164	166	168	170	162
Binge Alcohol Use ²	161	155	166	171	169	170	166
Heavy Alcohol Use ²	112	114	120	120	125	115	122
MALE							
TOBACCO PRODUCTS¹	100	106	106	110	108	116	108
Cigarettes	99	109	105	107	108	116	105
Smokeless Tobacco	56	55	58	60	62	61	63
Cigars	73	72	83	79	79	81	79
Pipe Tobacco	22	21	27	33	30	30	29
ALCOHOL	111	100	110	101	108	110	109
Binge Alcohol Use ²	110	105	113	111	115	115	117
Heavy Alcohol Use ²	87	91	88	96	95	94	94
FEMALE							
TOBACCO PRODUCTS¹	101	93	102	101	102	96	101
Cigarettes	102	92	101	101	98	94	100
Smokeless Tobacco	8	11	10	12	13	13	14
Cigars	38	43	44	41	44	43	47
Pipe Tobacco	12	12	12	16	11	14	17
ALCOHOL	102	114	104	112	111	114	106
Binge Alcohol Use ²	99	100	96	104	103	109	101
Heavy Alcohol Use ²	55	59	60	56	63	59	62

*Low precision; no estimate reported.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, 2007, and 2008.

Table 8.24D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Standard Errors of Percentages, 2002-2008

Gender/Substance	2002	2003	2004	2005	2006	2007	2008
TOTAL							
TOBACCO PRODUCTS¹	0.48	0.48	0.50	0.48	0.49	0.49	0.47
Cigarettes	0.48	0.47	0.49	0.47	0.47	0.48	0.45
Smokeless Tobacco	0.19	0.18	0.19	0.20	0.20	0.19	0.21
Cigars	0.27	0.26	0.30	0.28	0.29	0.28	0.29
Pipe Tobacco	0.08	0.08	0.09	0.11	0.10	0.10	0.10
ALCOHOL	0.53	0.50	0.51	0.51	0.51	0.52	0.49
Binge Alcohol Use ²	0.52	0.49	0.52	0.53	0.52	0.52	0.50
Heavy Alcohol Use ²	0.36	0.36	0.37	0.37	0.38	0.35	0.37
MALE							
TOBACCO PRODUCTS¹	0.64	0.67	0.66	0.67	0.66	0.71	0.65
Cigarettes	0.64	0.69	0.65	0.66	0.66	0.70	0.63
Smokeless Tobacco	0.36	0.34	0.36	0.37	0.37	0.37	0.38
Cigars	0.47	0.45	0.51	0.48	0.48	0.49	0.47
Pipe Tobacco	0.14	0.13	0.17	0.20	0.18	0.18	0.17
ALCOHOL	0.71	0.63	0.68	0.62	0.65	0.67	0.66
Binge Alcohol Use ²	0.71	0.66	0.70	0.68	0.70	0.70	0.71
Heavy Alcohol Use ²	0.56	0.57	0.54	0.59	0.58	0.57	0.57
FEMALE							
TOBACCO PRODUCTS¹	0.65	0.59	0.64	0.63	0.63	0.59	0.61
Cigarettes	0.66	0.58	0.63	0.63	0.60	0.58	0.61
Smokeless Tobacco	0.05	0.07	0.06	0.08	0.08	0.08	0.08
Cigars	0.25	0.27	0.27	0.26	0.27	0.26	0.28
Pipe Tobacco	0.08	0.08	0.08	0.10	0.07	0.08	0.10
ALCOHOL	0.66	0.72	0.65	0.69	0.69	0.70	0.65
Binge Alcohol Use ²	0.64	0.63	0.60	0.65	0.63	0.67	0.62
Heavy Alcohol Use ²	0.35	0.37	0.37	0.35	0.39	0.36	0.38

*Low precision; no estimate reported.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, 2007, and 2008.

Table 8.24P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: *P* Values from Tests of Differences of Percentages, 2008 Versus 2002, 2003, 2004, 2005, 2006, and 2007

Gender/Substance	2002	2003	2004	2005	2006	2007	2008
TOTAL							
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0000	0.0000	0.0002	0.4754	
Cigarettes	0.0000	0.0000	0.0000	0.0000	0.0000	0.4668	
Smokeless Tobacco	0.0547	0.0103	0.1093	0.3658	0.5586	0.6583	
Cigars	0.4752	0.7557	0.0005	0.0837	0.0442	0.1919	
Pipe Tobacco	0.0129	0.0003	0.2615	0.6628	0.6315	0.1937	
ALCOHOL	0.3304	0.7961	0.3375	0.7060	0.2754	0.9543	
Binge Alcohol Use ²	0.9060	0.4332	0.7770	0.2507	0.1048	0.2659	
Heavy Alcohol Use ²	0.3801	0.2166	0.2504	0.1062	0.0348	0.5795	
MALE							
TOBACCO PRODUCTS¹	0.0003	0.0026	0.0018	0.0035	0.0221	0.2290	
Cigarettes	0.0000	0.0000	0.0000	0.0002	0.0103	0.3006	
Smokeless Tobacco	0.0954	0.0092	0.1402	0.3262	0.5342	0.5285	
Cigars	0.4950	0.8866	0.0006	0.0976	0.0368	0.0860	
Pipe Tobacco	0.1348	0.0070	0.8265	0.3415	0.6132	0.6188	
ALCOHOL	0.3590	0.0049	0.5343	0.0321	0.1020	0.3096	
Binge Alcohol Use ²	0.0730	0.0026	0.0813	0.0007	0.0627	0.1428	
Heavy Alcohol Use ²	0.1092	0.1022	0.0815	0.0222	0.1664	0.9699	
FEMALE							
TOBACCO PRODUCTS¹	0.0000	0.0000	0.0001	0.0005	0.0006	0.7669	
Cigarettes	0.0000	0.0000	0.0000	0.0002	0.0003	0.9436	
Smokeless Tobacco	0.1269	0.8290	0.3422	0.8874	0.9050	0.5234	
Cigars	0.8993	0.6487	0.2283	0.4832	0.5859	0.6561	
Pipe Tobacco	0.0228	0.0107	0.0053	0.3847	0.0303	0.0729	
ALCOHOL	0.0142	0.0233	0.0334	0.0071	0.9860	0.3447	
Binge Alcohol Use ²	0.0326	0.0417	0.1232	0.0582	0.6047	0.9303	
Heavy Alcohol Use ²	0.5390	0.9674	0.7713	0.7265	0.0458	0.3426	

*Low precision; no estimate reported.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, 2007, and 2008.

Table 8.25C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2008

Gender/Substance	2002	2003	2004	2005	2006	2007	2008
TOTAL							
TOBACCO PRODUCTS¹	797	748	752	798	798	801	845
Cigarettes	754	738	718	726	757	762	781
Smokeless Tobacco	270	274	236	256	268	266	301
Cigars	331	332	311	335	332	311	351
Pipe Tobacco	160	126	142	141	161	166	142
ALCOHOL	953	887	933	936	915	985	955
Binge Alcohol Use ²	702	639	654	650	697	712	700
Heavy Alcohol Use ²	355	349	356	347	393	403	389
MALE							
TOBACCO PRODUCTS¹	553	546	552	554	566	572	613
Cigarettes	506	521	512	493	525	547	545
Smokeless Tobacco	252	247	226	232	259	237	285
Cigars	311	302	280	309	305	289	324
Pipe Tobacco	132	124	135	136	156	164	128
ALCOHOL	630	577	610	598	584	641	615
Binge Alcohol Use ²	527	492	492	507	541	562	564
Heavy Alcohol Use ²	323	306	307	302	333	356	339
FEMALE							
TOBACCO PRODUCTS¹	488	477	469	515	477	486	542
Cigarettes	475	460	466	508	471	469	538
Smokeless Tobacco	99	125	63	85	74	111	73
Cigars	99	117	107	98	117	104	100
Pipe Tobacco	89	32	39	37	44	26	50
ALCOHOL	628	582	654	615	614	657	653
Binge Alcohol Use ²	393	347	361	370	356	364	380
Heavy Alcohol Use ²	158	155	160	146	147	157	182

*Low precision; no estimate reported.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, 2007, and 2008.

Table 8.25D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Standard Errors of Percentages, 2002-2008

Gender/Substance	2002	2003	2004	2005	2006	2007	2008
TOTAL							
TOBACCO PRODUCTS¹	0.44	0.41	0.41	0.43	0.42	0.42	0.44
Cigarettes	0.42	0.41	0.39	0.39	0.40	0.40	0.41
Smokeless Tobacco	0.15	0.15	0.13	0.14	0.14	0.14	0.16
Cigars	0.18	0.18	0.17	0.18	0.18	0.16	0.18
Pipe Tobacco	0.09	0.07	0.08	0.08	0.09	0.09	0.07
ALCOHOL	0.53	0.49	0.51	0.51	0.49	0.52	0.50
Binge Alcohol Use ²	0.39	0.35	0.36	0.35	0.37	0.37	0.36
Heavy Alcohol Use ²	0.20	0.19	0.19	0.19	0.21	0.21	0.20
MALE							
TOBACCO PRODUCTS¹	0.65	0.63	0.63	0.62	0.63	0.63	0.67
Cigarettes	0.59	0.60	0.59	0.56	0.58	0.60	0.59
Smokeless Tobacco	0.29	0.29	0.26	0.26	0.29	0.26	0.31
Cigars	0.36	0.35	0.32	0.35	0.34	0.32	0.35
Pipe Tobacco	0.15	0.14	0.15	0.15	0.17	0.18	0.14
ALCOHOL	0.74	0.67	0.70	0.67	0.65	0.70	0.67
Binge Alcohol Use ²	0.62	0.57	0.56	0.57	0.60	0.62	0.61
Heavy Alcohol Use ²	0.38	0.35	0.35	0.34	0.37	0.39	0.37
FEMALE							
TOBACCO PRODUCTS¹	0.52	0.50	0.49	0.53	0.49	0.49	0.54
Cigarettes	0.51	0.49	0.49	0.53	0.48	0.47	0.54
Smokeless Tobacco	0.11	0.13	0.07	0.09	0.08	0.11	0.07
Cigars	0.11	0.12	0.11	0.10	0.12	0.11	0.10
Pipe Tobacco	0.10	0.03	0.04	0.04	0.05	0.03	0.05
ALCOHOL	0.67	0.61	0.68	0.64	0.63	0.66	0.65
Binge Alcohol Use ²	0.42	0.37	0.38	0.38	0.36	0.37	0.38
Heavy Alcohol Use ²	0.17	0.16	0.17	0.15	0.15	0.16	0.18

*Low precision; no estimate reported.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, 2007, and 2008.

Table 8.25P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: *P* Values from Tests of Differences of Percentages, 2008 Versus 2002, 2003, 2004, 2005, 2006, and 2007

Gender/Substance	2002	2003	2004	2005	2006	2007	2008
TOTAL							
TOBACCO PRODUCTS¹	0.0109	0.1257	0.7282	0.2457	0.0944	0.7742	
Cigarettes	0.0164	0.1527	0.6154	0.3550	0.1567	0.6344	
Smokeless Tobacco	0.7273	0.5138	0.0044	0.0983	0.4746	0.1533	
Cigars	0.5044	0.7917	0.4399	0.2789	0.4058	0.9168	
Pipe Tobacco	0.3731	0.9704	0.7054	0.0728	0.0183	0.2705	
ALCOHOL	0.2262	0.0013	0.0115	0.7164	0.0981	0.3287	
Binge Alcohol Use ²	0.2255	0.0267	0.0576	0.0393	0.1586	0.7828	
Heavy Alcohol Use ²	0.2059	0.2026	0.5424	0.0238	0.3362	0.5887	
MALE							
TOBACCO PRODUCTS¹	0.0128	0.2699	0.4316	0.2701	0.0441	0.5337	
Cigarettes	0.0206	0.1903	0.3150	0.4514	0.0860	0.3541	
Smokeless Tobacco	0.5032	0.2136	0.0030	0.0517	0.5612	0.1241	
Cigars	0.3194	0.8382	0.4141	0.2439	0.8300	0.7141	
Pipe Tobacco	0.3759	0.7147	0.4681	0.0421	0.0091	0.1176	
ALCOHOL	0.5274	0.2865	0.2052	0.8253	0.1553	0.0851	
Binge Alcohol Use ²	0.2559	0.0592	0.1278	0.0118	0.2559	0.7604	
Heavy Alcohol Use ²	0.8288	0.2465	0.5127	0.1122	0.7614	0.9257	
FEMALE							
TOBACCO PRODUCTS¹	0.1783	0.2240	0.7641	0.5533	0.6933	0.8033	
Cigarettes	0.2007	0.4186	0.7762	0.5691	0.7151	0.7884	
Smokeless Tobacco	0.2427	0.1231	0.9501	0.5199	0.5751	0.9836	
Cigars	0.5375	0.1119	0.8163	0.9156	0.0351	0.4180	
Pipe Tobacco	0.7492	0.2463	0.3686	0.7372	0.7678	0.1232	
ALCOHOL	0.2712	0.0002	0.0158	0.7179	0.2742	0.7985	
Binge Alcohol Use ²	0.6728	0.2318	0.2329	0.8802	0.3620	0.9778	
Heavy Alcohol Use ²	0.0290	0.6559	1.0000	0.0539	0.0996	0.2874	

*Low precision; no estimate reported.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, 2007, and 2008.

Table 8.26C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2008

Gender/Substance	2002	2003	2004	2005	2006	2007	2008
TOTAL							
TOBACCO PRODUCTS¹	829	784	793	839	837	838	868
Cigarettes	790	774	765	772	796	802	804
Smokeless Tobacco	276	283	243	264	281	278	309
Cigars	346	333	323	351	347	331	362
Pipe Tobacco	163	130	145	145	164	169	147
ALCOHOL	972	909	950	969	955	1,010	974
Binge Alcohol Use ²	730	675	699	701	741	763	734
Heavy Alcohol Use ²	389	376	383	378	425	426	412
MALE							
TOBACCO PRODUCTS¹	565	561	568	576	580	592	629
Cigarettes	521	537	532	514	540	570	563
Smokeless Tobacco	260	255	233	239	271	248	293
Cigars	322	306	291	324	316	306	331
Pipe Tobacco	136	127	138	141	159	168	132
ALCOHOL	643	590	616	616	600	658	626
Binge Alcohol Use ²	540	517	511	529	561	585	587
Heavy Alcohol Use ²	346	320	318	322	353	374	357
FEMALE							
TOBACCO PRODUCTS¹	509	498	494	541	505	505	551
Cigarettes	497	481	488	537	497	487	546
Smokeless Tobacco	99	126	64	87	75	112	74
Cigars	110	121	119	109	129	112	112
Pipe Tobacco	90	35	41	40	46	28	53
ALCOHOL	640	593	662	635	640	671	664
Binge Alcohol Use ²	411	375	389	401	395	391	397
Heavy Alcohol Use ²	175	170	173	162	168	169	192

*Low precision; no estimate reported.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, 2007, and 2008.

Table 8.26D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Standard Errors of Percentages, 2002-2008

Gender/Substance	2002	2003	2004	2005	2006	2007	2008
TOTAL							
TOBACCO PRODUCTS¹	0.39	0.37	0.37	0.39	0.38	0.38	0.39
Cigarettes	0.38	0.36	0.36	0.35	0.36	0.36	0.36
Smokeless Tobacco	0.13	0.13	0.11	0.12	0.13	0.13	0.14
Cigars	0.16	0.16	0.15	0.16	0.16	0.15	0.16
Pipe Tobacco	0.08	0.06	0.07	0.07	0.07	0.08	0.07
ALCOHOL	0.46	0.43	0.44	0.44	0.43	0.45	0.43
Binge Alcohol Use ²	0.35	0.32	0.32	0.32	0.34	0.34	0.33
Heavy Alcohol Use ²	0.19	0.18	0.18	0.17	0.19	0.19	0.18
MALE							
TOBACCO PRODUCTS¹	0.56	0.55	0.55	0.55	0.55	0.55	0.58
Cigarettes	0.52	0.53	0.51	0.49	0.51	0.53	0.52
Smokeless Tobacco	0.26	0.25	0.22	0.23	0.25	0.23	0.27
Cigars	0.32	0.30	0.28	0.31	0.30	0.28	0.30
Pipe Tobacco	0.13	0.12	0.13	0.13	0.15	0.16	0.12
ALCOHOL	0.64	0.58	0.59	0.59	0.56	0.61	0.58
Binge Alcohol Use ²	0.53	0.51	0.49	0.50	0.53	0.54	0.54
Heavy Alcohol Use ²	0.34	0.31	0.31	0.31	0.33	0.35	0.33
FEMALE							
TOBACCO PRODUCTS¹	0.47	0.45	0.44	0.48	0.44	0.44	0.47
Cigarettes	0.45	0.44	0.44	0.48	0.43	0.42	0.47
Smokeless Tobacco	0.09	0.11	0.06	0.08	0.07	0.10	0.06
Cigars	0.10	0.11	0.11	0.10	0.11	0.10	0.10
Pipe Tobacco	0.08	0.03	0.04	0.04	0.04	0.02	0.05
ALCOHOL	0.59	0.54	0.59	0.56	0.56	0.58	0.57
Binge Alcohol Use ²	0.38	0.34	0.35	0.36	0.35	0.34	0.34
Heavy Alcohol Use ²	0.16	0.15	0.16	0.14	0.15	0.15	0.16

*Low precision; no estimate reported.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, 2007, and 2008.

Table 8.26P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: *P* Values from Tests of Differences of Percentages, 2008 Versus 2002, 2003, 2004, 2005, 2006, and 2007

Gender/Substance	2002	2003	2004	2005	2006	2007	2008
TOTAL							
TOBACCO PRODUCTS¹	0.0004	0.0130	0.1894	0.0459	0.0194	0.6726	
Cigarettes	0.0002	0.0058	0.0906	0.0519	0.0290	0.5398	
Smokeless Tobacco	0.4545	0.2510	0.0019	0.0760	0.4312	0.1417	
Cigars	0.6223	0.6717	0.0680	0.1153	0.1620	0.7825	
Pipe Tobacco	0.6781	0.4310	0.8839	0.0613	0.0274	0.4182	
ALCOHOL	0.1806	0.0022	0.0091	0.7519	0.1692	0.3476	
Binge Alcohol Use ²	0.2518	0.0704	0.1102	0.1155	0.3876	1.0000	
Heavy Alcohol Use ²	0.3765	0.4662	0.9071	0.1300	0.8160	0.7287	
MALE							
TOBACCO PRODUCTS¹	0.0021	0.0937	0.1642	0.0949	0.0158	0.4038	
Cigarettes	0.0010	0.0209	0.0593	0.1305	0.0272	0.2671	
Smokeless Tobacco	0.3174	0.0821	0.0015	0.0385	0.5058	0.1073	
Cigars	0.4048	0.9215	0.0751	0.1129	0.4417	0.9113	
Pipe Tobacco	0.5589	0.8752	0.4421	0.0272	0.0081	0.1531	
ALCOHOL	0.6468	0.5846	0.2584	0.5532	0.2818	0.1290	
Binge Alcohol Use ²	0.4774	0.2632	0.3111	0.1007	0.5177	0.9954	
Heavy Alcohol Use ²	0.8137	0.5601	0.9413	0.4229	0.8891	0.9508	
FEMALE							
TOBACCO PRODUCTS¹	0.0215	0.0366	0.5819	0.1995	0.2973	0.7703	
Cigarettes	0.0163	0.0715	0.5516	0.1963	0.2937	0.7866	
Smokeless Tobacco	0.3319	0.1325	0.9210	0.5104	0.5652	0.9110	
Cigars	0.5667	0.0833	0.4440	0.6405	0.0313	0.5714	
Pipe Tobacco	0.9014	0.0558	0.0887	0.5511	0.3679	0.0351	
ALCOHOL	0.1441	0.0001	0.0074	0.9519	0.3027	0.9175	
Binge Alcohol Use ²	0.3699	0.1195	0.1623	0.5859	0.5237	0.9916	
Heavy Alcohol Use ²	0.0284	0.7178	0.9822	0.0648	0.4446	0.5029	

*Low precision; no estimate reported.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, 2005, 2006, 2007, and 2008.