

Table 6.8A (REVISED) Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness: Numbers in Thousands, 2008

Drug	Total¹	Serious Mental Illness	No Serious Mental Illness
ILLCIT DRUGS²	30,892	2,289	28,596
Marijuana and Hashish	22,633	1,625	21,002
Cocaine	4,977	488	4,489
Crack	1,079	160	919
Heroin	415	62	353
Hallucinogens	2,982	278	2,704
LSD	623	55	568
PCP	37	3	34
Ecstasy	1,770	180	1,589
Inhalants	1,079	69	1,010
Nonmedical Use of Psychotherapeutics ^{3,4}	13,271	1,359	11,911
Pain Relievers	10,304	1,041	9,262
OxyContin®	1,236	126	1,110
Tranquilizers	4,616	591	4,024
Stimulants ⁴	2,275	332	1,943
Methamphetamine ⁴	760	125	635
Sedatives	513	128	385
ILLCIT DRUGS OTHER THAN MARIJUANA²	17,159	1,630	15,529

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.8B (REVISED) Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness: Percentages, 2008

Drug	Total ¹	Serious Mental Illness	No Serious Mental Illness
ILLCIT DRUGS²	13.7	27.5	13.2
Marijuana and Hashish	10.1	19.5	9.7
Cocaine	2.2	5.9	2.1
Crack	0.5	1.9	0.4
Heroin	0.2	0.7	0.2
Hallucinogens	1.3	3.3	1.2
LSD	0.3	0.7	0.3
PCP	0.0	0.0	0.0
Ecstasy	0.8	2.2	0.7
Inhalants	0.5	0.8	0.5
Nonmedical Use of Psychotherapeutics ^{3,4}	5.9	16.3	5.5
Pain Relievers	4.6	12.5	4.3
OxyContin®	0.5	1.5	0.5
Tranquilizers	2.1	7.1	1.9
Stimulants ⁴	1.0	4.0	0.9
Methamphetamine ⁴	0.3	1.5	0.3
Sedatives	0.2	1.5	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA²	7.6	19.6	7.2

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.9A (REVISED) Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Mental Illness: Numbers in Thousands, 2008

Substance	Total ¹	Serious Mental Illness	No Serious Mental Illness
PAST YEAR			
Tobacco Products ²	79,803	4,650	75,140
Cigarettes	66,368	4,296	62,059
Smokeless Tobacco	10,607	503	10,104
Cigars	24,155	1,177	22,978
Alcohol	157,239	6,095	151,097
PAST MONTH			
Tobacco Products ²	68,085	4,173	63,900
Cigarettes	57,551	3,894	53,646
Smokeless Tobacco	8,150	298	7,852
Cigars	12,274	616	11,658
Pipe Tobacco	1,707	123	1,585
Alcohol	125,213	4,553	120,616
Binge Alcohol Use ³	56,159	2,274	53,868
Heavy Alcohol Use ³	16,877	933	15,928

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.9B (REVISED) Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Mental Illness: Percentages, 2008

Substance	Total ¹	Serious Mental Illness	No Serious Mental Illness
PAST YEAR			
Tobacco Products ²	35.5	55.8	34.7
Cigarettes	29.5	51.6	28.7
Smokeless Tobacco	4.7	6.0	4.7
Cigars	10.7	14.1	10.6
Alcohol	69.9	73.2	69.8
PAST MONTH			
Tobacco Products ²	30.3	50.1	29.5
Cigarettes	25.6	46.7	24.8
Smokeless Tobacco	3.6	3.6	3.6
Cigars	5.5	7.4	5.4
Pipe Tobacco	0.8	1.5	0.7
Alcohol	55.7	54.6	55.7
Binge Alcohol Use ³	25.0	27.3	24.9
Heavy Alcohol Use ³	7.5	11.2	7.4

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ Estimates in the Total column represent persons aged 18 or older, including those with unknown past year SMI information.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.