

**Table 7.47C Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008**

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	105	93	17	15	26	26	100	89
From More Than One Doctor	33	36	6	5	8	9	31	35
Wrote Fake Prescription	6	11	1	1	3	3	5	10
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	11	11	3	3	3	7	10	8
From Friend or Relative for Free	156	159	24	25	56	58	142	142
Bought from Friend or Relative	58	50	10	11	25	25	52	42
Took from Friend or Relative without Asking	41	41	11	12	16	16	36	36
Bought from Drug Dealer or Other Stranger	40	42	8	8	15	15	36	38
Bought on the Internet	11	14	1	1	3	3	11	14
Some Other Way	39	43	10	9	15	13	34	38

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.47D Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	0.77	0.76	1.04	1.02	0.63	0.65	1.31	1.31
From More Than One Doctor	0.28	0.32	0.41	0.40	0.22	0.23	0.48	0.55
Wrote Fake Prescription	0.05	0.10	0.07	0.08	0.08	0.08	0.08	0.17
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.09	0.10	0.20	0.23	0.08	0.18	0.16	0.13
From Friend or Relative for Free	0.90	0.94	1.22	1.28	0.88	0.95	1.52	1.56
Bought from Friend or Relative	0.49	0.44	0.68	0.75	0.61	0.64	0.80	0.68
Took from Friend or Relative without Asking	0.35	0.36	0.68	0.81	0.42	0.40	0.56	0.58
Bought from Drug Dealer or Other Stranger	0.35	0.37	0.54	0.54	0.39	0.41	0.57	0.62
Bought on the Internet	0.10	0.13	0.08	0.10	0.07	0.07	0.17	0.23
Some Other Way	0.33	0.37	0.66	0.61	0.40	0.36	0.53	0.61

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

**Table 7.47P Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007**

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	0.4346		0.3299		0.6723		0.4644	
From More Than One Doctor	0.2688		0.6231		0.6297		0.2033	
Wrote Fake Prescription	0.3608		0.7231		0.6951		0.3002	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.1707		0.3420		0.0055		0.9360	
From Friend or Relative for Free	0.8171		0.5921		0.5002		0.9566	
Bought from Friend or Relative	0.6420		0.4622		0.9288		0.4248	
Took from Friend or Relative without Asking	0.8174		0.0104		0.1660		0.8909	
Bought from Drug Dealer or Other Stranger	0.3025		0.8666		0.1015		0.7772	
Bought on the Internet	0.2002		0.3266		0.9753		0.2396	
Some Other Way	0.9974		0.0037		0.6846		0.5099	

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.48C Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Pain Relievers¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	136	141	17	17	42	46	129	130
From More Than One Doctor	24	22	4	4	7	8	22	20
Wrote Fake Prescription	4	2	0	0	3	2	3	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	14	14	3	2	2	3	14	13
From Friend or Relative for Free	35	30	8	8	14	13	31	26
Bought from Friend or Relative	24	21	5	5	12	11	20	16
Took from Friend or Relative without Asking	10	9	6	5	5	5	6	5
Bought from Drug Dealer or Other Stranger	16	14	4	4	9	8	13	11
Bought on the Internet	1	4	0	0	1	2	*	3
Some Other Way	13	8	5	4	4	5	11	5

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.48D Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	0.93	0.89	2.03	2.03	1.07	1.12	1.37	1.25
From More Than One Doctor	0.41	0.40	0.76	0.82	0.39	0.44	0.65	0.62
Wrote Fake Prescription	0.07	0.04	0.06	0.07	0.14	0.11	0.09	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.25	0.25	0.52	0.47	0.11	0.16	0.41	0.41
From Friend or Relative for Free	0.58	0.53	1.37	1.42	0.70	0.70	0.90	0.78
Bought from Friend or Relative	0.41	0.38	0.91	0.89	0.62	0.62	0.59	0.51
Took from Friend or Relative without Asking	0.18	0.16	1.07	0.96	0.28	0.28	0.18	0.15
Bought from Drug Dealer or Other Stranger	0.28	0.26	0.77	0.77	0.46	0.43	0.38	0.34
Bought on the Internet	0.03	0.07	0.07	0.07	0.08	0.09	*	0.10
Some Other Way	0.22	0.15	0.88	0.79	0.24	0.29	0.32	0.16

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.48P Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: *P* Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	0.3532		0.5583		0.6340		0.2690	
From More Than One Doctor	0.4491		0.4035		0.4603		0.6744	
Wrote Fake Prescription	0.1870		0.3787		0.3523		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.8210		0.5261		0.2122		0.4459	
From Friend or Relative for Free	0.1662		0.8833		0.8400		0.1235	
Bought from Friend or Relative	0.4502		0.7797		0.6817		0.5808	
Took from Friend or Relative without Asking	0.3179		0.8902		0.7718		0.1562	
Bought from Drug Dealer or Other Stranger	0.7328		0.8183		0.6260		0.4944	
Bought on the Internet	0.0961		0.1818		0.1369		*	
Some Other Way	0.1949		0.1005		0.6835		0.2513	

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.49C Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	80	79	12	11	20	19	78	75
From More Than One Doctor	35	38	5	4	9	9	34	36
Wrote Fake Prescription	10	6	1	1	3	3	9	5
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	13	10	3	3	4	4	12	8
From Friend or Relative for Free	100	100	16	15	35	34	94	91
Bought from Friend or Relative	55	56	11	9	25	24	48	50
Took from Friend or Relative without Asking	37	39	9	9	16	15	33	36
Bought from Drug Dealer or Other Stranger	43	40	7	7	16	17	40	36
Bought on the Internet	12	14	2	2	4	4	11	13
Some Other Way	30	26	8	6	11	10	27	23

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.49D Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	1.37	1.42	1.80	1.88	1.19	1.22	2.27	2.35
From More Than One Doctor	0.72	0.82	0.84	0.82	0.59	0.66	1.21	1.35
Wrote Fake Prescription	0.21	0.14	0.23	0.19	0.22	0.23	0.34	0.20
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.27	0.21	0.51	0.63	0.28	0.31	0.43	0.31
From Friend or Relative for Free	1.51	1.53	2.06	2.11	1.41	1.45	2.47	2.49
Bought from Friend or Relative	1.10	1.16	1.66	1.60	1.42	1.47	1.66	1.78
Took from Friend or Relative without Asking	0.80	0.85	1.40	1.68	1.02	0.98	1.21	1.34
Bought from Drug Dealer or Other Stranger	0.88	0.86	1.26	1.32	1.02	1.10	1.41	1.34
Bought on the Internet	0.25	0.31	0.36	0.38	0.25	0.29	0.41	0.51
Some Other Way	0.64	0.58	1.40	1.24	0.79	0.75	1.00	0.88

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.49P Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	0.3687		0.7392		0.6572		0.4387	
From More Than One Doctor	0.1815		0.8645		0.6227		0.2130	
Wrote Fake Prescription	0.3794		0.9308		0.7967		0.3186	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7965		0.0605		0.2639		0.6367	
From Friend or Relative for Free	0.2489		0.2432		0.5940		0.2230	
Bought from Friend or Relative	0.2714		0.5481		0.5342		0.1028	
Took from Friend or Relative without Asking	0.0939		0.0286		0.8153		0.1332	
Bought from Drug Dealer or Other Stranger	0.1652		0.4026		0.1640		0.4060	
Bought on the Internet	0.4733		0.4852		0.3144		0.6494	
Some Other Way	0.0914		0.0187		0.6811		0.3190	

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.50C Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Pain Relievers¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	86	84	11	10	28	27	83	79
From More Than One Doctor	30	27	5	5	10	10	27	25
Wrote Fake Prescription	15	9	1	2	3	3	14	8
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	19	16	2	3	4	4	18	15
From Friend or Relative for Free	36	34	7	6	14	13	32	31
Bought from Friend or Relative	27	27	6	5	14	14	22	23
Took from Friend or Relative without Asking	13	15	6	5	7	6	9	14
Bought from Drug Dealer or Other Stranger	18	18	5	5	11	10	14	15
Bought on the Internet	6	7	1	1	4	4	5	5
Some Other Way	10	14	4	3	5	4	8	12

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.50D Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Pain Relievers¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	1.53	1.58	3.01	2.93	1.64	1.72	2.33	2.39
From More Than One Doctor	1.16	1.10	1.92	2.19	1.14	1.21	1.84	1.72
Wrote Fake Prescription	0.59	0.37	0.55	0.79	0.35	0.44	0.99	0.56
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.74	0.66	0.95	1.33	0.43	0.55	1.24	1.07
From Friend or Relative for Free	1.35	1.39	2.50	2.67	1.52	1.56	2.10	2.13
Bought from Friend or Relative	1.05	1.14	2.20	2.24	1.52	1.62	1.50	1.67
Took from Friend or Relative without Asking	0.52	0.64	2.37	2.10	0.82	0.72	0.62	0.98
Bought from Drug Dealer or Other Stranger	0.74	0.75	1.91	2.19	1.28	1.26	0.96	1.05
Bought on the Internet	0.27	0.29	0.37	0.68	0.49	0.56	0.36	0.36
Some Other Way	0.43	0.58	1.74	1.42	0.69	0.56	0.55	0.90

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.50P Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	0.5277		0.1988		0.4728		0.3491	
From More Than One Doctor	0.5830		0.2237		0.4211		0.8610	
Wrote Fake Prescription	0.2834		0.5955		0.0234		0.1263	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.8522		0.0591		0.2004		0.4273	
From Friend or Relative for Free	0.7231		0.3845		0.6978		0.7371	
Bought from Friend or Relative	0.2622		0.8723		0.5145		0.2554	
Took from Friend or Relative without Asking	0.8829		0.2789		0.4052		0.3060	
Bought from Drug Dealer or Other Stranger	0.2982		0.0794		0.6458		0.5147	
Bought on the Internet	0.4643		0.1157		0.5571		0.9901	
Some Other Way	0.4454		0.5711		0.2741		0.1026	

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

**Table 7.51C Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008**

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	48	43	5	4	13	10	46	41
From More Than One Doctor	14	14	2	2	5	5	13	12
Wrote Fake Prescription	10	3	2	2	2	2	9	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	9	5	1	2	3	4	9	*
From Friend or Relative for Free	114	111	13	13	34	35	106	105
Bought from Friend or Relative	41	44	8	9	20	21	33	37
Took from Friend or Relative without Asking	40	45	5	6	10	10	38	43
Bought from Drug Dealer or Other Stranger	30	23	7	6	14	13	25	18
Bought on the Internet	12	43	1	1	3	2	12	43
Some Other Way	27	32	4	4	10	10	25	30

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.51D Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	0.96	0.87	1.34	1.11	0.74	0.64	1.57	1.42
From More Than One Doctor	0.28	0.28	0.49	0.54	0.29	0.33	0.45	0.44
Wrote Fake Prescription	0.20	0.06	0.48	0.45	0.14	0.13	0.33	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.20	0.09	0.32	0.42	0.21	0.26	0.31	*
From Friend or Relative for Free	1.47	1.54	2.46	2.36	1.40	1.46	2.29	2.45
Bought from Friend or Relative	0.81	0.88	1.91	1.90	1.08	1.13	1.14	1.28
Took from Friend or Relative without Asking	0.81	0.91	1.33	1.43	0.63	0.63	1.33	1.48
Bought from Drug Dealer or Other Stranger	0.62	0.48	1.77	1.55	0.82	0.77	0.87	0.63
Bought on the Internet	0.26	0.87	0.32	0.36	0.18	0.12	0.43	1.48
Some Other Way	0.55	0.64	1.06	0.89	0.59	0.63	0.87	1.03

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

**Table 7.51P Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007**

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	0.1823		0.4255		0.1014		0.4259	
From More Than One Doctor	0.6779		0.9972		0.3622		0.9757	
Wrote Fake Prescription	0.3142		0.8480		0.6598		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5517		0.6436		0.4241		*	
From Friend or Relative for Free	0.3615		0.5468		0.9747		0.2726	
Bought from Friend or Relative	0.0450		0.8703		0.5336		0.0324	
Took from Friend or Relative without Asking	0.2434		0.5041		0.6903		0.3127	
Bought from Drug Dealer or Other Stranger	0.2425		0.4495		0.6892		0.3440	
Bought on the Internet	0.1631		0.6982		0.1044		0.1349	
Some Other Way	0.9627		0.3191		0.4420		0.9532	

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.52C Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Tranquilizers¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	99	91	7	7	25	25	95	88
From More Than One Doctor	11	17	2	3	5	5	10	16
Wrote Fake Prescription	1	1	*	0	1	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	4	*	0	2	3	5	3
From Friend or Relative for Free	29	32	5	5	12	13	26	29
Bought from Friend or Relative	19	19	4	4	9	11	16	15
Took from Friend or Relative without Asking	8	9	3	4	4	3	*	*
Bought from Drug Dealer or Other Stranger	14	12	3	2	5	5	13	11
Bought on the Internet	7	3	0	0	2	3	7	0
Some Other Way	13	14	2	2	5	5	11	13

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.52D Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	1.49	1.62	3.91	3.80	1.85	1.99	1.96	2.22
From More Than One Doctor	0.44	0.67	1.34	2.26	0.72	0.64	0.56	0.96
Wrote Fake Prescription	0.03	0.04	*	0.26	0.11	0.11	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.19	0.17	*	0.29	0.23	0.43	0.28	0.17
From Friend or Relative for Free	1.06	1.18	3.43	3.06	1.46	1.53	1.43	1.66
Bought from Friend or Relative	0.71	0.73	2.72	2.49	1.13	1.31	0.90	0.88
Took from Friend or Relative without Asking	0.31	0.33	2.37	2.74	0.46	0.46	*	*
Bought from Drug Dealer or Other Stranger	0.53	0.47	2.29	1.61	0.64	0.60	0.72	0.67
Bought on the Internet	0.26	0.10	0.22	0.20	0.33	0.34	0.37	0.03
Some Other Way	0.48	0.55	1.22	1.58	0.61	0.59	0.64	0.77

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.52P Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: *P* Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	0.1153		0.5862		0.8454		0.1835	
From More Than One Doctor	0.0619		0.2302		0.2964		0.0394	
Wrote Fake Prescription	0.2509		*		0.3231		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.9887		*		0.2873		0.3381	
From Friend or Relative for Free	0.2191		0.1648		0.8398		0.1439	
Bought from Friend or Relative	0.9127		0.7235		0.1876		0.4312	
Took from Friend or Relative without Asking	0.2614		0.0256		0.4195		*	
Bought from Drug Dealer or Other Stranger	0.0686		0.0302		0.2774		0.3172	
Bought on the Internet	0.3000		0.4608		0.7198		0.3183	
Some Other Way	0.2216		0.5955		0.9288		0.2731	

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.53C Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	33	31	3	3	8	7	31	29
From More Than One Doctor	11	12	*	*	3	3	10	11
Wrote Fake Prescription	9	2	1	*	2	2	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	9	3	0	*	3	3	*	*
From Friend or Relative for Free	65	68	7	8	19	19	61	65
Bought from Friend or Relative	27	30	5	4	14	14	23	27
Took from Friend or Relative without Asking	23	24	3	4	9	8	21	22
Bought from Drug Dealer or Other Stranger	31	27	4	4	10	10	29	25
Bought on the Internet	12	24	1	1	3	2	11	*
Some Other Way	14	11	2	2	5	5	13	10

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.53D Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	1.84	1.74	2.81	2.19	1.37	1.30	2.95	2.66
From More Than One Doctor	0.65	0.70	*	*	0.53	0.60	1.06	1.10
Wrote Fake Prescription	0.58	0.12	1.18	*	0.40	0.39	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.55	0.20	0.10	*	0.50	0.58	*	*
From Friend or Relative for Free	2.46	2.37	4.55	4.31	2.39	2.48	3.89	3.68
Bought from Friend or Relative	1.60	1.74	3.85	3.12	2.06	2.26	2.27	2.53
Took from Friend or Relative without Asking	1.33	1.35	2.88	2.72	1.45	1.41	2.07	2.08
Bought from Drug Dealer or Other Stranger	1.74	1.55	3.47	2.71	1.63	1.73	2.77	2.35
Bought on the Internet	0.71	1.43	0.83	1.10	0.49	0.37	1.15	*
Some Other Way	0.84	0.69	2.06	1.56	0.93	0.87	1.29	1.01

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.53P Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	0.6197		0.4497		0.1571		0.9320	
From More Than One Doctor	0.3966		*		0.8200		0.4613	
Wrote Fake Prescription	0.2055		*		0.4081		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.4985		*		0.2572		*	
From Friend or Relative for Free	0.3027		0.6048		0.4556		0.4948	
Bought from Friend or Relative	0.3232		0.0737		0.9625		0.0544	
Took from Friend or Relative without Asking	0.7158		0.9588		0.8880		0.5757	
Bought from Drug Dealer or Other Stranger	0.9963		0.2728		0.8347		0.8837	
Bought on the Internet	0.3789		0.2543		0.1769		*	
Some Other Way	0.1677		0.7573		0.6621		0.2175	

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.54C Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	53	55	*	*	15	14	50	54
From More Than One Doctor	21	24	*	*	4	5	*	23
Wrote Fake Prescription	*	0	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	1	*	*	2	*	*	*
From Friend or Relative for Free	14	17	*	*	8	10	12	13
Bought from Friend or Relative	12	11	*	*	5	7	11	8
Took from Friend or Relative without Asking	5	7	*	*	3	3	4	6
Bought from Drug Dealer or Other Stranger	10	13	*	*	4	4	9	13
Bought on the Internet	7	1	*	*	2	1	*	0
Some Other Way	3	10	*	*	1	3	2	10

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.54D Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	2.85	2.77	*	*	2.95	3.44	4.01	3.76
From More Than One Doctor	2.28	2.35	*	*	1.53	1.80	*	3.41
Wrote Fake Prescription	*	0.02	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.22	0.15	*	*	0.72	*	*	*
From Friend or Relative for Free	1.64	1.77	*	*	2.71	3.24	2.05	2.04
Bought from Friend or Relative	1.45	1.18	*	*	1.99	2.57	1.87	1.27
Took from Friend or Relative without Asking	0.64	0.73	*	*	1.03	1.03	0.67	0.88
Bought from Drug Dealer or Other Stranger	1.10	1.34	*	*	1.68	1.55	1.45	1.91
Bought on the Internet	0.78	0.08	*	*	0.66	0.30	*	0.01
Some Other Way	0.37	1.10	*	*	0.46	1.05	0.44	1.57

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.54P Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	0.9912		*		0.0858		0.5549	
From More Than One Doctor	0.1109		*		0.1741		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.1945		*		*		*	
From Friend or Relative for Free	0.8181		*		0.4379		0.8678	
Bought from Friend or Relative	0.8371		*		0.3946		0.3098	
Took from Friend or Relative without Asking	0.9652		*		0.5749		0.6693	
Bought from Drug Dealer or Other Stranger	0.9775		*		0.4063		0.5355	
Bought on the Internet	0.1947		*		0.3395		*	
Some Other Way	0.1172		*		0.1288		0.1745	

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

**Table 7.55C Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008**

Source for Most Recent Nonmedical Use among Past Year Stimulant Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	28	27	7	4	10	8	25	25
From More Than One Doctor	5	6	2	2	2	4	4	*
Wrote Fake Prescription	7	3	1	2	2	2	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	6	6	2	2	3	4	4	4
From Friend or Relative for Free	70	58	11	11	28	25	62	51
Bought from Friend or Relative	22	28	6	6	16	18	14	20
Took from Friend or Relative without Asking	18	16	5	5	8	7	16	14
Bought from Drug Dealer or Other Stranger	16	16	5	4	8	9	12	13
Bought on the Internet	17	15	2	1	3	4	17	14
Some Other Way	19	11	4	3	4	5	19	9

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

**Table 7.55D Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008**

Source for Most Recent Nonmedical Use among Past Year Stimulant Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	1.24	1.32	1.92	1.53	1.07	0.94	2.52	2.78
From More Than One Doctor	0.24	0.32	0.53	0.60	0.28	0.44	0.45	*
Wrote Fake Prescription	0.34	0.13	0.40	0.65	0.19	0.22	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.26	0.30	0.71	0.72	0.36	0.44	0.43	0.48
From Friend or Relative for Free	1.97	1.92	2.55	2.83	1.83	1.90	3.99	3.87
Bought from Friend or Relative	1.03	1.30	1.76	1.98	1.49	1.75	1.57	2.30
Took from Friend or Relative without Asking	0.84	0.79	1.54	1.60	0.85	0.77	1.67	1.60
Bought from Drug Dealer or Other Stranger	0.73	0.80	1.40	1.51	0.88	1.10	1.31	1.47
Bought on the Internet	0.79	0.77	0.48	0.19	0.34	0.50	1.77	1.64
Some Other Way	0.88	0.55	1.18	1.20	0.42	0.59	2.00	1.06

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

**Table 7.55P Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007**

Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	0.9772		0.0814		0.5702		0.4152	
From More Than One Doctor	0.7011		0.6146		0.2717		*	
Wrote Fake Prescription	0.5179		0.4986		0.1408		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.8148		0.4665		0.3807		0.3565	
From Friend or Relative for Free	0.1980		0.3994		0.0478		0.4787	
Bought from Friend or Relative	0.0006		0.5532		0.1431		0.0012	
Took from Friend or Relative without Asking	0.6201		0.9238		0.2595		0.9700	
Bought from Drug Dealer or Other Stranger	0.0791		0.4581		0.1185		0.4189	
Bought on the Internet	0.1226		0.2069		0.4993		0.0952	
Some Other Way	0.1199		0.5716		0.8619		0.1297	

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.56C Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Stimulants ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	49	43	8	9	23	22	*	*
From More Than One Doctor	6	7	3	3	4	3	3	5
Wrote Fake Prescription	0	0	0	0	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	1	*	0	1	1	*	*
From Friend or Relative for Free	22	23	3	3	8	7	*	*
Bought from Friend or Relative	28	11	3	4	5	5	*	9
Took from Friend or Relative without Asking	3	3	3	3	2	1	*	*
Bought from Drug Dealer or Other Stranger	3	3	2	2	3	2	1	*
Bought on the Internet	5	4	1	*	3	2	5	*
Some Other Way	3	4	1	0	0	1	3	3

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.56D Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Stimulants ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	2.89	2.61	4.34	4.72	2.21	2.40	*	*
From More Than One Doctor	0.60	0.76	2.60	2.59	0.89	0.86	0.69	1.26
Wrote Fake Prescription	0.01	0.01	0.06	0.07	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.15	0.11	*	0.43	0.19	0.21	*	*
From Friend or Relative for Free	2.02	2.35	2.57	2.68	1.62	1.75	*	*
Bought from Friend or Relative	2.49	1.19	2.30	3.11	1.14	1.37	*	2.18
Took from Friend or Relative without Asking	0.33	0.35	2.37	2.42	0.37	0.35	*	*
Bought from Drug Dealer or Other Stranger	0.33	0.30	1.51	1.62	0.57	0.48	0.25	*
Bought on the Internet	0.53	0.46	0.78	*	0.57	0.48	1.01	*
Some Other Way	0.33	0.41	0.45	0.32	0.09	0.34	0.72	0.87

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.56P Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: *P* Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007

Source Where Friend or Relative Obtained Stimulants ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	0.1245		0.9496		0.8314		*	
From More Than One Doctor	0.6624		0.4380		0.7552		0.2624	
Wrote Fake Prescription	0.3630		0.4803		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7509		*		0.3671		*	
From Friend or Relative for Free	0.7984		0.3012		0.7443		*	
Bought from Friend or Relative	0.1455		0.0308		0.5908		*	
Took from Friend or Relative without Asking	0.6683		0.7071		0.9364		*	
Bought from Drug Dealer or Other Stranger	0.4961		0.9701		0.5911		*	
Bought on the Internet	0.0728		*		0.1538		*	
Some Other Way	0.1240		0.4025		0.2645		0.1629	

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.57C Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

Source for Past Month Nonmedical Use among Past Month Stimulant Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	18	17	5	3	7	6	*	*
From More Than One Doctor	6	7	2	1	4	5	*	*
Wrote Fake Prescription	2	2	1	2	*	2	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	5	2	1	2	3	*	*
From Friend or Relative for Free	38	37	5	5	15	12	*	*
Bought from Friend or Relative	15	20	3	4	12	12	9	15
Took from Friend or Relative without Asking	16	13	3	3	7	6	*	*
Bought from Drug Dealer or Other Stranger	11	12	3	3	7	7	*	9
Bought on the Internet	13	13	*	*	5	4	*	*
Some Other Way	13	4	2	2	2	2	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.57D Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source for Past Month Nonmedical Use among Past Month Stimulant Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	2.34	2.42	3.92	3.38	2.20	2.30	*	*
From More Than One Doctor	0.78	1.05	1.86	1.69	1.51	1.84	*	*
Wrote Fake Prescription	0.25	0.36	0.97	1.98	*	0.62	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.67	0.74	1.68	1.16	0.71	1.04	*	*
From Friend or Relative for Free	3.50	3.56	4.36	4.47	3.30	3.41	*	*
Bought from Friend or Relative	2.16	2.79	2.97	3.76	3.40	3.61	2.61	4.63
Took from Friend or Relative without Asking	2.13	1.79	2.54	3.38	2.40	2.01	*	*
Bought from Drug Dealer or Other Stranger	1.49	1.74	2.64	3.02	2.41	2.53	*	2.77
Bought on the Internet	1.78	1.90	*	*	1.60	1.63	*	*
Some Other Way	1.85	0.60	1.90	2.38	0.79	0.85	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.57P Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007

Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	0.4282		0.1256		0.6180		*	
From More Than One Doctor	0.5355		0.8194		0.3611		*	
Wrote Fake Prescription	0.2033		0.3379		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7405		0.5994		0.4868		*	
From Friend or Relative for Free	0.8218		0.9264		0.0270		*	
Bought from Friend or Relative	0.0086		0.0088		0.9160		0.0077	
Took from Friend or Relative without Asking	0.2873		0.0633		0.1978		*	
Bought from Drug Dealer or Other Stranger	0.1252		0.3842		0.3622		*	
Bought on the Internet	0.3999		*		0.9413		*	
Some Other Way	0.1342		0.2002		0.9796		*	

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.58C Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Stimulants ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	*	*	*	*	13	11	*	*
From More Than One Doctor	3	4	*	*	2	2	*	*
Wrote Fake Prescription	2	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	6	5	*	*
Bought from Friend or Relative	*	*	*	*	4	3	*	*
Took from Friend or Relative without Asking	2	5	*	*	2	2	*	*
Bought from Drug Dealer or Other Stranger	8	7	*	*	4	2	*	*
Bought on the Internet	5	4	*	*	*	*	*	*
Some Other Way	3	5	*	*	0	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.58D Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Stimulants ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	*	*	*	*	3.74	4.03	*	*
From More Than One Doctor	0.96	1.15	*	*	1.56	1.90	*	*
Wrote Fake Prescription	0.54	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.30	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	3.75	3.66	*	*
Bought from Friend or Relative	*	*	*	*	2.88	2.27	*	*
Took from Friend or Relative without Asking	0.72	1.49	*	*	1.54	1.51	*	*
Bought from Drug Dealer or Other Stranger	2.37	2.12	*	*	2.29	1.71	*	*
Bought on the Internet	1.39	1.38	*	*	*	*	*	*
Some Other Way	1.01	1.66	*	*	0.27	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.58P Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007

Source Where Friend or Relative Obtained Stimulants ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	*		*		0.7305		*	
From More Than One Doctor	0.6700		*		0.4443		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	*		*		0.4442		*	
Bought from Friend or Relative	*		*		0.0785		*	
Took from Friend or Relative without Asking	0.5021		*		0.6626		*	
Bought from Drug Dealer or Other Stranger	0.5831		*		0.0469		*	
Bought on the Internet	0.4623		*		*		*	
Some Other Way	0.4643		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.59C Source Where Methamphetamine was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From Friend or Relative for Free	53	35	8	6	20	14	50	31
Bought from Friend or Relative	40	37	5	4	11	9	38	35
Took from Friend or Relative without Asking	7	6	2	2	4	3	6	4
Bought from Drug Dealer or Other Stranger	38	32	5	3	11	9	36	30
Bought on the Internet	9	4	*	*	1	0	*	*
Some Other Way	9	9	1	1	4	3	*	9

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.59D Source Where Methamphetamine was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From Friend or Relative for Free	2.55	2.88	4.53	5.51	3.16	3.05	3.90	4.37
Bought from Friend or Relative	2.38	2.90	3.76	4.11	2.38	2.52	3.68	4.56
Took from Friend or Relative without Asking	0.52	0.57	1.94	2.78	0.86	1.09	0.68	0.64
Bought from Drug Dealer or Other Stranger	2.29	2.88	3.75	3.38	2.30	2.64	3.48	4.46
Bought on the Internet	0.65	0.43	*	*	0.12	0.07	*	*
Some Other Way	0.62	0.96	1.15	1.04	0.86	1.13	*	1.47

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.59P Source Where Methamphetamine was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From Friend or Relative for Free	0.0111		0.7904		1.0000		0.0049	
Bought from Friend or Relative	0.0452		0.6737		0.9186		0.0262	
Took from Friend or Relative without Asking	0.9146		0.7776		0.7192		0.7467	
Bought from Drug Dealer or Other Stranger	0.6489		0.1689		0.7328		0.4170	
Bought on the Internet	0.7223		*		0.6721		*	
Some Other Way	0.0619		0.5402		0.3551		*	

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.60C Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From Friend or Relative for Free	20	17	4	*	9	8	17	*
Bought from Friend or Relative	33	20	*	*	10	7	30	*
Took from Friend or Relative without Asking	4	2	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	29	17	*	*	11	7	*	*
Bought on the Internet	0	*	0	*	*	*	*	*
Some Other Way	5	5	*	*	3	3	4	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Methamphetamine include (a) past year users who reported obtaining this drug for their most recent use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's most recent use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.60D Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From Friend or Relative for Free	2.92	4.32	5.48	*	3.65	4.85	4.44	*
Bought from Friend or Relative	3.83	4.26	*	*	4.41	4.35	6.02	*
Took from Friend or Relative without Asking	0.72	0.70	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	4.01	4.27	*	*	4.49	4.56	*	*
Bought on the Internet	0.02	*	0.22	*	*	*	*	*
Some Other Way	0.89	1.33	*	*	1.74	2.38	1.13	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Methamphetamine include (a) past year users who reported obtaining this drug for their most recent use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's most recent use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.60P Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: *P* Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From Friend or Relative for Free	0.0099		*		0.1032		*	
Bought from Friend or Relative	0.1861		*		0.5434		*	
Took from Friend or Relative without Asking	0.8412		*		*		*	
Bought from Drug Dealer or Other Stranger	0.3048		*		0.2381		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.5109		*		0.6249		*	

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Methamphetamine include (a) past year users who reported obtaining this drug for their most recent use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's most recent use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.61C Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From Friend or Relative for Free	39	*	*	*	10	*	*	*
Bought from Friend or Relative	36	*	*	*	8	5	*	*
Took from Friend or Relative without Asking	12	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	33	*	*	*	8	*	*	*
Bought on the Internet	1	0	*	*	1	*	*	*
Some Other Way	6	7	*	*	3	2	4	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.61D Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From Friend or Relative for Free	5.10	*	*	*	6.00	*	*	*
Bought from Friend or Relative	4.91	*	*	*	5.33	5.26	*	*
Took from Friend or Relative without Asking	2.15	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	4.79	*	*	*	4.88	*	*	*
Bought on the Internet	0.13	0.03	*	*	0.41	*	*	*
Some Other Way	1.07	1.99	*	*	2.40	2.12	1.03	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.61P Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		0.2605		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	0.2042		*		*		*	
Some Other Way	0.1284		*		0.9566		*	

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.62C Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	3	3	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	3	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.62D Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	1.16	1.74	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	1.15	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.62P Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	0.5880		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

**Table 7.63C Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008**

Source for Most Recent Nonmedical Use among Past Year Sedative Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	32	21	4	3	5	4	32	*
From More Than One Doctor	4	9	2	1	2	*	4	9
Wrote Fake Prescription	5	5	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	4	*	1	1	2	*	*
From Friend or Relative for Free	55	49	5	6	10	12	53	*
Bought from Friend or Relative	21	9	3	3	5	5	*	7
Took from Friend or Relative without Asking	6	6	2	3	4	3	4	3
Bought from Drug Dealer or Other Stranger	9	10	2	3	3	2	8	9
Bought on the Internet	3	2	1	*	*	0	*	*
Some Other Way	7	8	2	2	2	2	7	8

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

**Table 7.63D Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008**

Source for Most Recent Nonmedical Use among Past Year Sedative Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	3.66	3.04	4.37	3.05	2.75	2.21	5.10	*
From More Than One Doctor	0.54	1.27	1.98	1.73	1.22	*	0.62	1.94
Wrote Fake Prescription	0.59	0.73	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.45	0.63	*	0.62	0.39	1.41	*	*
From Friend or Relative for Free	4.26	3.98	5.06	5.09	4.23	4.01	5.87	*
Bought from Friend or Relative	2.42	1.26	3.05	2.91	3.00	2.70	*	1.52
Took from Friend or Relative without Asking	0.74	0.83	2.11	3.60	2.39	1.98	0.69	0.70
Bought from Drug Dealer or Other Stranger	1.08	1.47	2.04	2.83	1.62	1.34	1.43	2.12
Bought on the Internet	0.38	0.34	0.86	*	*	0.06	*	*
Some Other Way	0.87	1.22	2.77	1.91	0.95	1.08	1.13	1.84

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

**Table 7.63P Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007**

Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	0.0386		0.0109		0.4956		*	
From More Than One Doctor	0.2337		0.4192		*		0.1130	
Wrote Fake Prescription	0.3824		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.1302		*		0.2980		*	
From Friend or Relative for Free	0.2918		0.1993		0.0629		*	
Bought from Friend or Relative	0.4158		0.9888		0.3721		*	
Took from Friend or Relative without Asking	0.4141		0.0764		0.1366		0.7664	
Bought from Drug Dealer or Other Stranger	0.1580		0.4661		0.2225		0.1288	
Bought on the Internet	0.3739		*		*		*	
Some Other Way	0.5557		0.2030		0.8641		0.4053	

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.64C Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Sedatives ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	48	43	*	*	8	9	*	*
From More Than One Doctor	4	4	*	*	1	1	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0	3	*	*	*	*	*	*
From Friend or Relative for Free	3	6	*	*	2	3	*	*
Bought from Friend or Relative	2	2	*	*	2	*	*	*
Took from Friend or Relative without Asking	1	0	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	1	1	*	*	*	*	*	*
Bought on the Internet	0	0	*	*	0	0	*	*
Some Other Way	*	2	*	*	1	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.64D Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Sedatives ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	2.50	2.46	*	*	4.05	4.40	*	*
From More Than One Doctor	1.06	1.05	*	*	1.35	1.65	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.01	0.80	*	*	*	*	*	*
From Friend or Relative for Free	0.81	1.70	*	*	3.06	3.14	*	*
Bought from Friend or Relative	0.61	0.63	*	*	2.10	*	*	*
Took from Friend or Relative without Asking	0.30	0.13	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	0.31	0.31	*	*	*	*	*	*
Bought on the Internet	0.04	0.03	*	*	0.20	0.14	*	*
Some Other Way	*	0.49	*	*	1.23	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.64P Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: *P* Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007

Source Where Friend or Relative Obtained Sedatives ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	0.5383		*		0.9756		*	
From More Than One Doctor	0.8643		*		0.9223		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.1263		*		*		*	
From Friend or Relative for Free	0.0871		*		0.9481		*	
Bought from Friend or Relative	0.6328		*		*		*	
Took from Friend or Relative without Asking	0.3594		*		*		*	
Bought from Drug Dealer or Other Stranger	0.8276		*		*		*	
Bought on the Internet	0.4634		*		0.2632		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.65C Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

Source for Past Month Nonmedical Use among Past Month Sedative Users¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	*	11	*	*	*	*	*	*
From More Than One Doctor	2	7	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	36	*	*	*	*	*	*
Bought from Friend or Relative	8	9	*	*	*	*	*	*
Took from Friend or Relative without Asking	7	5	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	8	8	*	*	*	*	*	*
Bought on the Internet	*	1	*	*	*	*	*	*
Some Other Way	6	4	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.65D Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source for Past Month Nonmedical Use among Past Month Sedative Users ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	*	4.12	*	*	*	*	*	*
From More Than One Doctor	0.58	2.67	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	5.35	*	*	*	*	*	*
Bought from Friend or Relative	2.59	3.39	*	*	*	*	*	*
Took from Friend or Relative without Asking	2.07	2.07	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	2.38	3.00	*	*	*	*	*	*
Bought on the Internet	*	0.24	*	*	*	*	*	*
Some Other Way	1.80	1.79	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.65P Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007

Source for Past Month Nonmedical Use among Past Month Sedative Users ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	*		*		*		*	
From More Than One Doctor	0.1368		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	0.4589		*		*		*	
Took from Friend or Relative without Asking	0.3629		*		*		*	
Bought from Drug Dealer or Other Stranger	0.3597		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.4591		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.66C Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Sedatives ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.66D Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2006-2007 and 2007-2008

Source Where Friend or Relative Obtained Sedatives ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.66P Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2007-2008 Versus 2006-2007

Source Where Friend or Relative Obtained Sedatives ¹	Total (2006-2007)	Total (2007-2008)	Aged 12-17 (2006-2007)	Aged 12-17 (2007-2008)	Aged 18-25 (2006-2007)	Aged 18-25 (2007-2008)	Aged 26+ (2006-2007)	Aged 26+ (2007-2008)
From One Doctor	*		*		*		*	
From More Than One Doctor	*		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.