

Table 6.5A (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Numbers in Thousands, 2008

Age Category	Total	Male	Female
TOTAL	8,331	2,868	5,463
18	161	43	118
19	141	32	109
20	157	66	92
21	139	34	105
22	162	54	107
23	172	59	113
24	161	36	124
25	143	50	94
26-29	789	264	526
30-34	895	338	556
35-39	1,113	337	776
40-44	928	384	544
45-49	1,038	410	628
50-54	814	175	638
55-59	520	179	341
60-64	525	207	317
65 or Older	474	198	276

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.5B (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Percentages, 2008

Age Category	Total	Male	Female
TOTAL	3.7	2.6	4.7
18	3.3	1.7	4.9
19	3.2	1.4	5.3
20	4.0	3.3	4.7
21	3.5	1.7	5.3
22	4.0	2.7	5.3
23	4.2	2.9	5.5
24	4.2	1.9	6.3
25	3.8	2.7	4.8
26-29	4.6	3.0	6.3
30-34	4.8	3.8	5.8
35-39	5.5	3.4	7.4
40-44	4.3	3.7	5.0
45-49	4.6	3.6	5.6
50-54	3.7	1.7	5.5
55-59	2.9	2.0	3.8
60-64	3.5	2.8	4.0
65 or Older	1.3	1.2	1.3

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.6A (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics:
Numbers in Thousands, 2008

Demographic Characteristic	Total	Aged 18-25	Aged 26-49	Aged 50+
TOTAL	8,331	1,236	4,763	2,332
GENDER				
Male	2,868	374	1,734	760
Female	5,463	862	3,029	1,572
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	7,553	1,005	4,394	2,154
White	6,515	798	3,757	1,959
Black or African American	657	127	440	90
American Indian or Alaska Native	28	*	19	*
Native Hawaiian or Other Pacific Islander	9	*	*	*
Asian	220	38	95	*
Two or More Races	124	27	83	*
Hispanic or Latino	778	231	368	179
EDUCATION				
< High School	1,171	208	670	293
High School Graduate	2,726	425	1,502	799
Some College	2,635	468	1,527	641
College Graduate	1,799	135	1,064	600
CURRENT EMPLOYMENT				
Full-Time	3,541	508	2,524	510
Part-Time	1,127	336	551	240
Unemployed	605	154	321	*
Other ¹	3,058	238	1,367	1,453

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.6B (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics: Percentages, 2008

Demographic Characteristic	Total	Aged 18-25	Aged 26-49	Aged 50+
TOTAL	3.7	3.8	4.8	2.5
GENDER				
Male	2.6	2.3	3.5	1.8
Female	4.7	5.3	6.0	3.2
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	3.9	3.7	5.3	2.5
White	4.2	3.9	5.9	2.8
Black or African American	2.5	2.7	3.6	1.0
American Indian or Alaska Native	3.0	*	4.8	*
Native Hawaiian or Other Pacific Islander	1.2	*	*	*
Asian	2.2	2.6	1.8	*
Two or More Races	4.9	6.5	9.5	*
Hispanic or Latino	2.6	4.0	2.2	2.3
EDUCATION				
< High School	3.3	3.5	5.0	1.8
High School Graduate	3.9	3.7	5.3	2.6
Some College	4.6	4.3	5.8	3.2
College Graduate	2.9	2.9	3.3	2.3
CURRENT EMPLOYMENT				
Full-Time	2.9	3.4	3.5	1.4
Part-Time	3.7	3.9	5.2	2.1
Unemployed	6.7	5.0	7.4	*
Other ¹	4.8	3.7	10.1	3.3

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.7A (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Geographic Characteristics:
Numbers in Thousands, 2008

Geographic Characteristic	Total	Aged 18-25	Aged 26-49	Aged 50+
TOTAL	8,331	1,236	4,763	2,332
GEOGRAPHIC REGION				
Northeast	1,619	229	856	534
New England	398	43	225	130
Middle Atlantic	1,221	186	631	404
Midwest	1,968	273	1,197	498
East North Central	1,424	197	877	350
West North Central	545	76	320	148
South	2,785	415	1,624	747
South Atlantic	1,396	217	830	350
East South Central	443	57	290	96
West South Central	946	140	505	301
West	1,959	319	1,086	553
Mountain	727	107	399	221
Pacific	1,232	212	687	332
COUNTY TYPE				
Large Metro	4,301	675	2,452	1,174
Small Metro	2,692	397	1,533	763
250K – 1 Mil. Pop.	1,771	264	1,046	460
< 250K Pop.	921	132	486	303
Nonmetro	1,338	164	778	395
Urbanized	631	90	316	225
Less Urbanized	620	67	386	167
Completely Rural	86	8	76	2

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.7B (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Geographic Characteristics: Percentages, 2008

Geographic Characteristic	Total	Aged 18-25	Aged 26-49	Aged 50+
TOTAL	3.7	3.8	4.8	2.5
GEOGRAPHIC REGION				
Northeast	3.9	3.8	4.7	3.0
New England	3.6	2.8	4.7	2.8
Middle Atlantic	4.0	4.2	4.7	3.1
Midwest	4.0	3.8	5.5	2.4
East North Central	4.1	3.9	5.8	2.4
West North Central	3.7	3.4	5.0	2.4
South	3.4	3.5	4.5	2.2
South Atlantic	3.2	3.6	4.4	1.9
East South Central	3.3	3.1	5.0	1.7
West South Central	3.8	3.6	4.4	3.1
West	3.8	4.0	4.6	2.7
Mountain	4.6	4.5	5.6	3.5
Pacific	3.4	3.8	4.2	2.4
COUNTY TYPE				
Large Metro	3.6	3.8	4.4	2.6
Small Metro	4.0	3.9	5.3	2.6
250K – 1 Mil. Pop.	3.9	4.0	5.5	2.4
< 250K Pop.	4.0	3.7	5.0	3.1
Nonmetro	3.6	3.3	5.2	2.3
Urbanized	4.1	3.9	5.2	3.2
Less Urbanized	3.4	2.8	5.3	2.0
Completely Rural	2.4	2.3	5.1	0.1

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.