

Table 7.24C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2007 and 2008

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
TOBACCO PRODUCTS¹	N/A	N/A	364	401
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	147	163	246	264
Cigars	252	261	251	266
Pipe Tobacco	124	88	117	113
ALCOHOL	721	700	1,355	1,424
Binge Alcohol Use ²	558	534	658	660
Heavy Alcohol Use ²	317	308	300	285

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.24D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2007 and 2008

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
TOBACCO PRODUCTS¹	N/A	N/A	0.19	0.21
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.24	0.27	0.13	0.14
Cigars	0.39	0.42	0.13	0.14
Pipe Tobacco	0.20	0.15	0.06	0.06
ALCOHOL	0.73	0.72	0.48	0.46
Binge Alcohol Use ²	0.68	0.73	0.32	0.30
Heavy Alcohol Use ²	0.48	0.49	0.16	0.15

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.24P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
TOBACCO PRODUCTS¹	N/A		0.8682	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.3331		0.3016	
Cigars	0.5111		0.2596	
Pipe Tobacco	0.2390		0.9931	
ALCOHOL	0.6074		0.4118	
Binge Alcohol Use ²	0.6926		0.7194	
Heavy Alcohol Use ²	0.6116		0.9227	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.25C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2007 and 2008

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
TOBACCO PRODUCTS¹	N/A	N/A	37	34
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	24	22	25	20
Cigars	36	32	27	27
Pipe Tobacco	17	14	8	10
ALCOHOL	57	49	75	71
Binge Alcohol Use ²	51	42	52	45
Heavy Alcohol Use ²	31	24	21	18

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.25D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Percentages, 2007 and 2008

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
TOBACCO PRODUCTS¹	N/A	N/A	0.16	0.15
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.90	0.92	0.11	0.09
Cigars	1.23	1.25	0.12	0.12
Pipe Tobacco	0.67	0.61	0.04	0.04
ALCOHOL	1.34	1.38	0.30	0.29
Binge Alcohol Use ²	1.37	1.37	0.22	0.19
Heavy Alcohol Use ²	1.11	0.98	0.09	0.08

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.25P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
TOBACCO PRODUCTS¹	N/A		0.2037	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.9076		0.2314	
Cigars	0.3451		0.5069	
Pipe Tobacco	0.8361		0.8290	
ALCOHOL	0.2754		0.0405	
Binge Alcohol Use ²	0.2237		0.1758	
Heavy Alcohol Use ²	0.0515		0.8793	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.26C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2007 and 2008

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
TOBACCO PRODUCTS¹	N/A	N/A	76	73
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	55	55	38	40
Cigars	86	82	65	59
Pipe Tobacco	27	27	19	19
ALCOHOL	193	186	245	244
Binge Alcohol Use ²	173	166	170	167
Heavy Alcohol Use ²	104	110	74	77

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.26D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Percentages, 2007 and 2008

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
TOBACCO PRODUCTS¹	N/A	N/A	0.34	0.31
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.42	0.44	0.18	0.19
Cigars	0.58	0.59	0.30	0.26
Pipe Tobacco	0.22	0.23	0.09	0.09
ALCOHOL	0.64	0.62	0.67	0.65
Binge Alcohol Use ²	0.77	0.74	0.58	0.58
Heavy Alcohol Use ²	0.68	0.74	0.32	0.33

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.26P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
TOBACCO PRODUCTS¹	N/A		0.8865	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.8558		0.2344	
Cigars	0.6098		0.2174	
Pipe Tobacco	0.5071		0.1642	
ALCOHOL	0.8122		0.9770	
Binge Alcohol Use ²	0.4700		0.0778	
Heavy Alcohol Use ²	0.7375		0.8128	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.27C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2007 and 2008

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
TOBACCO PRODUCTS¹	N/A	N/A	347	389
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	132	149	238	260
Cigars	226	246	233	258
Pipe Tobacco	120	84	115	111
ALCOHOL	660	665	1,306	1,373
Binge Alcohol Use ²	504	497	619	628
Heavy Alcohol Use ²	293	284	289	277

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.27D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2007 and 2008

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
TOBACCO PRODUCTS¹	N/A	N/A	0.23	0.26
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.29	0.32	0.16	0.18
Cigars	0.47	0.52	0.16	0.17
Pipe Tobacco	0.26	0.18	0.08	0.08
ALCOHOL	0.93	0.91	0.60	0.59
Binge Alcohol Use ²	0.84	0.90	0.40	0.38
Heavy Alcohol Use ²	0.59	0.60	0.20	0.19

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

Table 7.27P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2007	2008	2007	2008
TOBACCO PRODUCTS¹	N/A		0.7726	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.2095		0.3403	
Cigars	0.2640		0.4037	
Pipe Tobacco	0.1774		0.8001	
ALCOHOL	0.5749		0.3839	
Binge Alcohol Use ²	0.6633		0.4371	
Heavy Alcohol Use ²	0.4073		0.9112	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.