

**Table 7.19C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2007 and 2008**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2007	2008	2007	2008	2007	2008	2007	2008
<b>ILLICIT DRUGS<sup>1</sup></b>	306	291	124	147	245	248	153	138
Marijuana and Hashish	258	274	96	130	199	214	97	91
Cocaine	121	100	52	21	49	67	25	17
Crack	61	49	46	10	21	17	5	17
Heroin	31	35	18	10	4	14	*	*
Hallucinogens	60	59	25	14	33	23	17	23
LSD	32	15	5	6	4	7	5	6
PCP	9	7	8	*	*	3	4	*
Ecstasy	46	48	8	9	27	14	11	7
Inhalants	35	47	4	10	44	46	34	25
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	151	128	83	89	143	122	119	98
Pain Relievers	132	109	72	70	113	113	108	88
OxyContin <sup>®</sup>	30	44	40	36	9	11	17	7
Tranquilizers	74	76	47	57	74	55	52	32
Stimulants <sup>3</sup>	69	49	48	27	31	36	27	22
Methamphetamine <sup>3</sup>	63	25	47	24	17	28	9	16
Sedatives	26	23	15	9	54	19	29	21
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	195	164	99	92	166	142	124	104

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.19D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2007 and 2008**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2007	2008	2007	2008	2007	2008	2007	2008
<b>ILLICIT DRUGS<sup>1</sup></b>	0.66	0.66	0.61	0.72	0.27	0.27	0.15	0.13
Marijuana and Hashish	0.58	0.63	0.48	0.64	0.22	0.23	0.10	0.09
Cocaine	0.29	0.25	0.26	0.11	0.06	0.08	0.02	0.02
Crack	0.15	0.12	0.23	0.05	0.02	0.02	0.00	0.02
Heroin	0.08	0.09	0.09	0.05	0.00	0.02	*	*
Hallucinogens	0.15	0.15	0.13	0.07	0.04	0.03	0.02	0.02
LSD	0.08	0.04	0.03	0.03	0.01	0.01	0.00	0.01
PCP	0.02	0.02	0.04	*	*	0.00	0.00	*
Ecstasy	0.12	0.12	0.04	0.05	0.03	0.02	0.01	0.01
Inhalants	0.09	0.12	0.02	0.05	0.05	0.05	0.03	0.02
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.35	0.31	0.41	0.45	0.16	0.14	0.12	0.10
Pain Relievers	0.31	0.26	0.36	0.36	0.13	0.13	0.11	0.09
OxyContin <sup>®</sup>	0.08	0.11	0.20	0.18	0.01	0.01	0.02	0.01
Tranquilizers	0.18	0.19	0.24	0.29	0.08	0.06	0.05	0.03
Stimulants <sup>3</sup>	0.17	0.12	0.24	0.14	0.04	0.04	0.03	0.02
Methamphetamine <sup>3</sup>	0.15	0.06	0.23	0.12	0.02	0.03	0.01	0.02
Sedatives	0.06	0.06	0.08	0.05	0.06	0.02	0.03	0.02
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.44	0.39	0.48	0.46	0.19	0.16	0.12	0.10

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.19P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2007	2008	2007	2008	2007	2008	2007	2008
<b>ILLICIT DRUGS<sup>1</sup></b>	0.9330		0.7297		0.4650		0.2158	
Marijuana and Hashish	0.3127		0.6263		0.3187		0.6902	
Cocaine	0.2226		0.0172		0.1854		0.5658	
Crack	0.0954		0.0049		0.8892		0.4937	
Heroin	0.1948		0.4668		0.3797		*	
Hallucinogens	0.3865		0.5786		0.8741		0.7420	
LSD	0.6784		0.2211		0.1101		0.9776	
PCP	0.6424		*		*		*	
Ecstasy	0.3177		0.8782		0.6676		0.6036	
Inhalants	0.9610		0.0825		0.4827		0.3314	
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.0402		0.5487		0.4348		0.1024	
Pain Relievers	0.0536		0.9545		0.9120		0.2934	
OxyContin <sup>®</sup>	0.3129		0.6329		0.8202		0.4013	
Tranquilizers	0.6672		0.1942		0.3015		0.1359	
Stimulants <sup>3</sup>	0.0937		0.4890		0.5029		0.9172	
Methamphetamine <sup>3</sup>	0.0011		0.5172		0.3766		0.5975	
Sedatives	0.3366		0.6753		0.3425		0.5960	
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.0739		0.8391		0.7987		0.1036	

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.20C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2007 and 2008**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2007	2008	2007	2008	2007	2008	2007	2008
<b>ILLICIT DRUGS<sup>1</sup></b>	40	40	23	22	32	34	42	34
Marijuana and Hashish	38	38	22	20	28	29	24	23
Cocaine	10	11	4	4	4	5	6	2
Crack	4	2	1	0	1	1	3	*
Heroin	1	6	0	3	1	*	*	*
Hallucinogens	14	15	6	9	7	13	6	6
LSD	6	7	4	5	1	4	2	3
PCP	3	3	*	*	1	3	3	1
Ecstasy	8	11	4	8	5	6	3	3
Inhalants	12	11	4	6	10	12	20	17
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	23	19	11	12	15	17	26	22
Pain Relievers	20	17	10	11	15	13	23	20
OxyContin <sup>®</sup>	6	6	3	4	4	2	3	3
Tranquilizers	15	11	4	5	6	7	11	9
Stimulants <sup>3</sup>	9	7	6	4	4	10	7	5
Methamphetamine <sup>3</sup>	5	5	3	2	3	2	4	1
Sedatives	3	5	3	4	2	3	8	4
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	26	26	13	15	18	22	35	26

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.20D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2007 and 2008**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2007	2008	2007	2008	2007	2008	2007	2008
<b>ILLICIT DRUGS<sup>1</sup></b>	1.90	1.81	1.99	2.01	1.12	1.20	0.20	0.16
Marijuana and Hashish	1.88	1.87	1.89	1.91	1.01	1.09	0.12	0.11
Cocaine	0.63	0.79	0.43	0.44	0.18	0.24	0.03	0.01
Crack	0.28	0.19	0.15	0.04	0.06	0.04	0.01	*
Heroin	0.06	0.44	0.03	0.28	0.06	*	*	*
Hallucinogens	0.88	1.08	0.62	0.94	0.28	0.54	0.03	0.03
LSD	0.41	0.54	0.41	0.52	0.06	0.17	0.01	0.02
PCP	0.21	0.19	*	*	0.06	0.12	0.01	0.01
Ecstasy	0.54	0.79	0.38	0.86	0.22	0.27	0.01	0.02
Inhalants	0.74	0.77	0.41	0.64	0.39	0.50	0.10	0.08
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	1.38	1.31	1.09	1.22	0.60	0.71	0.13	0.10
Pain Relievers	1.23	1.17	0.98	1.12	0.58	0.57	0.11	0.10
OxyContin <sup>®</sup>	0.40	0.45	0.35	0.40	0.15	0.10	0.01	0.01
Tranquilizers	0.93	0.81	0.40	0.51	0.24	0.28	0.05	0.05
Stimulants <sup>3</sup>	0.61	0.55	0.58	0.43	0.18	0.43	0.03	0.02
Methamphetamine <sup>3</sup>	0.34	0.39	0.32	0.17	0.12	0.10	0.02	0.00
Sedatives	0.19	0.38	0.35	0.40	0.07	0.13	0.04	0.02
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	1.50	1.63	1.30	1.50	0.68	0.89	0.17	0.13

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.20P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2007	2008	2007	2008	2007	2008	2007	2008
<b>ILLICIT DRUGS<sup>1</sup></b>	0.0679		0.5936		0.0026		0.0282	
Marijuana and Hashish	0.1514		0.4160		0.0036		0.3573	
Cocaine	0.5378		0.9329		0.3729		0.0962	
Crack	0.5423		0.2730		0.6594		*	
Heroin	0.1019		0.3341		*		*	
Hallucinogens	0.0694		0.2194		0.0342		0.3495	
LSD	0.3418		0.2414		0.1050		0.8382	
PCP	0.6066		*		0.1352		0.6602	
Ecstasy	0.0280		0.1420		0.6953		0.5665	
Inhalants	0.9344		0.2605		0.8605		0.3924	
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.3667		0.5497		0.2598		0.0637	
Pain Relievers	0.2871		0.8581		0.6758		0.3626	
OxyContin <sup>®</sup>	0.5746		0.5955		0.5079		0.4998	
Tranquilizers	0.3005		0.6968		0.6449		0.5668	
Stimulants <sup>3</sup>	0.7728		0.0095		0.0439		0.3758	
Methamphetamine <sup>3</sup>	0.1958		0.1564		0.7894		0.2200	
Sedatives	0.1098		0.8794		0.3812		0.2707	
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.6700		0.9952		0.0533		0.0448	

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.21C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2007 and 2008**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2007	2008	2007	2008	2007	2008	2007	2008
<b>ILLICIT DRUGS<sup>1</sup></b>	115	115	37	33	81	81	27	29
Marijuana and Hashish	106	107	34	28	73	72	24	25
Cocaine	35	35	6	7	14	14	3	4
Crack	10	8	5	3	1	4	*	3
Heroin	9	12	4	6	3	1	*	1
Hallucinogens	30	34	7	7	15	16	5	6
LSD	10	13	4	4	4	6	5	1
PCP	3	3	4	*	*	*	3	*
Ecstasy	22	28	5	5	7	10	3	6
Inhalants	15	15	1	1	7	9	9	3
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	59	59	18	18	37	36	16	17
Pain Relievers	53	51	15	16	31	31	14	16
OxyContin <sup>®</sup>	16	14	3	6	7	5	4	3
Tranquilizers	31	32	11	9	18	18	7	6
Stimulants <sup>3</sup>	26	23	9	7	13	15	6	4
Methamphetamine <sup>3</sup>	17	9	5	6	2	7	3	4
Sedatives	11	6	5	2	4	13	2	4
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	70	70	19	20	42	43	19	18

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.21D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2007 and 2008**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2007	2008	2007	2008	2007	2008	2007	2008
<b>ILLICIT DRUGS<sup>1</sup></b>	0.79	0.88	1.28	1.23	0.62	0.62	0.26	0.28
Marijuana and Hashish	0.78	0.85	1.22	1.10	0.58	0.57	0.24	0.24
Cocaine	0.36	0.36	0.25	0.29	0.13	0.13	0.03	0.04
Crack	0.10	0.09	0.19	0.14	0.01	0.04	*	0.03
Heroin	0.10	0.13	0.18	0.23	0.03	0.01	*	0.01
Hallucinogens	0.31	0.35	0.29	0.31	0.13	0.15	0.05	0.06
LSD	0.11	0.14	0.14	0.17	0.04	0.05	0.04	0.01
PCP	0.03	0.03	0.17	*	*	*	0.03	*
Ecstasy	0.23	0.29	0.21	0.19	0.07	0.09	0.03	0.06
Inhalants	0.15	0.16	0.03	0.04	0.07	0.08	0.08	0.03
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.56	0.56	0.70	0.73	0.33	0.32	0.15	0.16
Pain Relievers	0.51	0.49	0.58	0.66	0.28	0.28	0.14	0.15
OxyContin <sup>®</sup>	0.17	0.14	0.13	0.23	0.07	0.05	0.04	0.03
Tranquilizers	0.33	0.33	0.43	0.35	0.17	0.17	0.06	0.06
Stimulants <sup>3</sup>	0.27	0.24	0.35	0.30	0.12	0.14	0.06	0.04
Methamphetamine <sup>3</sup>	0.18	0.10	0.20	0.26	0.02	0.06	0.03	0.04
Sedatives	0.12	0.06	0.22	0.09	0.04	0.12	0.02	0.04
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.63	0.63	0.74	0.81	0.37	0.37	0.18	0.17

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.



**Table 7.21P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2007	2008	2007	2008	2007	2008	2007	2008
<b>ILLICIT DRUGS<sup>1</sup></b>	0.2740		0.9230		0.4130		0.4208	
Marijuana and Hashish	0.8694		0.8395		0.7504		0.2071	
Cocaine	0.1149		0.9692		0.8558		0.3032	
Crack	0.4230		0.5688		0.1679		*	
Heroin	0.1146		0.6699		0.4338		*	
Hallucinogens	0.2186		0.8274		0.3990		0.9477	
LSD	0.1286		0.6508		0.4702		0.2210	
PCP	0.6791		*		*		*	
Ecstasy	0.1106		0.5767		0.1021		0.4641	
Inhalants	0.8630		0.9303		0.7312		0.0484	
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.3874		0.9216		0.3216		0.9138	
Pain Relievers	0.6117		0.8888		0.7508		0.7849	
OxyContin <sup>®</sup>	0.4515		0.6988		0.5925		0.1710	
Tranquilizers	0.9200		0.7984		0.9516		0.9320	
Stimulants <sup>3</sup>	0.3605		0.9155		0.1075		0.6531	
Methamphetamine <sup>3</sup>	0.0062		0.7912		0.3317		0.6518	
Sedatives	0.0227		0.6756		0.1613		0.2811	
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.1654		0.6266		0.3130		0.4997	

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.22C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2007 and 2008**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2007	2008	2007	2008	2007	2008	2007	2008
<b>ILLICIT DRUGS<sup>1</sup></b>	262	262	120	138	226	229	142	129
Marijuana and Hashish	219	248	92	124	182	194	91	85
Cocaine	115	90	51	20	46	66	24	16
Crack	60	48	46	9	21	17	4	16
Heroin	30	33	17	8	*	14	*	*
Hallucinogens	48	47	23	7	29	10	15	22
LSD	30	*	*	*	*	*	*	5
PCP	8	6	6	*	*	*	*	*
Ecstasy	39	37	5	*	26	8	10	*
Inhalants	30	44	2	8	43	43	26	18
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	136	108	81	86	136	114	115	93
Pain Relievers	119	92	71	67	106	106	104	85
OxyContin <sup>®</sup>	25	42	40	35	3	9	16	5
Tranquilizers	65	66	46	56	72	52	51	30
Stimulants <sup>3</sup>	64	42	47	25	28	32	24	21
Methamphetamine <sup>3</sup>	60	22	47	23	17	28	7	16
Sedatives	24	22	14	8	53	14	28	20
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	178	140	97	88	157	135	118	99

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.22D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2007 and 2008**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2007	2008	2007	2008	2007	2008	2007	2008
<b>ILLICIT DRUGS<sup>1</sup></b>	0.80	0.81	0.70	0.82	0.30	0.30	0.20	0.18
Marijuana and Hashish	0.70	0.78	0.54	0.74	0.24	0.25	0.13	0.12
Cocaine	0.38	0.30	0.31	0.12	0.06	0.09	0.03	0.02
Crack	0.20	0.16	0.28	0.06	0.03	0.02	0.01	0.02
Heroin	0.10	0.11	0.11	0.05	*	0.02	*	*
Hallucinogens	0.16	0.16	0.14	0.05	0.04	0.01	0.02	0.03
LSD	0.10	*	*	*	*	*	*	0.01
PCP	0.03	0.02	0.04	*	*	*	*	*
Ecstasy	0.13	0.12	0.03	*	0.04	0.01	0.01	*
Inhalants	0.10	0.15	0.01	0.05	0.06	0.06	0.04	0.02
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.44	0.36	0.48	0.53	0.18	0.15	0.16	0.13
Pain Relievers	0.39	0.31	0.42	0.41	0.14	0.14	0.15	0.12
OxyContin <sup>®</sup>	0.09	0.14	0.24	0.22	0.00	0.01	0.02	0.01
Tranquilizers	0.22	0.22	0.28	0.35	0.10	0.07	0.07	0.04
Stimulants <sup>3</sup>	0.22	0.14	0.28	0.16	0.04	0.04	0.03	0.03
Methamphetamine <sup>3</sup>	0.20	0.07	0.28	0.14	0.02	0.04	0.01	0.02
Sedatives	0.08	0.07	0.09	0.05	0.07	0.02	0.04	0.03
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.57	0.46	0.57	0.53	0.21	0.18	0.17	0.14

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.22P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2008 Versus 2007**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2007	2008	2007	2008	2007	2008	2007	2008
<b>ILLICIT DRUGS<sup>1</sup></b>	0.6090		0.6588		0.7369		0.3662	
Marijuana and Hashish	0.1958		0.4689		0.4606		0.5778	
Cocaine	0.4132		0.0150		0.2124		0.6698	
Crack	0.1267		0.0061		0.7253		0.5141	
Heroin	0.5480		0.2814		*		*	
Hallucinogens	1.0000		0.2013		0.1085		0.9741	
LSD	*		*		*		*	
PCP	0.6130		*		*		*	
Ecstasy	0.8703		*		0.2062		*	
Inhalants	0.8466		0.1595		0.3948		0.9872	
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.1131		0.4961		0.2469		0.1963	
Pain Relievers	0.1124		1.0000		0.8779		0.3517	
OxyContin <sup>®</sup>	0.1635		0.6993		0.1955		0.5305	
Tranquilizers	0.4596		0.1809		0.2832		0.1638	
Stimulants <sup>3</sup>	0.1847		0.7189		0.6524		0.8229	
Methamphetamine <sup>3</sup>	0.0066		0.5460		0.4628		0.4651	
Sedatives	0.6816		0.7920		0.1757		0.6793	
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.1993		0.7642		0.4808		0.3044	

\*Low precision; no estimate reported.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.23C Illicit Drug Use in the Past Month, by Past Month Cigarette Use and Levels of Past Month Alcohol Use and Age Group: Standard Errors of Numbers in Thousands, 2007 and 2008**

<b>Cigarette Use/Level of Alcohol Use</b>	<b>Total (2007)</b>	<b>Total (2008)</b>	<b>Aged 12-17 (2007)</b>	<b>Aged 12-17 (2008)</b>	<b>Aged 18-25 (2007)</b>	<b>Aged 18-25 (2008)</b>	<b>Aged 26+ (2007)</b>	<b>Aged 26+ (2008)</b>
<b>TOTAL</b>	445	427	68	61	127	127	394	393
Heavy Alcohol Use <sup>1</sup>	219	195	27	26	90	91	191	167
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	232	227	37	35	91	85	206	208
Alcohol Use But Not Binge Use <sup>1</sup>	203	231	24	27	46	55	196	220
No Alcohol Use	195	200	51	41	45	45	184	190
<b>PAST MONTH CIGARETTE USE</b>	336	327	46	45	122	119	294	298
Heavy Alcohol Use <sup>1</sup>	175	155	25	21	75	75	151	134
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	187	168	28	28	68	70	167	154
Alcohol Use But Not Binge Use <sup>1</sup>	127	169	15	17	37	37	120	163
No Alcohol Use	124	147	23	22	37	33	120	138
<b>NO PAST MONTH CIGARETTE USE</b>	295	283	53	47	88	85	272	261
Heavy Alcohol Use <sup>1</sup>	129	105	10	14	40	42	122	95
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	128	155	23	23	55	47	116	146
Alcohol Use But Not Binge Use <sup>1</sup>	155	153	18	20	31	39	151	145
No Alcohol Use	153	138	42	34	27	29	142	129

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.23D Illicit Drug Use in the Past Month, by Past Month Cigarette Use and Levels of Past Month Alcohol Use and Age Group: Standard Errors of Percentages, 2007 and 2008**

<b>Cigarette Use/Level of Alcohol Use</b>	<b>Total (2007)</b>	<b>Total (2008)</b>	<b>Aged 12-17 (2007)</b>	<b>Aged 12-17 (2008)</b>	<b>Aged 18-25 (2007)</b>	<b>Aged 18-25 (2008)</b>	<b>Aged 26+ (2007)</b>	<b>Aged 26+ (2008)</b>
<b>TOTAL</b>	0.18	0.17	0.27	0.24	0.39	0.38	0.21	0.20
Heavy Alcohol Use <sup>1</sup>	1.03	1.03	2.85	2.76	1.17	1.16	1.38	1.30
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.53	0.53	1.53	1.63	0.78	0.79	0.66	0.65
Alcohol Use But Not Binge Use <sup>1</sup>	0.29	0.32	1.40	1.57	0.68	0.74	0.31	0.34
No Alcohol Use	0.16	0.16	0.23	0.18	0.34	0.35	0.21	0.22
<b>PAST MONTH CIGARETTE USE</b>	0.52	0.51	1.39	1.40	0.69	0.77	0.60	0.61
Heavy Alcohol Use <sup>1</sup>	1.39	1.44	3.33	3.22	1.40	1.42	1.96	1.89
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.97	0.94	2.43	2.35	1.13	1.25	1.26	1.23
Alcohol Use But Not Binge Use <sup>1</sup>	0.93	1.15	4.06	4.37	1.60	1.69	1.05	1.31
No Alcohol Use	0.61	0.72	1.99	2.01	1.28	1.23	0.70	0.82
<b>NO PAST MONTH CIGARETTE USE</b>	0.15	0.15	0.22	0.19	0.38	0.36	0.18	0.18
Heavy Alcohol Use <sup>1</sup>	1.59	1.35	4.95	5.14	1.85	1.82	2.07	1.65
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.53	0.62	1.89	1.99	1.01	0.95	0.63	0.75
Alcohol Use But Not Binge Use <sup>1</sup>	0.27	0.26	1.35	1.48	0.66	0.76	0.29	0.28
No Alcohol Use	0.15	0.13	0.20	0.16	0.26	0.28	0.20	0.18

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 7.23P Illicit Drug Use in the Past Month, by Past Month Cigarette Use and Levels of Past Month Alcohol Use and Age Group: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007**

<b>Cigarette Use/Level of Alcohol Use</b>	<b>Total (2007)</b>	<b>Total (2008)</b>	<b>Aged 12-17 (2007)</b>	<b>Aged 12-17 (2008)</b>	<b>Aged 18-25 (2007)</b>	<b>Aged 18-25 (2008)</b>	<b>Aged 26+ (2007)</b>	<b>Aged 26+ (2008)</b>
<b>TOTAL</b>	0.9192		0.4228		0.8154		0.7215	
Heavy Alcohol Use <sup>1</sup>	0.1844		0.0376		0.3254		0.3292	
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.6155		0.0448		0.5122		0.4787	
Alcohol Use But Not Binge Use <sup>1</sup>	0.1613		0.0540		0.0517		0.3795	
No Alcohol Use	0.4308		0.0227		0.6617		0.6800	
<b>PAST MONTH CIGARETTE USE</b>	0.7375		0.3908		0.3615		0.4609	
Heavy Alcohol Use <sup>1</sup>	0.1029		0.1550		0.1372		0.3225	
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.7254		0.1068		0.2004		0.7789	
Alcohol Use But Not Binge Use <sup>1</sup>	0.0243		0.4885		0.1091		0.0359	
No Alcohol Use	0.7297		0.5936		0.9230		0.6588	
<b>NO PAST MONTH CIGARETTE USE</b>	0.9213		0.6260		0.3055		0.9318	
Heavy Alcohol Use <sup>1</sup>	0.9246		0.0138		0.6928		0.6908	
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.1099		0.1855		0.9937		0.0709	
Alcohol Use But Not Binge Use <sup>1</sup>	0.8428		0.0435		0.0631		0.3946	
No Alcohol Use	0.2158		0.0282		0.4208		0.3662	

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.