

**Table 1.11C Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2007 and 2008**

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
<b>TOTAL</b>	971	928	596	554	445	427
12	27	30	20	24	15	15
13	38	35	30	28	18	16
14	42	42	35	35	23	22
15	49	53	43	43	32	29
16	57	54	49	46	37	32
17	59	62	51	52	38	37
18	84	84	72	71	52	53
19	86	87	72	74	53	54
20	90	81	74	67	54	49
21	92	85	75	69	57	50
22	81	86	60	64	46	47
23	81	93	60	56	42	41
24	86	81	59	59	45	43
25	77	80	49	52	40	39
26-29	309	337	185	196	135	139
30-34	319	322	176	171	136	138
35-39	368	360	183	179	129	136
40-44	376	418	170	179	125	120
45-49	407	423	169	169	128	135
50-54	550	539	241	206	184	154
55-59	502	484	179	184	132	159
60-64	342	406	130	148	80	116
65 or Older	296	325	87	103	71	94

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.11D Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2007 and 2008**

<b>Age Category</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	0.39	0.37	0.24	0.22	0.18	0.17
12	0.63	0.70	0.48	0.58	0.36	0.38
13	0.82	0.81	0.68	0.66	0.44	0.38
14	0.90	0.91	0.78	0.79	0.53	0.52
15	0.92	0.98	0.85	0.87	0.69	0.63
16	1.01	0.97	0.92	0.89	0.76	0.68
17	1.08	1.11	1.00	0.97	0.82	0.77
18	1.20	1.18	1.13	1.15	0.95	0.95
19	1.30	1.27	1.24	1.24	1.04	1.06
20	1.29	1.30	1.30	1.28	1.13	1.09
21	1.26	1.15	1.28	1.22	1.11	1.03
22	1.30	1.27	1.25	1.21	1.06	1.00
23	1.23	1.33	1.20	1.12	0.97	0.88
24	1.27	1.20	1.21	1.21	1.00	1.00
25	1.25	1.23	1.15	1.13	0.96	0.92
26-29	1.21	1.20	0.97	0.99	0.75	0.76
30-34	1.18	1.11	0.87	0.81	0.69	0.70
35-39	1.17	1.14	0.81	0.81	0.58	0.65
40-44	1.17	1.23	0.72	0.78	0.56	0.55
45-49	1.08	1.07	0.70	0.71	0.54	0.58
50-54	1.67	1.62	1.06	0.91	0.83	0.69
55-59	1.92	1.80	0.93	0.99	0.70	0.87
60-64	1.96	1.98	0.95	0.96	0.59	0.76
65 or Older	0.81	0.87	0.24	0.28	0.19	0.25

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.11P Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007**

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
<b>TOTAL</b>	0.1099		0.5763		0.9192	
12	0.1493		0.0121		0.4312	
13	0.2966		0.7500		0.3771	
14	0.9195		0.8964		0.9378	
15	0.6552		0.4028		0.5796	
16	0.2091		0.3909		0.2160	
17	0.7523		0.4625		0.7141	
18	0.7135		0.9790		0.7787	
19	0.6492		0.5991		0.9416	
20	0.4727		0.8480		0.8527	
21	0.4473		0.2432		0.1317	
22	0.7930		0.2226		0.6321	
23	0.0467		0.1012		0.8523	
24	0.2128		0.6241		0.3920	
25	0.3619		0.0432		0.9538	
26-29	0.0486		0.7822		0.8310	
30-34	0.9589		0.6671		0.8189	
35-39	0.5760		0.5810		0.1473	
40-44	0.3742		0.8752		0.3715	
45-49	0.2833		0.6474		0.8095	
50-54	0.6870		0.0797		0.1794	
55-59	0.9213		0.7985		0.4191	
60-64	0.0249		0.5762		0.2488	
65 or Older	0.0240		0.3806		0.2430	

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.12C Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2007 and 2008**

<b>Age Category</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	963	917	492	474	360	377
12	9	9	9	8	7	6
13	18	20	15	19	9	12
14	29	26	25	24	17	15
15	39	43	34	36	25	25
16	48	44	42	38	31	28
17	54	55	45	46	35	33
18	75	75	65	64	49	50
19	78	80	65	67	50	50
20	86	77	69	64	49	45
21	84	79	66	64	50	46
22	74	81	55	58	42	43
23	78	87	53	49	39	36
24	83	78	55	52	40	37
25	72	76	45	46	36	33
26-29	290	316	160	176	116	131
30-34	290	307	145	150	114	119
35-39	340	348	144	145	110	113
40-44	352	394	136	136	102	92
45-49	391	396	138	146	109	107
50-54	532	518	198	161	154	139
55-59	487	464	129	156	90	145
60-64	318	378	91	119	39	106
65 or Older	261	276	47	54	41	51

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.12D Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2007 and 2008**

<b>Age Category</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	0.39	0.37	0.20	0.19	0.15	0.15
12	0.23	0.23	0.21	0.22	0.17	0.15
13	0.43	0.49	0.35	0.46	0.21	0.29
14	0.65	0.61	0.58	0.57	0.40	0.37
15	0.78	0.87	0.72	0.76	0.55	0.56
16	0.94	0.89	0.83	0.79	0.66	0.60
17	1.05	1.02	0.93	0.90	0.77	0.71
18	1.17	1.12	1.10	1.07	0.92	0.91
19	1.28	1.25	1.15	1.16	1.01	1.01
20	1.31	1.35	1.24	1.29	1.05	1.02
21	1.31	1.18	1.22	1.17	1.04	0.98
22	1.32	1.36	1.21	1.17	0.99	0.95
23	1.28	1.33	1.12	1.03	0.91	0.80
24	1.35	1.25	1.17	1.14	0.92	0.86
25	1.30	1.27	1.07	1.04	0.90	0.79
26-29	1.20	1.24	0.87	0.93	0.66	0.72
30-34	1.16	1.15	0.74	0.73	0.58	0.61
35-39	1.16	1.21	0.66	0.69	0.51	0.55
40-44	1.13	1.23	0.61	0.60	0.47	0.42
45-49	1.10	1.09	0.58	0.63	0.47	0.47
50-54	1.68	1.63	0.90	0.71	0.70	0.63
55-59	1.92	1.80	0.68	0.85	0.48	0.79
60-64	1.89	1.96	0.67	0.78	0.29	0.69
65 or Older	0.72	0.74	0.13	0.15	0.11	0.14

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.12P Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007**

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
<b>TOTAL</b>	0.4107		0.4709		0.2151	
12	0.5866		0.6795		0.8299	
13	0.3758		0.1103		0.7652	
14	0.8549		0.6654		0.7870	
15	0.2869		0.6795		0.9108	
16	0.4673		0.5742		0.6261	
17	0.7045		0.5023		0.6939	
18	0.4945		0.4963		0.5878	
19	0.6536		0.6081		0.7449	
20	0.5891		0.9897		0.8231	
21	0.5140		0.7191		0.4210	
22	0.7829		0.4734		0.6363	
23	0.0534		0.0823		0.6081	
24	0.5669		0.9626		0.6410	
25	0.0977		0.3107		0.9639	
26-29	0.1170		0.5119		0.4544	
30-34	0.7516		0.4934		0.2905	
35-39	0.4040		0.2470		0.4189	
40-44	0.3454		0.9140		0.5214	
45-49	0.6671		0.7330		0.7940	
50-54	0.4640		0.2054		0.4274	
55-59	0.6397		0.5304		0.0627	
60-64	0.0691		0.2219		0.0267	
65 or Older	0.3580		0.5770		0.5536	

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.13C Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2007 and 2008**

<b>Age Category</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	659	668	242	207	149	131
12	3	1	2	1	2	*
13	5	5	5	3	3	1
14	6	5	6	4	3	2
15	13	13	13	12	4	8
16	19	15	14	12	7	5
17	21	24	18	16	9	8
18	37	31	30	24	11	11
19	36	40	28	30	14	15
20	40	38	29	26	15	15
21	42	41	28	27	17	19
22	42	40	26	25	11	10
23	47	44	27	23	11	10
24	42	41	22	22	12	11
25	44	40	25	18	14	11
26-29	166	188	83	93	46	52
30-34	169	151	76	76	50	55
35-39	212	199	70	67	42	39
40-44	245	243	59	60	42	34
45-49	263	280	84	69	66	47
50-54	373	358	111	59	54	39
55-59	255	309	60	33	38	29
60-64	163	183	2	51	*	26
65 or Older	110	107	39	16	34	16

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.13D Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2007 and 2008**

<b>Age Category</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	0.27	0.27	0.10	0.08	0.06	0.05
12	0.08	0.03	0.05	0.02	0.04	*
13	0.12	0.13	0.12	0.07	0.08	0.02
14	0.16	0.12	0.15	0.10	0.07	0.06
15	0.30	0.30	0.28	0.28	0.10	0.18
16	0.41	0.34	0.31	0.28	0.17	0.13
17	0.49	0.54	0.41	0.36	0.21	0.18
18	0.74	0.61	0.60	0.48	0.22	0.22
19	0.78	0.85	0.61	0.65	0.32	0.35
20	0.89	0.89	0.67	0.63	0.36	0.36
21	0.94	0.94	0.66	0.64	0.40	0.48
22	0.97	0.90	0.66	0.59	0.29	0.26
23	1.06	0.96	0.65	0.55	0.28	0.25
24	0.96	0.95	0.56	0.54	0.30	0.28
25	1.07	0.97	0.66	0.47	0.38	0.30
26-29	0.90	1.02	0.49	0.53	0.28	0.30
30-34	0.83	0.76	0.40	0.41	0.27	0.29
35-39	0.89	0.91	0.33	0.33	0.20	0.19
40-44	0.98	0.99	0.28	0.28	0.20	0.16
45-49	0.95	0.99	0.37	0.31	0.29	0.21
50-54	1.49	1.39	0.51	0.27	0.25	0.18
55-59	1.26	1.51	0.33	0.19	0.20	0.16
60-64	1.18	1.15	0.01	0.34	*	0.17
65 or Older	0.30	0.29	0.11	0.04	0.09	0.04

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.



**Table 1.13P Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007**

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
<b>TOTAL</b>	0.5181		0.0964		0.2298	
12	0.1622		0.3146		*	
13	0.8437		0.3805		0.3345	
14	0.5114		0.2137		0.7298	
15	0.8551		0.7770		0.0967	
16	0.0678		0.1299		0.2799	
17	0.3690		0.0692		0.6128	
18	0.5575		0.3160		0.3719	
19	0.3083		0.6472		0.8697	
20	0.7539		0.3627		0.9956	
21	0.9105		0.2444		0.6411	
22	0.5347		0.1033		0.1293	
23	0.1173		0.0840		0.2254	
24	0.2876		0.8421		0.5744	
25	0.4161		0.0612		0.5102	
26-29	0.3295		0.1918		0.5026	
30-34	0.3751		0.9045		0.9934	
35-39	0.4785		0.7076		0.9772	
40-44	0.6312		0.8620		0.5026	
45-49	0.2323		0.2437		0.1006	
50-54	0.5831		0.1238		0.8037	
55-59	0.0067		0.3911		0.8603	
60-64	0.1262		0.0684		*	
65 or Older	0.8595		0.1824		0.3217	

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.14C Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2007 and 2008**

<b>Age Category</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	339	322	123	94	80	55
12	*	*	*	*	*	*
13	2	*	2	*	0	*
14	4	3	3	1	2	*
15	6	6	6	5	2	2
16	7	6	6	3	3	2
17	8	9	7	4	3	1
18	10	13	6	10	4	4
19	15	12	8	7	5	3
20	15	17	10	8	3	4
21	20	16	6	7	1	6
22	19	18	9	10	4	2
23	18	16	8	5	4	3
24	18	17	10	6	3	3
25	18	16	8	6	5	3
26-29	68	85	29	35	21	10
30-34	93	85	27	34	15	28
35-39	103	106	40	34	29	12
40-44	126	135	39	36	29	19
45-49	135	135	60	44	43	33
50-54	174	159	66	34	29	20
55-59	111	133	37	8	27	*
60-64	43	49	*	23	*	9
65 or Older	62	16	23	16	13	*

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.14D Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2007 and 2008**

<b>Age Category</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	0.14	0.13	0.05	0.04	0.03	0.02
12	*	*	*	*	*	*
13	0.05	*	0.04	*	0.01	*
14	0.10	0.07	0.08	0.01	0.05	*
15	0.14	0.13	0.14	0.11	0.05	0.04
16	0.17	0.14	0.13	0.08	0.07	0.04
17	0.19	0.20	0.16	0.10	0.07	0.02
18	0.21	0.26	0.12	0.21	0.08	0.08
19	0.33	0.28	0.17	0.17	0.11	0.08
20	0.36	0.42	0.25	0.21	0.08	0.11
21	0.48	0.41	0.15	0.18	0.03	0.14
22	0.49	0.44	0.24	0.25	0.10	0.04
23	0.46	0.39	0.20	0.12	0.10	0.07
24	0.45	0.45	0.26	0.15	0.07	0.07
25	0.48	0.43	0.22	0.16	0.14	0.09
26-29	0.41	0.50	0.18	0.21	0.13	0.06
30-34	0.48	0.45	0.14	0.18	0.08	0.15
35-39	0.48	0.50	0.19	0.17	0.14	0.06
40-44	0.57	0.62	0.18	0.17	0.14	0.09
45-49	0.58	0.58	0.26	0.19	0.19	0.15
50-54	0.79	0.70	0.30	0.16	0.14	0.09
55-59	0.59	0.73	0.20	0.05	0.15	*
60-64	0.32	0.32	*	0.15	*	0.06
65 or Older	0.17	0.04	0.06	0.04	0.04	*

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.14P Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007**

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
<b>TOTAL</b>	0.6557		0.0213		0.0081	
12	*		*		*	
13	*		*		*	
14	0.5145		0.0530		*	
15	0.8420		0.5038		0.9260	
16	0.2839		0.0608		0.2492	
17	0.4550		0.5798		0.3844	
18	0.9769		0.6218		0.8365	
19	0.6286		0.8499		0.6018	
20	0.6254		0.6316		0.8258	
21	0.4740		0.9813		0.1033	
22	0.8961		0.7965		0.1067	
23	0.2529		0.1002		0.7568	
24	0.8409		0.2147		0.7424	
25	0.2037		0.4223		0.5972	
26-29	0.2414		0.7781		0.0870	
30-34	0.9743		0.3793		0.2595	
35-39	0.9590		0.5642		0.0941	
40-44	0.9833		0.7437		0.1536	
45-49	0.6953		0.0796		0.0908	
50-54	0.1889		0.1508		0.7891	
55-59	0.2695		0.0950		*	
60-64	0.9251		*		*	
65 or Older	0.0960		0.5457		*	

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.15C Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2007 and 2008**

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
<b>TOTAL</b>	638	648	140	129	74	68
12	5	8	4	6	1	3
13	8	9	6	9	5	6
14	11	13	9	11	5	8
15	22	19	19	17	8	11
16	22	22	18	19	9	11
17	23	26	19	21	10	11
18	41	37	29	32	14	18
19	42	42	31	31	13	15
20	46	40	31	31	14	19
21	51	41	30	25	13	14
22	43	46	24	29	9	13
23	54	46	20	20	9	7
24	53	46	23	19	10	11
25	47	45	23	18	11	8
26-29	188	214	64	53	28	36
30-34	207	178	61	48	43	30
35-39	193	210	37	40	25	23
40-44	196	202	18	21	12	*
45-49	207	231	19	19	17	5
50-54	330	301	18	10	10	2
55-59	233	312	*	26	*	7
60-64	123	217	*	*	*	*
65 or Older	71	97	9	4	*	*

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.15D Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2007 and 2008**

<b>Age Category</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	0.26	0.26	0.06	0.05	0.03	0.03
12	0.13	0.21	0.11	0.16	0.03	0.09
13	0.20	0.24	0.16	0.23	0.12	0.15
14	0.27	0.31	0.22	0.26	0.12	0.19
15	0.50	0.42	0.43	0.38	0.19	0.25
16	0.48	0.49	0.39	0.42	0.19	0.26
17	0.52	0.58	0.46	0.48	0.25	0.24
18	0.81	0.72	0.58	0.63	0.30	0.37
19	0.89	0.89	0.67	0.68	0.30	0.34
20	0.99	0.91	0.70	0.74	0.34	0.48
21	1.05	0.91	0.70	0.59	0.32	0.35
22	1.00	0.99	0.61	0.67	0.24	0.32
23	1.17	0.97	0.48	0.48	0.23	0.18
24	1.16	1.05	0.56	0.49	0.25	0.28
25	1.10	1.07	0.60	0.47	0.29	0.20
26-29	0.96	1.12	0.38	0.31	0.17	0.21
30-34	0.97	0.83	0.32	0.26	0.23	0.16
35-39	0.81	0.93	0.18	0.20	0.12	0.11
40-44	0.82	0.85	0.08	0.10	0.06	*
45-49	0.82	0.89	0.09	0.09	0.08	0.02
50-54	1.35	1.23	0.08	0.04	0.05	0.01
55-59	1.17	1.54	*	0.15	*	0.04
60-64	0.90	1.32	*	*	*	*
65 or Older	0.20	0.26	0.02	0.01	*	*

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.15P Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007**

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
<b>TOTAL</b>	0.0901		0.5367		0.5712	
12	0.1856		0.3180		0.0655	
13	0.8080		0.6692		0.6721	
14	0.9951		0.9582		0.0865	
15	0.5929		0.9713		0.1565	
16	0.2006		0.1144		0.0477	
17	0.1898		0.7565		0.8595	
18	0.5235		0.1351		0.6268	
19	0.5533		0.6548		0.6133	
20	0.4890		0.5565		0.0936	
21	0.0518		0.7046		0.5136	
22	0.7307		0.3918		0.1359	
23	0.0002		0.4150		0.2435	
24	0.0089		0.2490		0.6460	
25	0.9763		0.6229		0.1788	
26-29	0.4376		0.0764		0.3859	
30-34	0.5529		0.1902		0.2725	
35-39	0.2338		0.6219		0.8836	
40-44	0.8230		0.9383		*	
45-49	0.0469		0.6679		0.1035	
50-54	0.0784		0.3363		0.4235	
55-59	0.0031		*		*	
60-64	0.0010		*		*	
65 or Older	0.6526		0.6531		*	

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.16C Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2007 and 2008**

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
<b>TOTAL</b>	503	488	106	115	67	71
12	22	24	15	17	11	10
13	30	28	21	21	11	9
14	28	31	19	20	10	12
15	30	29	19	17	10	10
16	29	28	17	18	10	10
17	27	30	16	16	8	8
18	31	32	17	18	7	10
19	34	34	18	15	7	5
20	33	33	14	17	7	10
21	33	32	13	12	7	4
22	35	34	12	9	9	3
23	37	35	9	9	6	3
24	35	31	8	12	2	4
25	39	33	11	6	5	5
26-29	143	162	36	25	23	11
30-34	161	138	26	25	12	11
35-39	164	177	43	34	5	22
40-44	164	168	26	26	17	21
45-49	155	186	18	39	16	38
50-54	232	191	42	5	31	5
55-59	167	187	25	39	25	39
60-64	70	127	*	*	*	2
65 or Older	92	101	*	8	*	8

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.



**Table 1.16D Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2007 and 2008**

<b>Age Category</b>	<b>Lifetime (2007)</b>	<b>Lifetime (2008)</b>	<b>Past Year (2007)</b>	<b>Past Year (2008)</b>	<b>Past Month (2007)</b>	<b>Past Month (2008)</b>
<b>TOTAL</b>	0.20	0.20	0.04	0.05	0.03	0.03
12	0.52	0.58	0.37	0.44	0.27	0.26
13	0.69	0.66	0.49	0.50	0.26	0.23
14	0.64	0.70	0.46	0.47	0.25	0.29
15	0.65	0.63	0.42	0.40	0.22	0.23
16	0.63	0.62	0.39	0.40	0.22	0.22
17	0.63	0.67	0.37	0.37	0.20	0.18
18	0.62	0.61	0.37	0.37	0.14	0.21
19	0.72	0.72	0.41	0.33	0.17	0.11
20	0.76	0.77	0.34	0.41	0.18	0.26
21	0.76	0.75	0.32	0.30	0.18	0.10
22	0.85	0.79	0.32	0.23	0.23	0.06
23	0.86	0.77	0.22	0.22	0.15	0.08
24	0.83	0.77	0.19	0.31	0.04	0.12
25	0.95	0.80	0.28	0.17	0.13	0.13
26-29	0.79	0.89	0.21	0.15	0.14	0.06
30-34	0.78	0.69	0.14	0.13	0.06	0.06
35-39	0.74	0.79	0.20	0.17	0.02	0.11
40-44	0.72	0.74	0.12	0.12	0.08	0.10
45-49	0.65	0.76	0.08	0.17	0.07	0.17
50-54	1.02	0.84	0.19	0.02	0.14	0.02
55-59	0.87	1.01	0.14	0.22	0.14	0.22
60-64	0.52	0.82	*	*	*	0.01
65 or Older	0.25	0.27	*	0.02	*	0.02

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.16P Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007**

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
<b>TOTAL</b>	0.5790		0.7412		0.8406	
12	0.3002		0.0477		0.3937	
13	0.3488		0.9281		0.7006	
14	0.5399		0.7617		0.6469	
15	0.1499		0.4700		0.4809	
16	0.9931		0.8370		0.4877	
17	0.2347		0.2752		0.1569	
18	0.3249		0.9595		0.6294	
19	0.8562		0.4728		0.2710	
20	0.6195		0.1856		0.6973	
21	0.4049		0.7677		0.6942	
22	0.0813		0.1480		0.0292	
23	0.3544		0.8633		0.2764	
24	0.0540		0.1098		0.1440	
25	0.0172		0.3220		0.6018	
26-29	0.9044		0.3817		0.3625	
30-34	0.0281		0.5484		0.8117	
35-39	1.0000		0.4095		0.1816	
40-44	0.7163		0.8255		0.4432	
45-49	0.1536		0.2315		0.2396	
50-54	0.0536		0.1433		0.3571	
55-59	0.0945		0.6083		0.6083	
60-64	0.0158		*		*	
65 or Older	0.8776		*		*	

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.17C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2007 and 2008**

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
<b>TOTAL</b>	696	760	380	353	251	226
12	18	21	14	16	10	11
13	24	19	18	16	11	10
14	26	25	22	20	15	12
15	37	32	33	25	20	16
16	39	36	34	31	19	17
17	39	41	31	32	20	17
18	58	51	49	43	28	23
19	57	57	43	45	27	27
20	64	60	46	47	29	27
21	67	58	47	41	29	24
22	60	62	37	43	25	25
23	59	60	39	36	22	23
24	57	61	35	40	21	26
25	51	57	32	35	20	21
26-29	214	241	116	126	70	72
30-34	209	211	120	109	69	70
35-39	238	232	132	117	81	78
40-44	239	241	113	121	82	79
45-49	248	261	95	101	66	73
50-54	342	356	147	130	111	62
55-59	295	320	136	97	96	63
60-64	194	230	99	78	75	48
65 or Older	162	215	78	82	64	72

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.17D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2007 and 2008**

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
<b>TOTAL</b>	0.28	0.30	0.15	0.14	0.10	0.09
12	0.44	0.52	0.33	0.40	0.25	0.28
13	0.56	0.48	0.43	0.41	0.26	0.24
14	0.61	0.59	0.51	0.46	0.35	0.28
15	0.77	0.67	0.69	0.56	0.43	0.35
16	0.80	0.78	0.70	0.68	0.41	0.40
17	0.82	0.88	0.67	0.69	0.47	0.39
18	0.99	0.92	0.90	0.80	0.56	0.46
19	1.13	1.11	0.91	0.91	0.60	0.60
20	1.22	1.21	0.97	1.07	0.66	0.66
21	1.25	1.12	1.00	0.90	0.68	0.57
22	1.28	1.20	0.88	0.94	0.62	0.59
23	1.23	1.12	0.90	0.80	0.54	0.54
24	1.16	1.22	0.82	0.92	0.51	0.66
25	1.11	1.21	0.80	0.85	0.52	0.55
26-29	1.02	1.17	0.67	0.70	0.42	0.42
30-34	0.95	0.98	0.61	0.56	0.36	0.37
35-39	0.96	0.98	0.60	0.55	0.38	0.38
40-44	0.95	0.97	0.51	0.55	0.38	0.37
45-49	0.91	0.95	0.41	0.44	0.29	0.32
50-54	1.38	1.45	0.66	0.59	0.50	0.28
55-59	1.45	1.55	0.72	0.54	0.52	0.35
60-64	1.34	1.43	0.73	0.51	0.55	0.32
65 or Older	0.45	0.58	0.22	0.22	0.18	0.19

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.17P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007**

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
<b>TOTAL</b>	0.2716		0.0180		0.0300	
12	0.1715		0.1296		0.7772	
13	0.2368		0.8074		0.8253	
14	0.8623		0.6210		0.3972	
15	0.0259		0.0142		0.3757	
16	0.4560		0.8713		0.5928	
17	0.5546		0.5417		0.0105	
18	0.0085		0.0601		0.0270	
19	0.8308		0.9146		0.7334	
20	0.7607		0.9324		0.4163	
21	0.0234		0.0235		0.4626	
22	0.5874		0.1722		0.6320	
23	0.7845		0.7728		0.4750	
24	0.5090		0.1204		0.1999	
25	0.2198		0.4440		0.6590	
26-29	0.3477		0.5770		0.4066	
30-34	0.9115		0.2382		0.6579	
35-39	0.9726		0.3305		0.7000	
40-44	0.6701		0.8732		0.4362	
45-49	0.2884		0.8196		0.8996	
50-54	0.8096		0.3303		0.1155	
55-59	0.8479		0.1490		0.2219	
60-64	0.4160		0.2196		0.2757	
65 or Older	0.0055		0.8775		0.5512	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.18C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2007 and 2008**

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
<b>TOTAL</b>	562	597	316	296	215	196
12	16	20	13	15	10	10
13	20	17	16	15	9	8
14	24	23	19	18	12	10
15	34	30	29	24	17	14
16	37	33	31	28	17	15
17	34	35	27	28	19	16
18	51	48	42	39	23	21
19	53	51	40	38	24	23
20	60	56	41	43	27	24
21	60	54	40	37	24	21
22	56	56	34	38	23	21
23	54	53	33	33	19	20
24	51	54	32	35	17	22
25	48	52	29	32	17	16
26-29	189	210	99	111	65	68
30-34	184	183	104	97	62	61
35-39	198	187	120	100	70	69
40-44	189	193	99	103	67	64
45-49	180	181	84	88	57	62
50-54	218	246	129	98	100	56
55-59	195	223	98	68	76	39
60-64	123	178	72	69	56	39
65 or Older	88	159	61	78	44	72

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.18D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2007 and 2008**

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
<b>TOTAL</b>	0.23	0.24	0.13	0.12	0.09	0.08
12	0.40	0.49	0.32	0.39	0.23	0.26
13	0.48	0.43	0.38	0.37	0.22	0.20
14	0.56	0.54	0.46	0.43	0.28	0.25
15	0.71	0.64	0.63	0.52	0.37	0.32
16	0.76	0.72	0.65	0.62	0.38	0.35
17	0.75	0.77	0.62	0.61	0.44	0.36
18	0.93	0.87	0.81	0.73	0.48	0.42
19	1.07	1.04	0.86	0.81	0.54	0.52
20	1.17	1.17	0.86	1.00	0.61	0.59
21	1.21	1.07	0.91	0.84	0.57	0.50
22	1.25	1.17	0.81	0.85	0.57	0.50
23	1.16	1.05	0.78	0.74	0.48	0.48
24	1.12	1.15	0.77	0.83	0.43	0.56
25	1.07	1.16	0.72	0.79	0.45	0.42
26-29	0.95	1.07	0.57	0.63	0.38	0.39
30-34	0.85	0.89	0.53	0.50	0.33	0.32
35-39	0.85	0.83	0.56	0.48	0.33	0.34
40-44	0.80	0.81	0.45	0.47	0.31	0.30
45-49	0.71	0.74	0.37	0.39	0.25	0.27
50-54	0.96	1.07	0.58	0.44	0.46	0.25
55-59	1.03	1.16	0.52	0.38	0.41	0.22
60-64	0.90	1.15	0.54	0.45	0.42	0.26
65 or Older	0.24	0.43	0.17	0.21	0.12	0.19

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.

**Table 1.18P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2008 Versus 2007**

Age Category	Lifetime (2007)	Lifetime (2008)	Past Year (2007)	Past Year (2008)	Past Month (2007)	Past Month (2008)
<b>TOTAL</b>	0.0606		0.1199		0.1032	
12	0.0926		0.1309		0.7244	
13	0.6478		0.8153		0.8987	
14	0.5499		0.9468		0.5347	
15	0.2283		0.1693		0.7451	
16	0.3753		0.7826		0.3857	
17	0.6641		0.3749		0.0052	
18	0.2563		0.2698		0.1467	
19	0.8850		0.5428		0.6757	
20	0.9068		0.8731		0.5845	
21	0.0565		0.0415		0.4213	
22	0.8816		0.1791		0.9803	
23	0.8256		0.8234		0.6180	
24	0.3169		0.4737		0.2178	
25	0.5283		0.4765		0.6696	
26-29	0.2570		0.9260		0.3910	
30-34	0.8592		0.3902		0.5575	
35-39	0.5854		0.3077		0.7078	
40-44	0.2841		0.9653		0.6873	
45-49	0.4876		0.8485		0.7626	
50-54	0.2623		0.2754		0.2231	
55-59	0.8009		0.2575		0.1668	
60-64	0.1657		0.8584		0.7027	
65 or Older	0.0017		0.3942		0.1557	

\*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007 and 2008.