

Table 7.47A Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|------------------------------|------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|---------------------------------|---------------------------------|
| From One Doctor | 2,138 | 1,997 | 287 | 255 | 514 | 515 | 1,337 | 1,228 |
| From More Than One Doctor | 241 | 263 | 34 | 35 | 54 | 49 | 153 | 178 |
| Wrote Fake Prescription | 18 | 26 | 2 | 2 | 7 | 6 | 9 | 17 |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 40 | 58 | 8 | 10 | 9 ^b | 26 | 23 | 21 |
| From Friend or Relative for Free | 6,449 | 6,211 | 661 ^a | 610 | 2,112 | 2,050 | 3,677 | 3,550 |
| Bought from Friend or Relative | 1,047 | 988 | 124 | 124 | 475 | 469 | 448 | 396 |
| Took from Friend or Relative without Asking | 613 | 603 | 150 | 168 | 215 | 189 | 249 | 245 |
| Bought from Drug Dealer or Other Stranger | 460 | 481 | 66 | 63 | 213 | 235 | 181 | 183 |
| Bought on the Internet | 31 | 43 | 2 | 3 | 7 | 7 | 22 | 33 |
| Some Other Way | 463 | 447 | 105 ^b | 78 | 135 | 128 | 222 | 242 |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.47B Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|------------------------------|------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|---------------------------------|---------------------------------|
| From One Doctor | 18.6 | 18.0 | 19.9 | 18.9 | 13.7 | 14.0 | 21.1 | 20.1 |
| From More Than One Doctor | 2.1 | 2.4 | 2.3 | 2.6 | 1.5 | 1.3 | 2.4 | 2.9 |
| Wrote Fake Prescription | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.3 |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 0.4 | 0.5 | 0.6 | 0.8 | 0.2 ^b | 0.7 | 0.4 | 0.3 |
| From Friend or Relative for Free | 56.1 | 55.9 | 46.0 | 45.3 | 56.4 | 55.8 | 58.2 | 58.3 |
| Bought from Friend or Relative | 9.1 | 8.9 | 8.6 | 9.2 | 12.7 | 12.8 | 7.1 | 6.5 |
| Took from Friend or Relative without Asking | 5.3 | 5.4 | 10.4 ^a | 12.5 | 5.7 | 5.2 | 3.9 | 4.0 |
| Bought from Drug Dealer or Other Stranger | 4.0 | 4.3 | 4.6 | 4.6 | 5.7 | 6.4 | 2.9 | 3.0 |
| Bought on the Internet | 0.3 | 0.4 | 0.1 | 0.2 | 0.2 | 0.2 | 0.4 | 0.5 |
| Some Other Way | 4.0 | 4.0 | 7.3 ^b | 5.8 | 3.6 | 3.5 | 3.5 | 4.0 |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.48A Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Pain Relievers ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 4,675 | 4,489 | 329 | 310 | 1,408 | 1,339 | 2,938 | 2,841 |
| From More Than One Doctor | 178 | 188 | 26 | 27 | 55 | 58 | 97 | 102 |
| Wrote Fake Prescription | 8 | 3 | 0 | 0 | 5 | 3 | 3 | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 34 | 31 | 7 | 5 | 5 | 8 | 22 | 18 |
| From Friend or Relative for Free | 401 ^a | 343 | 70 | 66 | 157 | 148 | 175 | 129 |
| Bought from Friend or Relative | 244 | 220 | 35 | 31 | 128 | 119 | 81 | 70 |
| Took from Friend or Relative without Asking | 67 | 76 | 37 | 33 | 23 | 23 | 8 | 19 |
| Bought from Drug Dealer or Other Stranger | 99 | 89 | 22 | 20 | 45 | 47 | 32 | 23 |
| Bought on the Internet | 2 | 7 | 0 | 0 | 2 | 2 | * | 5 |
| Some Other Way | 72 | 52 | 27 ^a | 18 | 22 | 23 | 23 | 10 |

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.48B Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Pain Relievers¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|------------------------------|------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|---------------------------------|---------------------------------|
| From One Doctor | 80.9 | 81.7 | 59.6 | 60.7 | 76.2 | 75.6 | 86.9 | 88.3 |
| From More Than One Doctor | 3.1 | 3.4 | 4.6 | 5.3 | 3.0 | 3.3 | 2.9 | 3.2 |
| Wrote Fake Prescription | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.2 | 0.1 | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 0.6 | 0.6 | 1.2 | 1.0 | 0.3 | 0.5 | 0.7 | 0.6 |
| From Friend or Relative for Free | 6.9 | 6.2 | 12.7 | 12.9 | 8.5 | 8.3 | 5.2 | 4.0 |
| Bought from Friend or Relative | 4.2 | 4.0 | 6.3 | 6.0 | 6.9 | 6.7 | 2.4 | 2.2 |
| Took from Friend or Relative without Asking | 1.2 | 1.4 | 6.6 | 6.5 | 1.2 | 1.3 | 0.2 | 0.6 |
| Bought from Drug Dealer or Other Stranger | 1.7 | 1.6 | 4.0 | 3.9 | 2.4 | 2.6 | 0.9 | 0.7 |
| Bought on the Internet | 0.0 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | * | 0.1 |
| Some Other Way | 1.2 | 0.9 | 4.8 | 3.6 | 1.2 | 1.3 | 0.7 | 0.3 |

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.49A Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Past Month Nonmedical Use among Past Month Pain Reliever Users¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|------------------------------|------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|---------------------------------|---------------------------------|
| From One Doctor | 1,245 | 1,227 | 138 | 129 | 285 | 277 | 821 | 821 |
| From More Than One Doctor | 244 | 273 | 22 | 21 | 65 | 65 | 156 | 186 |
| Wrote Fake Prescription | 29 | 20 | 3 | 3 | 9 | 9 | 17 | 9 |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 49 | 50 | 8 | 12 | 14 | 18 | 28 | 20 |
| From Friend or Relative for Free | 2,714 | 2,626 | 283 ^a | 247 | 895 | 858 | 1,536 | 1,521 |
| Bought from Friend or Relative | 1,013 | 1,008 | 116 | 101 | 475 | 437 | 422 | 470 |
| Took from Friend or Relative without Asking | 485 | 510 | 97 | 106 | 185 | 172 | 203 | 232 |
| Bought from Drug Dealer or Other Stranger | 507 | 536 | 65 | 65 | 236 | 245 | 205 | 226 |
| Bought on the Internet | 39 | 44 | 4 | 3 | 11 | 14 | 24 | 27 |
| Some Other Way | 285 ^a | 225 | 69 ^b | 48 | 80 | 72 | 137 | 105 |

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.49B Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Past Month Nonmedical Use among Past Month Pain Reliever Users ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 26.2 | 27.4 | 24.9 | 25.5 | 19.5 | 20.0 | 30.0 | 31.7 |
| From More Than One Doctor | 5.1 | 6.1 | 4.1 | 4.2 | 4.4 | 4.7 | 5.7 | 7.2 |
| Wrote Fake Prescription | 0.6 | 0.4 | 0.5 | 0.5 | 0.6 | 0.6 | 0.6 | 0.3 |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 1.0 | 1.1 | 1.5 | 2.4 | 0.9 | 1.3 | 1.0 | 0.8 |
| From Friend or Relative for Free | 57.0 | 58.6 | 51.0 | 48.6 | 61.2 | 61.9 | 56.1 | 58.7 |
| Bought from Friend or Relative | 21.3 | 22.5 | 20.8 | 19.9 | 32.5 | 31.6 | 15.4 | 18.2 |
| Took from Friend or Relative without Asking | 10.2 | 11.4 | 17.4 ^a | 20.9 | 12.7 | 12.4 | 7.4 | 9.0 |
| Bought from Drug Dealer or Other Stranger | 10.7 | 12.0 | 11.7 | 12.8 | 16.2 | 17.7 | 7.5 | 8.7 |
| Bought on the Internet | 0.8 | 1.0 | 0.7 | 0.6 | 0.7 | 1.0 | 0.9 | 1.0 |
| Some Other Way | 6.2 | 5.1 | 13.0 ^a | 9.9 | 5.6 | 5.3 | 5.1 | 4.1 |

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.50A Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Pain Relievers ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 1,930 | 1,832 | 137 | 127 | 601 | 563 | 1,192 | 1,142 |
| From More Than One Doctor | 233 | 238 | 27 | 29 | 79 | 81 | 127 | 128 |
| Wrote Fake Prescription | 40 | 24 | 3 | 3 | 7 ^a | 11 | 30 | 10 |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 57 | 52 | 6 | 9 | 12 | 16 | 40 | 28 |
| From Friend or Relative for Free | 397 | 370 | 55 | 53 | 153 | 141 | 189 | 176 |
| Bought from Friend or Relative | 310 | 322 | 40 | 34 | 145 | 144 | 126 | 144 |
| Took from Friend or Relative without Asking | 99 | 97 | 39 | 29 | 40 | 33 | 20 | 35 |
| Bought from Drug Dealer or Other Stranger | 164 | 176 | 26 | 29 | 90 | 90 | 48 | 58 |
| Bought on the Internet | 25 | 28 | 1 | 3 | 14 | 15 | 10 | 10 |
| Some Other Way | 61 | 68 | 18 | 14 | 25 | 18 | 18 | 36 |

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.50B Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Pain Relievers ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 78.1 | 77.3 | 56.8 | 60.6 | 74.6 | 73.4 | 83.7 | 81.8 |
| From More Than One Doctor | 9.5 | 10.2 | 11.4 | 14.0 | 9.8 | 10.7 | 9.0 | 9.4 |
| Wrote Fake Prescription | 1.6 | 1.0 | 1.1 | 1.6 | 0.8 ^a | 1.4 | 2.2 | 0.8 |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 2.3 | 2.2 | 2.4 | 4.1 | 1.5 | 2.1 | 2.8 | 2.0 |
| From Friend or Relative for Free | 16.2 | 15.8 | 23.2 | 25.6 | 19.2 | 18.6 | 13.4 | 12.8 |
| Bought from Friend or Relative | 12.7 | 13.8 | 16.8 | 16.4 | 18.0 | 18.9 | 8.9 | 10.5 |
| Took from Friend or Relative without Asking | 4.1 | 4.2 | 16.4 | 13.9 | 5.0 | 4.4 | 1.4 | 2.6 |
| Bought from Drug Dealer or Other Stranger | 6.7 | 7.5 | 10.8 | 13.9 | 11.3 | 11.8 | 3.4 | 4.2 |
| Bought on the Internet | 1.0 | 1.2 | 0.5 | 1.4 | 1.7 | 2.0 | 0.7 | 0.7 |
| Some Other Way | 2.6 | 3.0 | 8.2 | 7.1 | 3.2 | 2.5 | 1.3 | 2.7 |

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

**Table 7.51A Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008**

| Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|------------------------------|------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|---------------------------------|---------------------------------|
| From One Doctor | 483 | 433 | 28 | 25 | 112 | 92 | 343 | 316 |
| From More Than One Doctor | 45 | 50 | 4 | 4 | 12 | 17 | 29 | 29 |
| Wrote Fake Prescription | 15 | 6 | 3 | 4 | 3 | 3 | 9 | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 19 | 13 | 2 | 3 | 8 | 10 | 9 | * |
| From Friend or Relative for Free | 2,925 | 2,879 | 170 | 181 | 861 | 861 | 1,893 | 1,837 |
| Bought from Friend or Relative | 548 ^a | 626 | 70 | 73 | 298 | 309 | 180 ^a | 244 |
| Took from Friend or Relative without Asking | 240 | 274 | 38 | 43 | 84 | 87 | 118 | 143 |
| Bought from Drug Dealer or Other Stranger | 307 | 273 | 46 | 43 | 159 | 154 | 102 | 76 |
| Bought on the Internet | 34 | 91 | 2 | 3 | 8 | 4 | 24 | 84 |
| Some Other Way | 158 | 160 | 21 | 18 | 50 | 56 | 87 | 86 |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.51B Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 10.1 | 9.0 | 7.2 | 6.3 | 7.0 | 5.8 | 12.3 | 11.2 |
| From More Than One Doctor | 0.9 | 1.0 | 1.1 | 1.1 | 0.7 | 1.1 | 1.0 | 1.0 |
| Wrote Fake Prescription | 0.3 | 0.1 | 0.8 | 0.9 | 0.2 | 0.2 | 0.3 | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 0.4 | 0.3 | 0.6 | 0.8 | 0.5 | 0.6 | 0.3 | * |
| From Friend or Relative for Free | 61.3 | 59.9 | 44.3 | 45.7 | 54.0 | 54.1 | 67.8 | 65.2 |
| Bought from Friend or Relative | 11.5 ^a | 13.0 | 18.1 | 18.5 | 18.7 | 19.4 | 6.4 ^a | 8.7 |
| Took from Friend or Relative without Asking | 5.0 | 5.7 | 10.0 | 10.9 | 5.2 | 5.5 | 4.2 | 5.1 |
| Bought from Drug Dealer or Other Stranger | 6.4 | 5.7 | 12.0 | 10.7 | 10.0 | 9.7 | 3.6 | 2.7 |
| Bought on the Internet | 0.7 | 1.9 | 0.5 | 0.7 | 0.5 | 0.2 | 0.9 | 3.0 |
| Some Other Way | 3.3 | 3.3 | 5.4 | 4.5 | 3.1 | 3.5 | 3.1 | 3.0 |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.52A Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Tranquilizers ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 2,044 | 1,939 | 60 | 62 | 490 | 491 | 1,495 | 1,385 |
| From More Than One Doctor | 63 | 93 | 5 | 9 | 28 | 23 | 30 | 61 |
| Wrote Fake Prescription | 1 | 1 | * | 1 | 1 | 1 | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 10 | 9 | * | 0 | 3 | 6 | 7 | 3 |
| From Friend or Relative for Free | 234 | 257 | 31 | 28 | 107 | 111 | 95 | 119 |
| Bought from Friend or Relative | 147 | 145 | 16 | 18 | 69 | 81 | 63 | 46 |
| Took from Friend or Relative without Asking | 28 | 33 | 9 ^a | 17 | 12 | 9 | * | * |
| Bought from Drug Dealer or Other Stranger | 70 ^a | 54 | 11 | 7 | 25 | 20 | 34 | 26 |
| Bought on the Internet | 13 | 5 | 0 | 0 | 5 | 4 | 7 | 0 |
| Some Other Way | 30 | 38 | 3 | 4 | 12 | 12 | 15 | 21 |

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.52B Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Tranquilizers¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|------------------------------|------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|---------------------------------|---------------------------------|
| From One Doctor | 77.5 | 75.3 | 44.3 | 42.1 | 65.1 | 64.7 | 85.3 | 83.0 |
| From More Than One Doctor | 2.4 | 3.6 | 3.5 | 6.1 | 3.8 | 3.0 | 1.7 ^a | 3.7 |
| Wrote Fake Prescription | 0.0 | 0.1 | * | 0.3 | 0.1 | 0.1 | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 0.4 | 0.4 | * | 0.3 | 0.4 | 0.8 | 0.4 | 0.2 |
| From Friend or Relative for Free | 8.9 | 10.0 | 23.2 | 18.7 | 14.3 | 14.6 | 5.4 | 7.1 |
| Bought from Friend or Relative | 5.6 | 5.6 | 11.5 | 12.4 | 9.1 | 10.7 | 3.6 | 2.8 |
| Took from Friend or Relative without Asking | 1.1 | 1.3 | 7.0 ^a | 11.8 | 1.6 | 1.2 | * | * |
| Bought from Drug Dealer or Other Stranger | 2.7 | 2.1 | 8.3 ^a | 5.0 | 3.3 | 2.7 | 1.9 | 1.6 |
| Bought on the Internet | 0.5 | 0.2 | 0.2 | 0.2 | 0.7 | 0.6 | 0.4 | 0.0 |
| Some Other Way | 1.1 | 1.5 | 2.0 | 3.0 | 1.6 | 1.6 | 0.9 | 1.3 |

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.53A Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 225 | 217 | 12 | 12 | 58 | 44 | 156 | 161 |
| From More Than One Doctor | 25 | 34 | * | * | 7 | 8 | 17 | 25 |
| Wrote Fake Prescription | 14 | 3 | 2 | * | 3 | 2 | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 13 | 7 | 0 | * | 4 | 6 | * | * |
| From Friend or Relative for Free | 955 | 1,014 | 52 | 63 | 288 | 279 | 615 | 671 |
| Bought from Friend or Relative | 254 | 282 | 26 | 22 | 144 | 136 | 84 ^a | 124 |
| Took from Friend or Relative without Asking | 118 | 125 | 14 | 16 | 49 | 45 | 55 | 64 |
| Bought from Drug Dealer or Other Stranger | 180 | 183 | 19 | 18 | 79 | 76 | 82 | 89 |
| Bought on the Internet | 26 | 46 | 1 | 3 | 6 | 3 | 18 | * |
| Some Other Way | 68 | 55 | 6 | 6 | 19 | 16 | 43 | 33 |

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.53B Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 13.8 | 13.0 | 10.8 | 9.1 | 10.4 | 8.5 | 16.1 | 15.9 |
| From More Than One Doctor | 1.5 | 2.0 | * | * | 1.3 | 1.5 | 1.8 | 2.5 |
| Wrote Fake Prescription | 0.9 | 0.2 | 1.8 | * | 0.5 | 0.4 | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 0.8 | 0.4 | 0.1 | * | 0.8 | 1.1 | * | * |
| From Friend or Relative for Free | 58.5 | 60.9 | 47.4 | 49.5 | 51.9 | 53.7 | 63.5 | 66.0 |
| Bought from Friend or Relative | 15.5 | 17.0 | 23.4 | 17.5 | 26.0 | 26.1 | 8.7 | 12.3 |
| Took from Friend or Relative without Asking | 7.2 | 7.5 | 12.4 | 12.6 | 8.7 | 8.6 | 5.7 | 6.3 |
| Bought from Drug Dealer or Other Stranger | 11.0 | 11.0 | 17.6 | 14.1 | 14.3 | 14.6 | 8.4 | 8.8 |
| Bought on the Internet | 1.6 | 2.8 | 1.1 | 2.1 | 1.1 | 0.7 | 1.9 | * |
| Some Other Way | 4.2 | 3.3 | 5.5 | 4.8 | 3.5 | 3.1 | 4.5 | 3.3 |

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.54A Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Tranquilizers ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 648 | 708 | * | * | 176 | 164 | 453 | 518 |
| From More Than One Doctor | 56 | 85 | * | * | 15 | 21 | * | 62 |
| Wrote Fake Prescription | * | 0 | * | * | * | * | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 3 | 2 | * | * | 3 | * | * | * |
| From Friend or Relative for Free | 106 | 120 | * | * | 56 | 63 | 39 | 45 |
| Bought from Friend or Relative | 69 | 72 | * | * | 32 | 37 | 31 | 24 |
| Took from Friend or Relative without Asking | 18 | 20 | * | * | 7 | 6 | 5 | 8 |
| Bought from Drug Dealer or Other Stranger | 44 | 47 | * | * | 19 | 16 | 19 | 27 |
| Bought on the Internet | 10 | 1 | * | * | 3 | 1 | * | 0 |
| Some Other Way | 6 | 23 | * | * | 2 | 6 | 3 | 16 |

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.54B Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Tranquilizers ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 75.2 | 75.2 | * | * | 69.0 | 63.4 | 80.4 | 82.4 |
| From More Than One Doctor | 6.5 | 9.0 | * | * | 6.0 | 8.1 | * | 9.9 |
| Wrote Fake Prescription | * | 0.0 | * | * | * | * | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 0.4 | 0.2 | * | * | 1.3 | * | * | * |
| From Friend or Relative for Free | 12.4 | 12.8 | * | * | 22.1 | 24.3 | 6.8 | 7.2 |
| Bought from Friend or Relative | 8.0 | 7.7 | * | * | 12.6 | 14.4 | 5.5 | 3.8 |
| Took from Friend or Relative without Asking | 2.1 | 2.1 | * | * | 2.7 | 2.1 | 0.9 | 1.2 |
| Bought from Drug Dealer or Other Stranger | 5.1 | 5.0 | * | * | 7.4 | 6.3 | 3.3 | 4.3 |
| Bought on the Internet | 1.2 | 0.1 | * | * | 1.2 | 0.5 | * | 0.0 |
| Some Other Way | 0.8 | 2.4 | * | * | 0.9 | 2.3 | 0.5 | 2.6 |

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

**Table 7.55A Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008**

| Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 203 | 187 | 40 ^a | 25 | 61 | 53 | 102 | 109 |
| From More Than One Doctor | 17 | 18 | 4 | 4 | 6 | 10 | 7 | * |
| Wrote Fake Prescription | 11 | 6 | 2 | 3 | 2 | 3 | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 20 | 19 | 6 | 4 | 8 | 10 | 5 | 5 |
| From Friend or Relative for Free | 1,161 ^a | 1,018 | 142 | 126 | 477 ^a | 423 | 542 | 468 |
| Bought from Friend or Relative | 290 ^a | 346 | 44 | 40 | 188 | 202 | 58 ^b | 103 |
| Took from Friend or Relative without Asking | 128 | 108 | 35 | 30 | 44 | 34 | 49 | 44 |
| Bought from Drug Dealer or Other Stranger | 125 | 141 | 25 | 24 | 57 | 69 | 43 | 48 |
| Bought on the Internet | 85 | 58 | 3 | 1 | 12 | 14 | 70 ^a | 44 |
| Some Other Way | 85 | 50 | 18 | 14 | 18 | 18 | 49 | 18 |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.55B Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 9.6 | 9.6 | 12.6 | 9.3 | 7.0 | 6.3 | 10.9 | 12.9 |
| From More Than One Doctor | 0.8 | 0.9 | 1.2 | 1.5 | 0.7 | 1.1 | 0.7 | * |
| Wrote Fake Prescription | 0.5 | 0.3 | 0.7 | 1.1 | 0.2 | 0.3 | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 0.9 | 1.0 | 2.0 | 1.3 | 0.9 | 1.2 | 0.6 | 0.6 |
| From Friend or Relative for Free | 54.6 | 52.1 | 44.4 | 46.7 | 54.6 ^a | 50.6 | 58.1 | 55.4 |
| Bought from Friend or Relative | 13.6 ^b | 17.7 | 13.7 | 14.7 | 21.6 | 24.2 | 6.2 ^b | 12.2 |
| Took from Friend or Relative without Asking | 6.0 | 5.5 | 10.9 | 11.1 | 5.0 | 4.1 | 5.3 | 5.2 |
| Bought from Drug Dealer or Other Stranger | 5.9 | 7.2 | 7.9 | 9.0 | 6.6 | 8.3 | 4.6 | 5.7 |
| Bought on the Internet | 4.0 | 3.0 | 0.9 | 0.3 | 1.4 | 1.7 | 7.5 | 5.2 |
| Some Other Way | 4.0 | 2.6 | 5.8 | 5.0 | 2.1 | 2.2 | 5.2 | 2.1 |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.56A Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Stimulants ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 722 | 667 | 72 | 66 | 336 | 296 | * | * |
| From More Than One Doctor | 29 | 28 | 11 | 8 | 14 | 11 | 4 | 8 |
| Wrote Fake Prescription | 0 | 0 | 0 | 0 | * | * | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 2 | 2 | * | 1 | 1 | 1 | * | * |
| From Friend or Relative for Free | 125 | 105 | 16 | 12 | 50 | 46 | * | * |
| Bought from Friend or Relative | 104 | 59 | 8 | 13 | 27 | 27 | * | 19 |
| Took from Friend or Relative without Asking | 8 | 9 | 5 | 6 | 3 | 3 | * | * |
| Bought from Drug Dealer or Other Stranger | 12 | 9 | 5 | 5 | 6 | 4 | 1 | * |
| Bought on the Internet | 14 ^a | 6 | 1 | * | 6 | 3 | 7 | * |
| Some Other Way | 6 | 8 | 1 | 1 | 0 | 2 | 5 | 5 |

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.56B Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Stimulants ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 70.6 | 74.7 | 59.2 | 59.5 | 75.9 | 75.4 | * | * |
| From More Than One Doctor | 2.8 | 3.1 | 8.9 | 7.4 | 3.1 | 2.9 | 0.9 | 2.1 |
| Wrote Fake Prescription | 0.0 | 0.0 | 0.1 | 0.1 | * | * | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 0.2 | 0.2 | * | 0.7 | 0.2 | 0.2 | * | * |
| From Friend or Relative for Free | 12.2 | 11.8 | 13.5 | 10.7 | 11.3 | 11.8 | * | * |
| Bought from Friend or Relative | 10.2 | 6.6 | 6.6 ^a | 11.5 | 6.1 | 6.8 | * | 5.0 |
| Took from Friend or Relative without Asking | 0.8 | 1.0 | 4.4 | 5.3 | 0.7 | 0.7 | * | * |
| Bought from Drug Dealer or Other Stranger | 1.2 | 1.0 | 4.2 | 4.3 | 1.4 | 1.1 | 0.3 | * |
| Bought on the Internet | 1.4 | 0.7 | 1.0 | * | 1.3 | 0.7 | 1.5 | * |
| Some Other Way | 0.6 | 0.9 | 0.9 | 0.6 | 0.1 | 0.5 | 1.0 | 1.4 |

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.57A Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|-------------------|-------------------|------------------------|------------------------|------------------------|------------------------|----------------------|----------------------|
| From One Doctor | 107 | 89 | 23 | 14 | 31 | 27 | * | * |
| From More Than One Doctor | 18 | 21 | 5 | 4 | 10 | 13 | * | * |
| Wrote Fake Prescription | 3 | 5 | 1 | 3 | * | 2 | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 12 | 12 | 3 | 2 | 3 | 5 | * | * |
| From Friend or Relative for Free | 359 | 343 | 42 | 35 | 154 | 128 | * | * |
| Bought from Friend or Relative | 130 ^a | 166 | 13 | 19 | 86 | 83 | 31 ^a | 64 |
| Took from Friend or Relative without Asking | 73 | 53 | 11 | 14 | 27 | 19 | * | * |
| Bought from Drug Dealer or Other Stranger | 55 | 72 | 9 | 10 | 35 | 41 | * | 21 |
| Bought on the Internet | 46 | 34 | * | * | 10 | 9 | * | * |
| Some Other Way | 36 | 15 | 6 | 7 | 6 | 6 | * | * |

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.57B Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|-------------------|-------------------|------------------------|------------------------|------------------------|------------------------|----------------------|----------------------|
| From One Doctor | 15.2 | 13.4 | 22.9 | 16.9 | 11.5 | 10.4 | * | * |
| From More Than One Doctor | 2.6 | 3.2 | 4.8 | 4.3 | 3.8 | 4.9 | * | * |
| Wrote Fake Prescription | 0.4 | 0.8 | 1.3 | 3.4 | * | 0.9 | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 1.6 | 1.8 | 2.9 | 1.8 | 1.2 | 2.0 | * | * |
| From Friend or Relative for Free | 50.7 | 51.4 | 41.7 | 41.2 | 56.5 ^a | 48.9 | * | * |
| Bought from Friend or Relative | 18.3 ^b | 24.9 | 13.5 ^b | 22.2 | 31.4 | 31.8 | 9.1 ^b | 20.1 |
| Took from Friend or Relative without Asking | 10.3 | 8.0 | 10.7 | 16.0 | 10.0 | 7.2 | * | * |
| Bought from Drug Dealer or Other Stranger | 7.7 | 10.7 | 9.4 | 11.9 | 13.0 | 15.7 | * | 6.4 |
| Bought on the Internet | 6.6 | 5.1 | * | * | 3.6 | 3.5 | * | * |
| Some Other Way | 5.1 | 2.3 | 5.7 | 8.2 | 2.2 | 2.2 | * | * |

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.58A Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Stimulants ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | * | * | * | * | 111 | 91 | * | * |
| From More Than One Doctor | 10 | 11 | * | * | 4 | 5 | * | * |
| Wrote Fake Prescription | 3 | * | * | * | * | * | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 1 | * | * | * | * | * | * | * |
| From Friend or Relative for Free | * | * | * | * | 32 | 23 | * | * |
| Bought from Friend or Relative | * | * | * | * | 18 ^a | 9 | * | * |
| Took from Friend or Relative without Asking | 6 | 9 | * | * | 5 | 4 | * | * |
| Bought from Drug Dealer or Other Stranger | 25 | 20 | * | * | 12 ^a | 5 | * | * |
| Bought on the Internet | 9 | 7 | * | * | * | * | * | * |
| Some Other Way | 6 | 9 | * | * | 0 | * | * | * |

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.58B Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Stimulants ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | * | * | * | * | 76.1 | 74.7 | * | * |
| From More Than One Doctor | 3.1 | 3.5 | * | * | 3.0 | 4.0 | * | * |
| Wrote Fake Prescription | 0.8 | * | * | * | * | * | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 0.3 | * | * | * | * | * | * | * |
| From Friend or Relative for Free | * | * | * | * | 21.6 | 18.5 | * | * |
| Bought from Friend or Relative | * | * | * | * | 12.1 | 7.1 | * | * |
| Took from Friend or Relative without Asking | 1.9 | 2.9 | * | * | 3.6 | 3.0 | * | * |
| Bought from Drug Dealer or Other Stranger | 7.6 | 6.2 | * | * | 8.1 ^a | 4.0 | * | * |
| Bought on the Internet | 2.7 | 2.3 | * | * | * | * | * | * |
| Some Other Way | 1.8 | 2.8 | * | * | 0.3 | * | * | * |

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.59A Source Where Methamphetamine was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From Friend or Relative for Free | 716 ^b | 419 | 67 ^b | 46 | 223 ^b | 158 | 425 ^b | 214 |
| Bought from Friend or Relative | 361 | 294 | 24 ^a | 15 | 97 ^b | 68 | 240 | 211 |
| Took from Friend or Relative without Asking | 27 | 19 | 7 | 6 | 9 | 7 | 11 | 6 |
| Bought from Drug Dealer or Other Stranger | 296 ^a | 212 | 21 ^a | 9 | 84 ^b | 57 | 191 | 146 |
| Bought on the Internet | 13 | 6 | * | * | 1 | 0 | * | * |
| Some Other Way | 26 | 27 | 3 | 2 | 13 | 12 | * | 14 |

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.59B Source Where Methamphetamine was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From Friend or Relative for Free | 49.7 ^a | 42.8 | 54.1 | 55.6 | 52.4 | 52.4 | 47.9 ^b | 36.2 |
| Bought from Friend or Relative | 25.1 ^a | 30.1 | 19.3 | 17.6 | 22.7 | 22.5 | 27.0 ^a | 35.7 |
| Took from Friend or Relative without Asking | 1.9 | 1.9 | 5.9 | 6.7 | 2.1 | 2.4 | 1.2 | 1.0 |
| Bought from Drug Dealer or Other Stranger | 20.5 | 21.7 | 16.6 | 10.8 | 19.6 | 18.8 | 21.5 | 24.7 |
| Bought on the Internet | 0.9 | 0.7 | * | * | 0.1 | 0.1 | * | * |
| Some Other Way | 1.8 | 2.8 | 2.5 | 1.8 | 3.0 | 3.9 | * | 2.4 |

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.60A Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Methamphetamine ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From Friend or Relative for Free | 141 | 114 | 19 | * | 54 | 46 | 68 | * |
| Bought from Friend or Relative | 254 ^b | 126 | * | * | 66 ^b | 40 | 163 | * |
| Took from Friend or Relative without Asking | 8 | 4 | * | * | * | * | * | * |
| Bought from Drug Dealer or Other Stranger | 184 ^b | 90 | * | * | 59 ^b | 32 | * | * |
| Bought on the Internet | 0 | * | 0 | * | * | * | * | * |
| Some Other Way | 20 | 14 | * | * | 10 | 8 | 8 | * |

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Methamphetamine include (a) past year users who reported obtaining this drug for their most recent use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's most recent use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.60B Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Methamphetamine ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From Friend or Relative for Free | 23.3 ^b | 32.9 | 30.9 | * | 28.2 | 35.6 | 19.3 | * |
| Bought from Friend or Relative | 41.8 | 36.3 | * | * | 34.0 | 31.1 | 46.1 | * |
| Took from Friend or Relative without Asking | 1.3 | 1.1 | * | * | * | * | * | * |
| Bought from Drug Dealer or Other Stranger | 30.4 | 25.8 | * | * | 30.4 | 24.6 | * | * |
| Bought on the Internet | 0.0 | * | 0.2 | * | * | * | * | * |
| Some Other Way | 3.3 | 4.0 | * | * | 5.4 | 6.5 | 2.1 | * |

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Methamphetamine include (a) past year users who reported obtaining this drug for their most recent use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's most recent use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.61A Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Past Month Nonmedical Use among Past Month Methamphetamine Users¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|------------------------------|------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|---------------------------------|---------------------------------|
| From Friend or Relative for Free | 308 | * | * | * | 68 | * | * | * |
| Bought from Friend or Relative | 242 | * | * | * | 44 ^b | 23 | * | * |
| Took from Friend or Relative without Asking | 22 | * | * | * | * | * | * | * |
| Bought from Drug Dealer or Other Stranger | 184 | * | * | * | 39 | * | * | * |
| Bought on the Internet | 1 | 0 | * | * | 1 | * | * | * |
| Some Other Way | 16 | 20 | * | * | 6 | 4 | 6 | * |

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.61B Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Past Month Nonmedical Use among Past Month Methamphetamine Users¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|--|------------------------------|------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|---------------------------------|---------------------------------|
| From Friend or Relative for Free | 55.3 | * | * | * | 54.7 | * | * | * |
| Bought from Friend or Relative | 43.2 | * | * | * | 35.8 | 29.4 | * | * |
| Took from Friend or Relative without Asking | 3.9 | * | * | * | * | * | * | * |
| Bought from Drug Dealer or Other Stranger | 33.0 | * | * | * | 31.4 | * | * | * |
| Bought on the Internet | 0.2 | 0.0 | * | * | 0.5 | * | * | * |
| Some Other Way | 3.0 | 5.7 | * | * | 5.2 | 5.3 | 1.5 | * |

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.62A Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Methamphetamine ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From Friend or Relative for Free | * | * | * | * | * | * | * | * |
| Bought from Friend or Relative | * | * | * | * | * | * | * | * |
| Took from Friend or Relative without Asking | 6 | 5 | * | * | * | * | * | * |
| Bought from Drug Dealer or Other Stranger | * | * | * | * | * | * | * | * |
| Bought on the Internet | 4 | * | * | * | * | * | * | * |
| Some Other Way | * | * | * | * | * | * | * | * |

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.62B Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Methamphetamine ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From Friend or Relative for Free | * | * | * | * | * | * | * | * |
| Bought from Friend or Relative | * | * | * | * | * | * | * | * |
| Took from Friend or Relative without Asking | 2.2 | 3.1 | * | * | * | * | * | * |
| Bought from Drug Dealer or Other Stranger | * | * | * | * | * | * | * | * |
| Bought on the Internet | 1.6 | * | * | * | * | * | * | * |
| Some Other Way | * | * | * | * | * | * | * | * |

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.63A Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Most Recent Nonmedical Use among Past Year Sedative Users¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|------------------------------|------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|---------------------------------|---------------------------------|
| From One Doctor | 176 ^a | 102 | 17 ^a | 9 | 20 | 16 | 140 | * |
| From More Than One Doctor | 12 | 20 | 3 | 2 | 4 | * | 5 | 17 |
| Wrote Fake Prescription | 6 | 8 | * | * | * | * | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 5 | 8 | * | 1 | 1 | 3 | * | * |
| From Friend or Relative for Free | 472 | 419 | 34 | 41 | 88 | 100 | 350 | * |
| Bought from Friend or Relative | 64 | 40 | 9 | 9 | 23 | 19 | * | 12 |
| Took from Friend or Relative without Asking | 22 | 23 | 5 | 11 | 10 | 6 | 6 | 5 |
| Bought from Drug Dealer or Other Stranger | 25 | 30 | 4 | 6 | 9 | 6 | 13 | 18 |
| Bought on the Internet | 5 | 3 | 1 | * | * | 0 | * | * |
| Some Other Way | 24 | 24 | 6 | 4 | 4 | 4 | 15 | 17 |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.63B Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 21.7 ^a | 15.1 | 20.3 ^a | 10.6 | 12.4 | 10.4 | 24.5 | * |
| From More Than One Doctor | 1.5 | 3.0 | 3.9 | 2.9 | 2.5 | * | 0.8 | 3.9 |
| Wrote Fake Prescription | 0.8 | 1.2 | * | * | * | * | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 0.6 | 1.2 | * | 0.9 | 0.5 | 2.1 | * | * |
| From Friend or Relative for Free | 58.0 | 61.9 | 42.4 | 48.8 | 54.8 | 63.3 | 61.2 | * |
| Bought from Friend or Relative | 7.9 | 5.9 | 10.9 | 10.9 | 14.3 | 11.8 | * | 2.7 |
| Took from Friend or Relative without Asking | 2.6 | 3.3 | 6.6 | 12.8 | 6.5 | 3.9 | 1.0 | 1.3 |
| Bought from Drug Dealer or Other Stranger | 3.1 | 4.4 | 4.7 | 6.8 | 5.5 | 3.9 | 2.2 | 4.1 |
| Bought on the Internet | 0.6 | 0.4 | 1.2 | * | * | 0.1 | * | * |
| Some Other Way | 3.0 | 3.6 | 6.8 | 4.1 | 2.2 | 2.4 | 2.7 | 4.0 |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.64A Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Sedatives ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 379 | 319 | * | * | 59 | 67 | * | * |
| From More Than One Doctor | 11 | 9 | * | * | 2 | 2 | * | * |
| Wrote Fake Prescription | * | * | * | * | * | * | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 0 | 4 | * | * | * | * | * | * |
| From Friend or Relative for Free | 12 | 21 | * | * | 8 | 9 | * | * |
| Bought from Friend or Relative | 6 | 4 | * | * | 3 | * | * | * |
| Took from Friend or Relative without Asking | 2 | 1 | * | * | * | * | * | * |
| Bought from Drug Dealer or Other Stranger | 2 | 1 | * | * | * | * | * | * |
| Bought on the Internet | 0 | 0 | * | * | 0 | 0 | * | * |
| Some Other Way | * | 3 | * | * | 2 | * | * | * |

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.64B Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Sedatives ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | 89.9 | 88.2 | * | * | 79.6 | 79.7 | * | * |
| From More Than One Doctor | 2.6 | 2.4 | * | * | 2.4 | 2.6 | * | * |
| Wrote Fake Prescription | * | * | * | * | * | * | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 0.0 | 1.2 | * | * | * | * | * | * |
| From Friend or Relative for Free | 3.0 | 5.7 | * | * | 10.1 | 10.3 | * | * |
| Bought from Friend or Relative | 1.3 | 1.1 | * | * | 4.5 | * | * | * |
| Took from Friend or Relative without Asking | 0.4 | 0.2 | * | * | * | * | * | * |
| Bought from Drug Dealer or Other Stranger | 0.4 | 0.3 | * | * | * | * | * | * |
| Bought on the Internet | 0.0 | 0.0 | * | * | 0.3 | 0.1 | * | * |
| Some Other Way | * | 0.8 | * | * | 2.3 | * | * | * |

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.65A Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Past Month Nonmedical Use among Past Month Sedative Users ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | * | 42 | * | * | * | * | * | * |
| From More Than One Doctor | 6 | 14 | * | * | * | * | * | * |
| Wrote Fake Prescription | * | * | * | * | * | * | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | * | * | * | * | * | * | * | * |
| From Friend or Relative for Free | * | 181 | * | * | * | * | * | * |
| Bought from Friend or Relative | 26 | 27 | * | * | * | * | * | * |
| Took from Friend or Relative without Asking | 17 | 19 | * | * | * | * | * | * |
| Bought from Drug Dealer or Other Stranger | 21 | 23 | * | * | * | * | * | * |
| Bought on the Internet | * | 1 | * | * | * | * | * | * |
| Some Other Way | 15 | 10 | * | * | * | * | * | * |

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.65B Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source for Past Month Nonmedical Use among Past Month Sedative Users ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | * | 16.4 | * | * | * | * | * | * |
| From More Than One Doctor | 1.7 | 5.5 | * | * | * | * | * | * |
| Wrote Fake Prescription | * | * | * | * | * | * | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | * | * | * | * | * | * | * | * |
| From Friend or Relative for Free | * | 70.1 | * | * | * | * | * | * |
| Bought from Friend or Relative | 7.9 | 10.3 | * | * | * | * | * | * |
| Took from Friend or Relative without Asking | 5.2 | 7.3 | * | * | * | * | * | * |
| Bought from Drug Dealer or Other Stranger | 6.4 | 9.1 | * | * | * | * | * | * |
| Bought on the Internet | * | 0.4 | * | * | * | * | * | * |
| Some Other Way | 4.8 | 3.9 | * | * | * | * | * | * |

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.66A Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Sedatives ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | * | * | * | * | * | * | * | * |
| From More Than One Doctor | * | * | * | * | * | * | * | * |
| Wrote Fake Prescription | * | * | * | * | * | * | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | * | * | * | * | * | * | * | * |
| From Friend or Relative for Free | * | * | * | * | * | * | * | * |
| Bought from Friend or Relative | * | * | * | * | * | * | * | * |
| Took from Friend or Relative without Asking | * | * | * | * | * | * | * | * |
| Bought from Drug Dealer or Other Stranger | * | * | * | * | * | * | * | * |
| Bought on the Internet | * | * | * | * | * | * | * | * |
| Some Other Way | * | * | * | * | * | * | * | * |

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.

Table 7.66B Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2006-2007 and 2007-2008

| Source Where Friend or Relative Obtained Sedatives ¹ | Total (2006-2007) | Total (2007-2008) | Aged 12-17 (2006-2007) | Aged 12-17 (2007-2008) | Aged 18-25 (2006-2007) | Aged 18-25 (2007-2008) | Aged 26+ (2006-2007) | Aged 26+ (2007-2008) |
|---|----------------------|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|-------------------------|-------------------------|
| From One Doctor | * | * | * | * | * | * | * | * |
| From More Than One Doctor | * | * | * | * | * | * | * | * |
| Wrote Fake Prescription | * | * | * | * | * | * | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | * | * | * | * | * | * | * | * |
| From Friend or Relative for Free | * | * | * | * | * | * | * | * |
| Bought from Friend or Relative | * | * | * | * | * | * | * | * |
| Took from Friend or Relative without Asking | * | * | * | * | * | * | * | * |
| Bought from Drug Dealer or Other Stranger | * | * | * | * | * | * | * | * |
| Bought on the Internet | * | * | * | * | * | * | * | * |
| Some Other Way | * | * | * | * | * | * | * | * |

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007-2008 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007-2008 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006, 2007, and 2008.