

Table 6.5C (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Numbers in Thousands, 2008

Age Category	Total	Male	Female
TOTAL	308	185	248
18	21	12	17
19	19	8	18
20	18	13	13
21	19	8	17
22	22	12	17
23	21	12	18
24	18	8	16
25	17	11	13
26-29	87	47	70
30-34	77	51	59
35-39	114	64	91
40-44	98	66	69
45-49	101	71	72
50-54	128	48	118
55-59	105	61	85
60-64	111	68	87
65 or Older	98	70	68

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.5D (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Standard Errors of Percentages, 2008

Age Category	Total	Male	Female
TOTAL	0.14	0.17	0.21
18	0.43	0.48	0.68
19	0.44	0.36	0.83
20	0.46	0.66	0.66
21	0.47	0.39	0.82
22	0.52	0.60	0.84
23	0.51	0.58	0.83
24	0.47	0.44	0.80
25	0.44	0.56	0.66
26-29	0.50	0.54	0.79
30-34	0.40	0.56	0.59
35-39	0.53	0.63	0.83
40-44	0.45	0.62	0.62
45-49	0.45	0.63	0.64
50-54	0.57	0.47	0.99
55-59	0.58	0.68	0.94
60-64	0.71	0.92	1.07
65 or Older	0.26	0.44	0.32

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.6C (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics:
Standard Errors of Numbers in Thousands, 2008

Demographic Characteristic	Total	Aged 18-25	Aged 26-49	Aged 50+
TOTAL	308	54	208	224
GENDER				
Male	185	30	138	126
Female	248	45	161	181
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	288	47	196	207
White	280	41	194	199
Black or African American	78	18	67	34
American Indian or Alaska Native	8	*	7	*
Native Hawaiian or Other Pacific Islander	5	*	*	*
Asian	68	11	26	*
Two or More Races	32	7	28	*
Hispanic or Latino	100	29	63	67
EDUCATION				
< High School	107	22	78	71
High School Graduate	174	33	121	131
Some College	164	33	116	117
College Graduate	152	18	99	114
CURRENT EMPLOYMENT				
Full-Time	197	35	163	98
Part-Time	107	29	70	74
Unemployed	79	19	52	*
Other ¹	217	23	113	180

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.6D (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics: Standard Errors of Percentages, 2008

Demographic Characteristic	Total	Aged 18-25	Aged 26-49	Aged 50+
TOTAL	0.14	0.16	0.21	0.24
GENDER				
Male	0.17	0.18	0.28	0.30
Female	0.21	0.28	0.32	0.37
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.15	0.17	0.24	0.25
White	0.17	0.20	0.28	0.28
Black or African American	0.30	0.38	0.53	0.38
American Indian or Alaska Native	0.89	*	1.75	*
Native Hawaiian or Other Pacific Islander	0.58	*	*	*
Asian	0.68	0.76	0.49	*
Two or More Races	1.25	1.59	3.03	*
Hispanic or Latino	0.33	0.51	0.37	0.87
EDUCATION				
< High School	0.30	0.36	0.56	0.44
High School Graduate	0.25	0.28	0.41	0.43
Some College	0.28	0.30	0.42	0.57
College Graduate	0.24	0.38	0.30	0.44
CURRENT EMPLOYMENT				
Full-Time	0.16	0.23	0.22	0.27
Part-Time	0.35	0.34	0.65	0.65
Unemployed	0.85	0.59	1.17	*
Other ¹	0.33	0.35	0.78	0.40

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

Table 6.7C (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2008

Geographic Characteristic	Total	Aged 18-25	Aged 26-49	Aged 50+
TOTAL	308	54	208	224
GEOGRAPHIC REGION				
Northeast	132	24	76	112
New England	47	7	36	35
Middle Atlantic	123	23	67	106
Midwest	124	24	87	87
East North Central	104	21	76	71
West North Central	70	12	44	51
South	185	32	132	123
South Atlantic	134	24	100	81
East South Central	68	10	49	41
West South Central	110	19	71	77
West	164	30	103	116
Mountain	81	15	50	65
Pacific	141	26	90	96
COUNTY TYPE				
Large Metro	233	46	156	161
Small Metro	174	28	117	124
250K – 1 Mil. Pop.	141	24	93	99
< 250K Pop.	107	15	73	76
Nonmetro	120	18	77	86
Urbanized	90	13	52	72
Less Urbanized	74	12	50	47
Completely Rural	26	3	26	2

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.

**Table 6.7D (REVISED) Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Geographic Characteristics:
Standard Errors of Percentages, 2008**

Geographic Characteristic	Total	Aged 18-25	Aged 26-49	Aged 50+
TOTAL	0.14	0.16	0.21	0.24
GEOGRAPHIC REGION				
Northeast	0.32	0.41	0.42	0.63
New England	0.43	0.48	0.77	0.76
Middle Atlantic	0.40	0.52	0.50	0.81
Midwest	0.25	0.32	0.40	0.43
East North Central	0.30	0.41	0.50	0.50
West North Central	0.47	0.52	0.68	0.82
South	0.23	0.27	0.36	0.36
South Atlantic	0.31	0.41	0.52	0.44
East South Central	0.50	0.56	0.84	0.73
West South Central	0.44	0.48	0.62	0.80
West	0.32	0.38	0.44	0.57
Mountain	0.51	0.62	0.70	1.04
Pacific	0.39	0.48	0.55	0.68
COUNTY TYPE				
Large Metro	0.19	0.25	0.27	0.34
Small Metro	0.25	0.27	0.39	0.42
250K – 1 Mil. Pop.	0.30	0.36	0.47	0.50
< 250K Pop.	0.45	0.41	0.69	0.79
Nonmetro	0.31	0.36	0.50	0.49
Urbanized	0.56	0.59	0.82	1.00
Less Urbanized	0.40	0.50	0.66	0.56
Completely Rural	0.66	0.89	1.66	0.14

*Low precision; no estimate reported.

NOTE: Some 2008 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Estimates for SMI presented in this table may differ from previously published SMI estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2008.