

**Table 1.103A Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 or Older: Numbers in Thousands, 2017 and 2018**

<b>Stimulant Subtype</b>	<b>Any Use in Past Year (2017)</b>	<b>Any Use in Past Year (2018)</b>	<b>Misuse in Past Year among Total Population (2017)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Past Year Any Users (2017)</b>	<b>Misuse in Past Year among Past Year Any Users (2018)</b>
<b>ANY PRESCRIPTION STIMULANT</b>	18,629	18,008	5,839 <sup>a</sup>	5,109	5,839 <sup>a</sup>	5,109
Amphetamine or Methylphenidate Products	14,214	13,749	5,589 <sup>a</sup>	4,835	5,589 <sup>a</sup>	4,835
Amphetamine Products	12,734	12,215	5,280 <sup>a</sup>	4,615	5,280 <sup>a</sup>	4,615
Methylphenidate Products	3,397	3,143	913	747	913	747
Anorectic (Weight-Loss) Stimulants	2,489	2,506	175	147	175	147
Provigil®	515	415	97	125	*	*
Any Other Prescription Stimulant <sup>1</sup>	2,829	2,626	128	84	128	84

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 1.103B Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 or Older: Percentages, 2017 and 2018**

<b>Stimulant Subtype</b>	<b>Any Use in Past Year (2017)</b>	<b>Any Use in Past Year (2018)</b>	<b>Misuse in Past Year among Total Population (2017)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Past Year Any Users (2017)</b>	<b>Misuse in Past Year among Past Year Any Users (2018)</b>
<b>ANY PRESCRIPTION STIMULANT</b>	6.8	6.6	2.1 <sup>a</sup>	1.9	31.3 <sup>a</sup>	28.4
Amphetamine or Methylphenidate Products	5.2	5.0	2.1 <sup>a</sup>	1.8	39.3 <sup>a</sup>	35.2
Amphetamine Products	4.7	4.5	1.9 <sup>a</sup>	1.7	41.5 <sup>a</sup>	37.8
Methylphenidate Products	1.2	1.1	0.3	0.3	26.9	23.8
Anorectic (Weight-Loss) Stimulants	0.9	0.9	0.1	0.1	7.0	5.9
Provigil®	0.2	0.2	0.0	0.0	*	*
Any Other Prescription Stimulant <sup>1</sup>	1.0	1.0	0.0	0.0	4.5	3.2

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 1.104A Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 to 17 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 to 17: Numbers in Thousands, 2017 and 2018**

<b>Stimulant Subtype</b>	<b>Any Use in Past Year (2017)</b>	<b>Any Use in Past Year (2018)</b>	<b>Misuse in Past Year among Total Population (2017)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Past Year Any Users (2017)</b>	<b>Misuse in Past Year among Past Year Any Users (2018)</b>
<b>ANY PRESCRIPTION STIMULANT</b>	1,787	1,752	452	369	452	369
Amphetamine or Methylphenidate Products	1,593	1,572	427	350	427	350
Amphetamine Products	1,243	1,223	401 <sup>a</sup>	320	401	320
Methylphenidate Products	584	561	103	74	103	74
Anorectic (Weight-Loss) Stimulants	84	51	13	7	*	*
Provigil®	15	19	3	3	*	*
Any Other Prescription Stimulant <sup>1</sup>	293	286	16	12	16	12

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 1.104B Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 to 17 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 to 17: Percentages, 2017 and 2018**

<b>Stimulant Subtype</b>	<b>Any Use in Past Year (2017)</b>	<b>Any Use in Past Year (2018)</b>	<b>Misuse in Past Year among Total Population (2017)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Past Year Any Users (2017)</b>	<b>Misuse in Past Year among Past Year Any Users (2018)</b>
<b>ANY PRESCRIPTION STIMULANT</b>	7.2	7.0	1.8	1.5	25.3 <sup>a</sup>	21.1
Amphetamine or Methylphenidate Products	6.4	6.3	1.7	1.4	26.8	22.3
Amphetamine Products	5.0	4.9	1.6	1.3	32.2 <sup>a</sup>	26.1
Methylphenidate Products	2.3	2.3	0.4	0.3	17.6	13.2
Anorectic (Weight-Loss) Stimulants	0.3	0.2	0.1	0.0	*	*
Provigil®	0.1	0.1	0.0	0.0	*	*
Any Other Prescription Stimulant <sup>1</sup>	1.2	1.2	0.1	0.0	5.6	4.1

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 1.105A Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 or Older: Numbers in Thousands, 2017 and 2018**

<b>Stimulant Subtype</b>	<b>Any Use in Past Year (2017)</b>	<b>Any Use in Past Year (2018)</b>	<b>Misuse in Past Year among Total Population (2017)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Past Year Any Users (2017)</b>	<b>Misuse in Past Year among Past Year Any Users (2018)</b>
<b>ANY PRESCRIPTION STIMULANT</b>	16,842	16,256	5,386 <sup>a</sup>	4,739	5,386 <sup>a</sup>	4,739
Amphetamine or Methylphenidate Products	12,621	12,177	5,162 <sup>a</sup>	4,486	5,162 <sup>a</sup>	4,486
Amphetamine Products	11,491	10,992	4,879 <sup>a</sup>	4,295	4,879 <sup>a</sup>	4,295
Methylphenidate Products	2,813	2,583	810	673	810	673
Anorectic (Weight-Loss) Stimulants	2,405	2,455	162	140	162	140
Provigil <sup>®</sup>	500	396	94	121	*	*
Any Other Prescription Stimulant <sup>1</sup>	2,536	2,340	112	73	112	73

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 1.105B Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 or Older: Percentages, 2017 and 2018**

<b>Stimulant Subtype</b>	<b>Any Use in Past Year (2017)</b>	<b>Any Use in Past Year (2018)</b>	<b>Misuse in Past Year among Total Population (2017)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Past Year Any Users (2017)</b>	<b>Misuse in Past Year among Past Year Any Users (2018)</b>
<b>ANY PRESCRIPTION STIMULANT</b>	6.8	6.5	2.2 <sup>a</sup>	1.9	32.0 <sup>a</sup>	29.2
Amphetamine or Methylphenidate Products	5.1	4.9	2.1 <sup>a</sup>	1.8	40.9 <sup>a</sup>	36.8
Amphetamine Products	4.6	4.4	2.0 <sup>a</sup>	1.7	42.5 <sup>a</sup>	39.1
Methylphenidate Products	1.1	1.0	0.3	0.3	28.8	26.1
Anorectic (Weight-Loss) Stimulants	1.0	1.0	0.1	0.1	6.7	5.7
Provigil®	0.2	0.2	0.0	0.0	*	*
Any Other Prescription Stimulant <sup>1</sup>	1.0	0.9	0.0	0.0	4.4	3.1

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 1.106A Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 to 25 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 to 25: Numbers in Thousands, 2017 and 2018**

<b>Stimulant Subtype</b>	<b>Any Use in Past Year (2017)</b>	<b>Any Use in Past Year (2018)</b>	<b>Misuse in Past Year among Total Population (2017)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Past Year Any Users (2017)</b>	<b>Misuse in Past Year among Past Year Any Users (2018)</b>
<b>ANY PRESCRIPTION STIMULANT</b>	5,038 <sup>a</sup>	4,473	2,545 <sup>a</sup>	2,216	2,545 <sup>a</sup>	2,216
Amphetamine or Methylphenidate Products	4,689 <sup>a</sup>	4,120	2,499 <sup>a</sup>	2,175	2,499 <sup>a</sup>	2,175
Amphetamine Products	4,358 <sup>a</sup>	3,740	2,411 <sup>a</sup>	2,092	2,411 <sup>a</sup>	2,092
Methylphenidate Products	1,099	1,058	403	397	403	397
Anorectic (Weight-Loss) Stimulants	190	195	38	16	*	*
Provigil®	43	48	18	26	*	*
Any Other Prescription Stimulant <sup>1</sup>	495	435	28	13	28	13

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 1.106B Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 to 25 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 to 25: Percentages, 2017 and 2018**

<b>Stimulant Subtype</b>	<b>Any Use in Past Year (2017)</b>	<b>Any Use in Past Year (2018)</b>	<b>Misuse in Past Year among Total Population (2017)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Past Year Any Users (2017)</b>	<b>Misuse in Past Year among Past Year Any Users (2018)</b>
<b>ANY PRESCRIPTION STIMULANT</b>	14.7 <sup>a</sup>	13.1	7.4 <sup>a</sup>	6.5	50.5	49.5
Amphetamine or Methylphenidate Products	13.7 <sup>a</sup>	12.1	7.3 <sup>a</sup>	6.4	53.3	52.8
Amphetamine Products	12.7 <sup>a</sup>	11.0	7.0 <sup>a</sup>	6.1	55.3	55.9
Methylphenidate Products	3.2	3.1	1.2	1.2	36.7	37.5
Anorectic (Weight-Loss) Stimulants	0.6	0.6	0.1	0.0	*	*
Provigil®	0.1	0.1	0.1	0.1	*	*
Any Other Prescription Stimulant <sup>1</sup>	1.4	1.3	0.1	0.0	5.7	3.0

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 1.107A Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 26 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 26 or Older: Numbers in Thousands, 2017 and 2018**

<b>Stimulant Subtype</b>	<b>Any Use in Past Year (2017)</b>	<b>Any Use in Past Year (2018)</b>	<b>Misuse in Past Year among Total Population (2017)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Past Year Any Users (2017)</b>	<b>Misuse in Past Year among Past Year Any Users (2018)</b>
<b>ANY PRESCRIPTION STIMULANT</b>	11,805	11,783	2,841	2,524	2,841	2,524
Amphetamine or Methylphenidate Products	7,932	8,057	2,663	2,310	2,663	2,310
Amphetamine Products	7,133	7,252	2,468	2,204	2,468	2,204
Methylphenidate Products	1,714	1,524	406	276	406	276
Anorectic (Weight-Loss) Stimulants	2,215	2,260	124	124	124	124
Provigil®	456	348	76	95	*	*
Any Other Prescription Stimulant <sup>1</sup>	2,041	1,905	84	59	84	59

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 1.107B Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 26 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 26 or Older: Percentages, 2017 and 2018**

<b>Stimulant Subtype</b>	<b>Any Use in Past Year (2017)</b>	<b>Any Use in Past Year (2018)</b>	<b>Misuse in Past Year among Total Population (2017)</b>	<b>Misuse in Past Year among Total Population (2018)</b>	<b>Misuse in Past Year among Past Year Any Users (2017)</b>	<b>Misuse in Past Year among Past Year Any Users (2018)</b>
<b>ANY PRESCRIPTION STIMULANT</b>	5.5	5.5	1.3	1.2	24.1	21.4
Amphetamine or Methylphenidate Products	3.7	3.8	1.3 <sup>a</sup>	1.1	33.6 <sup>a</sup>	28.7
Amphetamine Products	3.4	3.4	1.2	1.0	34.6 <sup>a</sup>	30.4
Methylphenidate Products	0.8	0.7	0.2	0.1	23.7	18.1
Anorectic (Weight-Loss) Stimulants	1.0	1.1	0.1	0.1	5.6	5.5
Provigil®	0.2	0.2	0.0	0.0	*	*
Any Other Prescription Stimulant <sup>1</sup>	1.0	0.9	0.0	0.0	4.1	3.1

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

<sup>a</sup> The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

<sup>1</sup> Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.