

Table 1.103C Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 or Older: Standard Errors of Numbers in Thousands, 2017 and 2018

Stimulant Subtype	Any Use in Past Year (2017)	Any Use in Past Year (2018)	Misuse in Past Year among Total Population (2017)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Past Year Any Users (2017)	Misuse in Past Year among Past Year Any Users (2018)
ANY PRESCRIPTION STIMULANT	369	375	191	178	194	180
Amphetamine or Methylphenidate Products	305	323	186	171	189	173
Amphetamine Products	301	305	179	167	181	169
Methylphenidate Products	146	131	72	57	73	57
Anorectic (Weight-Loss) Stimulants	157	140	33	28	33	28
Provigil®	72	56	26	29	*	*
Any Other Prescription Stimulant ¹	157	148	32	26	32	26

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.103D Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 or Older: Standard Errors of Percentages, 2017 and 2018

Stimulant Subtype	Any Use in Past Year (2017)	Any Use in Past Year (2018)	Misuse in Past Year among Total Population (2017)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Past Year Any Users (2017)	Misuse in Past Year among Past Year Any Users (2018)
ANY PRESCRIPTION STIMULANT	0.14	0.14	0.07	0.06	0.86	0.87
Amphetamine or Methylphenidate Products	0.11	0.12	0.07	0.06	0.99	1.03
Amphetamine Products	0.11	0.11	0.07	0.06	1.06	1.10
Methylphenidate Products	0.05	0.05	0.03	0.02	1.78	1.67
Anorectic (Weight-Loss) Stimulants	0.06	0.05	0.01	0.01	1.33	1.10
Provigil®	0.03	0.02	0.01	0.01	*	*
Any Other Prescription Stimulant ¹	0.06	0.05	0.01	0.01	1.09	0.96

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.103P Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 or Older: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

Stimulant Subtype	Any Use in Past Year (2017)	Any Use in Past Year (2018)	Misuse in Past Year among Total Population (2017)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Past Year Any Users (2017)	Misuse in Past Year among Past Year Any Users (2018)
ANY PRESCRIPTION STIMULANT	0.1563		0.0023		0.0128	
Amphetamine or Methylphenidate Products	0.2046		0.0011		0.0024	
Amphetamine Products	0.1550		0.0025		0.0122	
Methylphenidate Products	0.1628		0.0645		0.2101	
Anorectic (Weight-Loss) Stimulants	0.9920		0.4986		0.4970	
Provigil®	0.2852		0.4954		*	
Any Other Prescription Stimulant ¹	0.3027		0.2745		0.3583	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.104C Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 to 17 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 to 17: Standard Errors of Numbers in Thousands, 2017 and 2018

Stimulant Subtype	Any Use in Past Year (2017)	Any Use in Past Year (2018)	Misuse in Past Year among Total Population (2017)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Past Year Any Users (2017)	Misuse in Past Year among Past Year Any Users (2018)
ANY PRESCRIPTION STIMULANT	62	62	34	27	34	28
Amphetamine or Methylphenidate Products	59	58	33	27	34	27
Amphetamine Products	51	51	33	25	33	26
Methylphenidate Products	36	35	17	13	17	13
Anorectic (Weight-Loss) Stimulants	14	9	6	3	*	*
Provigil®	5	8	3	3	*	*
Any Other Prescription Stimulant ¹	25	26	6	5	6	5

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.104D Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 to 17 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 to 17: Standard Errors of Percentages, 2017 and 2018

Stimulant Subtype	Any Use in Past Year (2017)	Any Use in Past Year (2018)	Misuse in Past Year among Total Population (2017)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Past Year Any Users (2017)	Misuse in Past Year among Past Year Any Users (2018)
ANY PRESCRIPTION STIMULANT	0.25	0.25	0.14	0.11	1.61	1.42
Amphetamine or Methylphenidate Products	0.24	0.23	0.13	0.11	1.76	1.55
Amphetamine Products	0.21	0.21	0.13	0.10	2.12	1.86
Methylphenidate Products	0.15	0.14	0.07	0.05	2.69	2.25
Anorectic (Weight-Loss) Stimulants	0.06	0.04	0.02	0.01	*	*
Provigil®	0.02	0.03	0.01	0.01	*	*
Any Other Prescription Stimulant ¹	0.10	0.10	0.03	0.02	2.08	1.62

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.104P Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 to 17 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 to 17: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

Stimulant Subtype	Any Use in Past Year (2017)	Any Use in Past Year (2018)	Misuse in Past Year among Total Population (2017)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Past Year Any Users (2017)	Misuse in Past Year among Past Year Any Users (2018)
ANY PRESCRIPTION STIMULANT	0.7198		0.0597		0.0496	
Amphetamine or Methylphenidate Products	0.8214		0.0727		0.0522	
Amphetamine Products	0.8027		0.0518		0.0312	
Methylphenidate Products	0.6618		0.1832		0.2061	
Anorectic (Weight-Loss) Stimulants	0.0508		0.3780		*	
Provigil®	0.7039		0.9493		*	
Any Other Prescription Stimulant ¹	0.8694		0.5588		0.5779	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.105C Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 or Older: Standard Errors of Numbers in Thousands, 2017 and 2018

Stimulant Subtype	Any Use in Past Year (2017)	Any Use in Past Year (2018)	Misuse in Past Year among Total Population (2017)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Past Year Any Users (2017)	Misuse in Past Year among Past Year Any Users (2018)
ANY PRESCRIPTION STIMULANT	366	368	190	175	192	177
Amphetamine or Methylphenidate Products	299	316	185	169	187	170
Amphetamine Products	294	299	177	164	179	166
Methylphenidate Products	144	127	71	57	71	58
Anorectic (Weight-Loss) Stimulants	157	139	32	27	32	27
Provigil®	72	56	25	29	*	*
Any Other Prescription Stimulant ¹	155	146	31	25	31	25

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.105D Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 or Older: Standard Errors of Percentages, 2017 and 2018

Stimulant Subtype	Any Use in Past Year (2017)	Any Use in Past Year (2018)	Misuse in Past Year among Total Population (2017)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Past Year Any Users (2017)	Misuse in Past Year among Past Year Any Users (2018)
ANY PRESCRIPTION STIMULANT	0.15	0.15	0.08	0.07	0.95	0.95
Amphetamine or Methylphenidate Products	0.12	0.13	0.07	0.07	1.11	1.14
Amphetamine Products	0.12	0.12	0.07	0.07	1.16	1.21
Methylphenidate Products	0.06	0.05	0.03	0.02	2.06	2.00
Anorectic (Weight-Loss) Stimulants	0.06	0.06	0.01	0.01	1.34	1.11
Provigil®	0.03	0.02	0.01	0.01	*	*
Any Other Prescription Stimulant ¹	0.06	0.06	0.01	0.01	1.19	1.05

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.105P Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 or Older: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

Stimulant Subtype	Any Use in Past Year (2017)	Any Use in Past Year (2018)	Misuse in Past Year among Total Population (2017)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Past Year Any Users (2017)	Misuse in Past Year among Past Year Any Users (2018)
ANY PRESCRIPTION STIMULANT	0.1728		0.0062		0.0318	
Amphetamine or Methylphenidate Products	0.2166		0.0031		0.0082	
Amphetamine Products	0.1620		0.0073		0.0372	
Methylphenidate Products	0.1996		0.1199		0.3471	
Anorectic (Weight-Loss) Stimulants	0.8720		0.5808		0.5483	
Provigil®	0.2662		0.4983		*	
Any Other Prescription Stimulant ¹	0.3108		0.3151		0.4064	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.106C Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 to 25 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 to 25: Standard Errors of Numbers in Thousands, 2017 and 2018

Stimulant Subtype	Any Use in Past Year (2017)	Any Use in Past Year (2018)	Misuse in Past Year among Total Population (2017)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Past Year Any Users (2017)	Misuse in Past Year among Past Year Any Users (2018)
ANY PRESCRIPTION STIMULANT	129	127	99	93	108	104
Amphetamine or Methylphenidate Products	127	124	98	92	107	103
Amphetamine Products	122	118	97	88	105	99
Methylphenidate Products	60	64	38	42	39	43
Anorectic (Weight-Loss) Stimulants	24	24	11	7	*	*
Provigil®	11	12	7	9	*	*
Any Other Prescription Stimulant ¹	38	38	11	5	11	5

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.106D Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 to 25 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 to 25: Standard Errors of Percentages, 2017 and 2018

Stimulant Subtype	Any Use in Past Year (2017)	Any Use in Past Year (2018)	Misuse in Past Year among Total Population (2017)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Past Year Any Users (2017)	Misuse in Past Year among Past Year Any Users (2018)
ANY PRESCRIPTION STIMULANT	0.38	0.37	0.29	0.27	1.34	1.39
Amphetamine or Methylphenidate Products	0.37	0.36	0.29	0.27	1.37	1.47
Amphetamine Products	0.35	0.35	0.28	0.26	1.40	1.54
Methylphenidate Products	0.17	0.19	0.11	0.12	2.71	2.84
Anorectic (Weight-Loss) Stimulants	0.07	0.07	0.03	0.02	*	*
Provigil®	0.03	0.04	0.02	0.03	*	*
Any Other Prescription Stimulant ¹	0.11	0.11	0.03	0.02	2.10	1.20

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.106P Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 to 25 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 to 25: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

Stimulant Subtype	Any Use in Past Year (2017)	Any Use in Past Year (2018)	Misuse in Past Year among Total Population (2017)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Past Year Any Users (2017)	Misuse in Past Year among Past Year Any Users (2018)
ANY PRESCRIPTION STIMULANT	0.0026		0.0154		0.5950	
Amphetamine or Methylphenidate Products	0.0015		0.0161		0.7993	
Amphetamine Products	0.0002		0.0159		0.7678	
Methylphenidate Products	0.7085		0.9589		0.8316	
Anorectic (Weight-Loss) Stimulants	0.8438		0.0785		*	
Provigil®	0.7361		0.4932		*	
Any Other Prescription Stimulant ¹	0.3024		0.2138		0.2708	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.107C Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 26 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 26 or Older: Standard Errors of Numbers in Thousands, 2017 and 2018

Stimulant Subtype	Any Use in Past Year (2017)	Any Use in Past Year (2018)	Misuse in Past Year among Total Population (2017)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Past Year Any Users (2017)	Misuse in Past Year among Past Year Any Users (2018)
ANY PRESCRIPTION STIMULANT	343	329	153	134	154	134
Amphetamine or Methylphenidate Products	269	273	146	126	148	126
Amphetamine Products	262	258	138	125	139	125
Methylphenidate Products	127	109	60	39	60	39
Anorectic (Weight-Loss) Stimulants	153	137	30	27	30	27
Provigil®	71	54	24	27	*	*
Any Other Prescription Stimulant ¹	151	140	29	24	29	24

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.107D Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 26 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 26 or Older: Standard Errors of Percentages, 2017 and 2018

Stimulant Subtype	Any Use in Past Year (2017)	Any Use in Past Year (2018)	Misuse in Past Year among Total Population (2017)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Past Year Any Users (2017)	Misuse in Past Year among Past Year Any Users (2018)
ANY PRESCRIPTION STIMULANT	0.16	0.15	0.07	0.06	1.13	1.05
Amphetamine or Methylphenidate Products	0.13	0.13	0.07	0.06	1.46	1.37
Amphetamine Products	0.12	0.12	0.07	0.06	1.55	1.47
Methylphenidate Products	0.06	0.05	0.03	0.02	2.92	2.48
Anorectic (Weight-Loss) Stimulants	0.07	0.06	0.01	0.01	1.36	1.17
Provigil®	0.03	0.03	0.01	0.01	*	*
Any Other Prescription Stimulant ¹	0.07	0.07	0.01	0.01	1.39	1.26

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 1.107P Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 26 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 26 or Older: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

Stimulant Subtype	Any Use in Past Year (2017)	Any Use in Past Year (2018)	Misuse in Past Year among Total Population (2017)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Past Year Any Users (2017)	Misuse in Past Year among Past Year Any Users (2018)
ANY PRESCRIPTION STIMULANT	0.7850		0.0911		0.0847	
Amphetamine or Methylphenidate Products	0.8913		0.0494		0.0126	
Amphetamine Products	0.8861		0.1197		0.0451	
Methylphenidate Products	0.2280		0.0589		0.1420	
Anorectic (Weight-Loss) Stimulants	0.9037		0.9777		0.9509	
Provigil®	0.2279		0.6122		*	
Any Other Prescription Stimulant ¹	0.4455		0.5095		0.5987	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.