

Table 11.13C Type of Treatment Received in Past Year for Depression among Persons Aged 12 to 17 with Major Depressive Episode in Past Year: Standard Errors of Numbers in Thousands, 2004-2018

| Type of Treatment | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 |
|--|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| SAW OR TALKED TO A HEALTH PROFESSIONAL OR USED PRESCRIPTION MEDICATION | 39 | 38 | 37 | 40 | 35 | 35 | 35 | 35 | 37 | 43 | 50 | 51 | 51 | 54 | 59 |
| Saw or Talked to a Health Professional BUT Did Not Use Prescription Medication | 30 | 28 | 28 | 27 | 27 | 26 | 25 | 26 | 27 | 31 | 36 | 38 | 37 | 37 | 46 |
| Used Prescription Medication BUT Did Not See or Talk to a Health Professional | 11 | 10 | 9 | 11 | 10 | 9 | 12 | 10 | 9 | 17 | 14 | 14 | 11 | 13 | 12 |
| Saw or Talked to a Health Professional AND Used Prescription Medication | 25 | 24 | 21 | 24 | 21 | 20 | 20 | 21 | 23 | 23 | 32 | 30 | 34 | 37 | 33 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown past year treatment for depression data were excluded.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2004-2018.

Table 11.13D Type of Treatment Received in Past Year for Depression among Persons Aged 12 to 17 with Major Depressive Episode in Past Year: Standard Errors of Percentages, 2004-2018

| Type of Treatment | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 |
|--|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| SAW OR TALKED TO A HEALTH PROFESSIONAL OR USED PRESCRIPTION MEDICATION | 1.35 | 1.40 | 1.54 | 1.51 | 1.45 | 1.51 | 1.48 | 1.45 | 1.35 | 1.35 | 1.42 | 1.37 | 1.29 | 1.26 | 1.36 |
| Saw or Talked to a Health Professional BUT Did Not Use Prescription Medication | 1.21 | 1.19 | 1.30 | 1.17 | 1.22 | 1.19 | 1.22 | 1.20 | 1.10 | 1.10 | 1.19 | 1.12 | 1.09 | 1.04 | 1.20 |
| Used Prescription Medication BUT Did Not See or Talk to a Health Professional | 0.47 | 0.45 | 0.47 | 0.52 | 0.48 | 0.47 | 0.63 | 0.50 | 0.40 | 0.65 | 0.50 | 0.47 | 0.36 | 0.41 | 0.36 |
| Saw or Talked to a Health Professional AND Used Prescription Medication | 1.03 | 1.02 | 0.98 | 1.10 | 1.00 | 0.99 | 0.95 | 0.97 | 0.97 | 0.86 | 1.07 | 0.94 | 1.02 | 1.01 | 0.89 |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown past year treatment for depression data were excluded.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2004-2018.

Table 11.13P Type of Treatment Received in Past Year for Depression among Persons Aged 12 to 17 with Major Depressive Episode in Past Year: P Values from Tests of Differences of Percentages, 2018 Versus 2004-2017

| Type of Treatment | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 |
|--|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|------|
| SAW OR TALKED TO A HEALTH PROFESSIONAL OR USED PRESCRIPTION MEDICATION | 0.5172 | 0.0368 | 0.0495 | 0.1440 | 0.0655 | 0.0006 | 0.0406 | 0.0645 | 0.0409 | 0.1152 | 0.8367 | 0.2666 | 0.9120 | 0.8708 | |
| Saw or Talked to a Health Professional BUT Did Not Use Prescription Medication | 0.1375 | 0.0533 | 0.5739 | 0.0512 | 0.3233 | 0.0459 | 0.1571 | 0.2408 | 0.1667 | 0.3757 | 0.2467 | 0.2223 | 0.0558 | 0.1454 | |
| Used Prescription Medication BUT Did Not See or Talk to a Health Professional | 0.0468 | 0.4335 | 0.7833 | 0.4965 | 0.2286 | 0.8166 | 0.2755 | 0.5443 | 0.8093 | 0.0549 | 0.3656 | 0.2623 | 0.5861 | 0.7534 | |
| Saw or Talked to a Health Professional AND Used Prescription Medication | 0.9775 | 0.3427 | 0.0199 | 0.9295 | 0.0527 | 0.0076 | 0.0897 | 0.1559 | 0.1930 | 0.0126 | 0.4831 | 0.6453 | 0.0116 | 0.0695 | |

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Some 2006 to 2010 estimates that are used in the comparisons may differ from previously published estimates due to updates (see Section 3.3.5 in Chapter 3 of the *2018 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

NOTE: Respondents with unknown past year treatment for depression data were excluded.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2004-2018.