

Table 6.6C Source Where Friend or Relative Obtained Pain Relievers among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Pain Relievers from a Friend or Relative for Free in Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2017 and 2018

Source Where Friend or Relative Obtained Pain Relievers	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER	152	153	18	17	151	152	45	43	142	146
Prescription from One Doctor	150	150	18	16	150	149	44	42	142	143
Prescriptions from More Than One Doctor	25	20	*	5	25	19	8	10	23	16
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	7	10	*	2	7	10	7	3	0	*
GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE	44	50	8	6	44	49	19	17	39	46
From Friend or Relative for Free	35	46	5	5	35	46	13	13	33	44
Bought from Friend or Relative	25	14	4	*	24	13	13	7	21	11
Took from Friend or Relative without Asking	8	13	5	1	7	13	*	8	2	10
BOUGHT FROM DRUG DEALER OR OTHER STRANGER	16	19	4	4	16	19	11	8	11	18
SOME OTHER WAY¹	33	35	5	1	33	35	9	7	32	34

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

¹ Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 6.6D Source Where Friend or Relative Obtained Pain Relievers among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Pain Relievers from a Friend or Relative for Free in Past Year, by Age Group: Standard Errors of Percentages, 2017 and 2018

Source Where Friend or Relative Obtained Pain Relievers	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER	1.43	1.72	4.45	4.62	1.49	1.78	2.79	2.77	1.76	2.15
Prescription from One Doctor	1.55	1.79	4.57	5.53	1.62	1.85	2.85	2.90	1.94	2.22
Prescriptions from More Than One Doctor	0.66	0.58	*	3.33	0.70	0.58	1.05	1.46	0.83	0.63
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.20	0.30	*	1.33	0.20	0.31	0.87	0.48	0.02	*
GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE	1.14	1.39	3.68	4.06	1.19	1.43	2.31	2.38	1.38	1.72
From Friend or Relative for Free	0.92	1.31	2.57	3.07	0.96	1.35	1.60	1.82	1.16	1.64
Bought from Friend or Relative	0.65	0.40	1.90	*	0.68	0.40	1.57	0.97	0.75	0.42
Took from Friend or Relative without Asking	0.22	0.37	2.24	0.80	0.20	0.39	*	1.18	0.08	0.38
BOUGHT FROM DRUG DEALER OR OTHER STRANGER	0.44	0.55	2.01	2.33	0.45	0.57	1.40	1.19	0.40	0.69
SOME OTHER WAY¹	0.87	1.00	2.19	0.91	0.91	1.04	1.07	1.04	1.14	1.29

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

¹ Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 6.6P Source Where Friend or Relative Obtained Pain Relievers among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Pain Relievers from a Friend or Relative for Free in Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

Source Where Friend or Relative Obtained Pain Relievers	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER	0.9731		0.2750		0.8036		0.3031		0.4536	
Prescription from One Doctor	0.9373		0.6330		0.9330		0.4858		0.7889	
Prescriptions from More Than One Doctor	0.6084		*		0.3053		0.3430		0.1236	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5918		*		0.6550		0.6429		*	
GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE	0.6594		0.4178		0.7826		0.4549		0.9590	
From Friend or Relative for Free	0.7904		0.7648		0.7446		0.3762		0.5250	
Bought from Friend or Relative	0.0252		*		0.0210		0.2979		0.0424	
Took from Friend or Relative without Asking	0.2743		0.0633		0.0826		*		0.1475	
BOUGHT FROM DRUG DEALER OR OTHER STRANGER	0.4374		0.5669		0.4766		0.4251		0.1593	
SOME OTHER WAY¹	0.8062		0.0779		0.6817		0.8465		0.6424	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

¹ Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 6.8C Source Where Friend or Relative Obtained Stimulants among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Stimulants from a Friend or Relative for Free in Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2017 and 2018

Source Where Friend or Relative Obtained Stimulants	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER	117	113	15	*	116	112	64	57	97	93
Prescription from One Doctor	115	112	14	*	114	111	60	56	96	93
Prescriptions from More Than One Doctor	17	13	4	*	16	13	11	10	12	9
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	3	*	*	*	3	*	3	*	2	*
GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE	43	34	*	*	42	33	22	20	35	26
From Friend or Relative for Free	31	27	*	*	30	26	15	17	26	19
Bought from Friend or Relative	30	20	4	*	28	20	17	8	23	18
Took from Friend or Relative without Asking	3	7	*	*	2	6	2	6	*	*
BOUGHT FROM DRUG DEALER OR OTHER STRANGER	18	15	2	*	18	15	7	7	17	13
SOME OTHER WAY¹	9	9	1	*	9	9	4	4	7	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

¹ Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 6.8D Source Where Friend or Relative Obtained Stimulants among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Stimulants from a Friend or Relative for Free in Past Year, by Age Group: Standard Errors of Percentages, 2017 and 2018

Source Where Friend or Relative Obtained Stimulants	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER	1.66	1.45	5.10	*	1.71	1.49	1.95	1.97	2.70	2.14
Prescription from One Doctor	1.77	1.54	5.54	*	1.82	1.58	2.14	2.11	2.79	2.22
Prescriptions from More Than One Doctor	0.62	0.54	2.84	*	0.64	0.56	0.90	0.94	0.88	0.67
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.12	*	*	*	0.13	*	0.25	*	0.11	*
GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE	1.53	1.32	*	*	1.57	1.35	1.84	1.85	2.42	1.88
From Friend or Relative for Free	1.10	1.04	*	*	1.13	1.07	1.23	1.63	1.84	1.40
Bought from Friend or Relative	1.09	0.79	3.08	*	1.11	0.83	1.43	0.80	1.62	1.30
Took from Friend or Relative without Asking	0.11	0.30	*	*	0.10	0.24	0.21	0.57	*	*
BOUGHT FROM DRUG DEALER OR OTHER STRANGER	0.68	0.62	1.29	*	0.71	0.65	0.63	0.71	1.20	0.97
SOME OTHER WAY¹	0.33	0.37	0.96	*	0.34	0.38	0.37	0.37	0.54	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

¹ Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 6.8P Source Where Friend or Relative Obtained Stimulants among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Stimulants from a Friend or Relative for Free in Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2018 Versus 2017

Source Where Friend or Relative Obtained Stimulants	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER	0.2044		*		0.2287		0.9249		0.1379	
Prescription from One Doctor	0.2060		*		0.2601		0.8177		0.1361	
Prescriptions from More Than One Doctor	0.9660		*		0.8446		0.5965		0.8746	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*		*	
GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE	0.2494		*		0.2817		0.8913		0.2418	
From Friend or Relative for Free	0.6757		*		0.7198		0.2337		0.2250	
Bought from Friend or Relative	0.1304		*		0.1776		0.0533		0.6970	
Took from Friend or Relative without Asking	0.3011		*		0.4855		0.4569		*	
BOUGHT FROM DRUG DEALER OR OTHER STRANGER	0.6757		*		0.6469		0.4623		0.3645	
SOME OTHER WAY¹	0.7950		*		0.8178		0.8755		*	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

¹ Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 6.10C Source Where Friend or Relative Obtained Tranquilizers among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Tranquilizers from a Friend or Relative for Free in Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2017 and 2018

Source Where Friend or Relative Obtained Tranquilizers	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER	119	129	*	*	118	128	42	38	110	122
Prescription from One Doctor	118	128	*	*	117	128	42	37	109	121
Prescriptions from More Than One Doctor	18	10	*	*	18	9	5	6	17	7
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	6	4	*	*	6	4	*	*	*	*
GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE	41	37	*	*	40	36	20	23	34	27
From Friend or Relative for Free	32	27	*	*	32	26	16	14	27	22
Bought from Friend or Relative	26	23	*	*	25	22	11	18	23	13
Took from Friend or Relative without Asking	5	11	*	*	3	11	3	2	1	11
BOUGHT FROM DRUG DEALER OR OTHER STRANGER	19	22	*	*	17	21	12	16	12	14
SOME OTHER WAY¹	8	10	*	*	8	10	4	5	7	9

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

¹ Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 6.10D Source Where Friend or Relative Obtained Tranquilizers among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Tranquilizers from a Friend or Relative for Free in Past Year, by Age Group: Standard Errors of Percentages, 2017 and 2018

Source Where Friend or Relative Obtained Tranquilizers	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER	1.70	1.72	*	*	1.74	1.74	2.89	3.46	2.16	1.80
Prescription from One Doctor	1.83	1.75	*	*	1.88	1.78	2.98	3.47	2.40	1.85
Prescriptions from More Than One Doctor	0.73	0.39	*	*	0.76	0.39	0.73	0.86	1.06	0.43
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.27	0.17	*	*	0.27	0.18	*	*	*	*
GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE	1.56	1.45	*	*	1.61	1.46	2.58	3.03	2.03	1.56
From Friend or Relative for Free	1.29	1.07	*	*	1.33	1.07	2.16	1.99	1.67	1.26
Bought from Friend or Relative	1.03	0.90	*	*	1.06	0.92	1.52	2.53	1.39	0.74
Took from Friend or Relative without Asking	0.19	0.44	*	*	0.15	0.45	0.46	0.29	0.06	0.62
BOUGHT FROM DRUG DEALER OR OTHER STRANGER	0.77	0.87	*	*	0.74	0.89	1.65	2.29	0.77	0.80
SOME OTHER WAY¹	0.31	0.40	*	*	0.33	0.42	0.49	0.75	0.42	0.50

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

¹ Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 6.10P Source Where Friend or Relative Obtained Tranquilizers among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Tranquilizers from a Friend or Relative for Free in Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2018 Versus 2017

Source Where Friend or Relative Obtained Tranquilizers	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER	0.6720		*		0.6221		0.1733		0.2076	
Prescription from One Doctor	0.3640		*		0.3436		0.1049		0.0536	
Prescriptions from More Than One Doctor	0.1603		*		0.1679		0.5364		0.0854	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.6376		*		0.7623		*		*	
GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE	0.4687		*		0.3364		0.4675		0.1461	
From Friend or Relative for Free	0.1955		*		0.1033		0.2069		0.2769	
Bought from Friend or Relative	0.7404		*		0.7640		0.0153		0.1785	
Took from Friend or Relative without Asking	0.6943		*		0.5294		0.2525		0.2642	
BOUGHT FROM DRUG DEALER OR OTHER STRANGER	0.8056		*		0.6291		0.3463		0.9665	
SOME OTHER WAY¹	0.6889		*		0.6611		0.4778		0.8961	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

¹ Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 6.12C Source Where Friend or Relative Obtained Sedatives among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Sedatives from a Friend or Relative for Free in Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2017 and 2018

Source Where Friend or Relative Obtained Sedatives	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER	64	61	*	*	64	60	*	*	*	*
Prescription from One Doctor	64	60	*	*	64	60	*	*	*	*
Prescriptions from More Than One Doctor	*	3	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*	*	*
GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE	16	12	*	*	16	11	*	*	*	*
From Friend or Relative for Free	11	10	*	*	11	10	*	*	*	*
Bought from Friend or Relative	*	3	*	*	*	3	*	*	*	*
Took from Friend or Relative without Asking	1	5	*	*	*	*	*	*	*	*
BOUGHT FROM DRUG DEALER OR OTHER STRANGER	4	*	*	*	*	*	*	*	*	*
SOME OTHER WAY¹	*	*	*	*	*	*	*	*	*	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

¹ Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 6.12D Source Where Friend or Relative Obtained Sedatives among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Sedatives from a Friend or Relative for Free in Past Year, by Age Group: Standard Errors of Percentages, 2017 and 2018

Source Where Friend or Relative Obtained Sedatives	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER	3.30	3.06	*	*	3.33	3.12	*	*	*	*
Prescription from One Doctor	3.48	3.07	*	*	3.51	3.14	*	*	*	*
Prescriptions from More Than One Doctor	*	0.57	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*	*	*
GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE	3.10	2.18	*	*	3.13	2.12	*	*	*	*
From Friend or Relative for Free	2.16	1.89	*	*	2.15	1.91	*	*	*	*
Bought from Friend or Relative	*	0.58	*	*	*	0.61	*	*	*	*
Took from Friend or Relative without Asking	0.15	1.00	*	*	*	*	*	*	*	*
BOUGHT FROM DRUG DEALER OR OTHER STRANGER	0.79	*	*	*	*	*	*	*	*	*
SOME OTHER WAY¹	*	*	*	*	*	*	*	*	*	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

¹ Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

Table 6.12P Source Where Friend or Relative Obtained Sedatives among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Sedatives from a Friend or Relative for Free in Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2018 Versus 2017

Source Where Friend or Relative Obtained Sedatives	Aged 12+ (2017)	Aged 12+ (2018)	Aged 12-17 (2017)	Aged 12-17 (2018)	Aged 18+ (2017)	Aged 18+ (2018)	Aged 18-25 (2017)	Aged 18-25 (2018)	Aged 26+ (2017)	Aged 26+ (2018)
GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER	0.9409		*		0.9449		*		*	
Prescription from One Doctor	0.9607		*		0.8832		*		*	
Prescriptions from More Than One Doctor	*		*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*		*	
GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE	0.5945		*		0.5260		*		*	
From Friend or Relative for Free	0.6853		*		0.7324		*		*	
Bought from Friend or Relative	*		*		*		*		*	
Took from Friend or Relative without Asking	0.2071		*		*		*		*	
BOUGHT FROM DRUG DEALER OR OTHER STRANGER	*		*		*		*		*	
SOME OTHER WAY¹	*		*		*		*		*	

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

¹ Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.