

**Table 8.36A Major Depressive Episode in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Numbers in Thousands, 2017 and 2018**

Age Category	Total (2017)	Total (2018)	Male (2017)	Male (2018)	Female (2017)	Female (2018)
<b>TOTAL</b>	17,297	17,720	6,266	6,272	11,031	11,447
<b>18-25</b>	4,416	4,609	1,618	1,665	2,799	2,943
18-20	1,731	1,818	658	631	1,073	1,187
21-25	2,685	2,790	960	1,034	1,725	1,756
<b>26 or Older</b>	12,880	13,111	4,648	4,607	8,232	8,504
26-49	7,635	7,990	2,744	2,956	4,891	5,034
26-29	1,668 <sup>a</sup>	1,996	564 <sup>a</sup>	797	1,104	1,199
30-34	1,676	1,911	718	696	958 <sup>a</sup>	1,215
35-39	1,432	1,467	457	565	974	902
40-44	1,469	1,319	485	483	984	836
45-49	1,390	1,295	520	415	870	881
50 or Older	5,245	5,121	1,904	1,651	3,341	3,470
50-54	1,297	1,382	521	530	776	852
55-59	1,421	1,281	509	437	911	843
60-64	1,107	971	396	262	710	709
65 or Older	1,420	1,488	478	422	943	1,066

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

<sup>a</sup> The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 8.36B Major Depressive Episode in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Percentages, 2017 and 2018**

Age Category	Total (2017)	Total (2018)	Male (2017)	Male (2018)	Female (2017)	Female (2018)
<b>TOTAL</b>	7.1	7.2	5.3	5.3	8.7	9.0
<b>18-25</b>	13.1	13.8	9.5	9.9	16.6	17.6
18-20	13.6	14.2	9.9	9.3	17.8	19.6
21-25	12.7	13.5	9.3	10.3	15.9	16.5
<b>26 or Older</b>	6.1	6.2	4.6	4.5	7.5	7.7
26-49	7.7	8.0	5.6	6.0	9.7	10.0
26-29	9.4 <sup>a</sup>	11.4	6.6 <sup>a</sup>	9.1	12.1	13.7
30-34	7.8	8.7	6.6	6.4	9.0 <sup>a</sup>	10.9
35-39	6.9	6.8	4.5	5.4	9.2	8.2
40-44	7.5	6.8	5.1	5.0	9.7	8.6
45-49	7.0	6.7	5.2	4.4	8.7	8.9
50 or Older	4.7	4.5	3.7	3.1	5.6	5.8
50-54	6.3	6.8	5.4	5.3	7.0	8.1
55-59	6.5	5.9	4.9	4.4	8.1	7.3
60-64	5.6	4.8	3.9	2.6	7.3	7.1
65 or Older	2.9	2.9	2.2	1.9	3.5	3.8

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

<sup>a</sup> The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 8.37A Major Depressive Episode with Severe Impairment in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Numbers in Thousands, 2017 and 2018**

Age Category	Total (2017)	Total (2018)	Male (2017)	Male (2018)	Female (2017)	Female (2018)
<b>TOTAL</b>	11,007	11,489	3,777	4,013	7,230	7,476
<b>18-25</b>	2,887	2,968	1,020	1,087	1,867	1,881
18-20	1,144	1,183	432	417	712	766
21-25	1,743	1,785	588	670	1,155	1,115
<b>26 or Older</b>	8,119	8,520	2,757	2,926	5,363	5,595
26-49	5,019	5,284	1,718	1,980	3,301	3,304
26-29	1,132	1,326	366	530	766	796
30-34	1,035	1,259	381	480	654	778
35-39	1,011	973	327	366	684	606
40-44	1,003	895	324	345	679	550
45-49	837	831	320	258	517	573
50 or Older	3,101	3,236	1,039	946	2,062	2,291
50-54	727	976	297	287	430 <sup>a</sup>	689
55-59	992	852	307	311	684	541
60-64	621	588	230	128	391	460
65 or Older	761	820	204	220	557	601

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

<sup>a</sup> The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.

**Table 8.37B Major Depressive Episode with Severe Impairment in Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Category: Percentages, 2017 and 2018**

Age Category	Total (2017)	Total (2018)	Male (2017)	Male (2018)	Female (2017)	Female (2018)
<b>TOTAL</b>	4.5	4.7	3.2	3.4	5.7	5.9
<b>18-25</b>	8.5	8.9	6.0	6.5	11.1	11.3
18-20	9.0	9.2	6.5	6.2	11.8	12.6
21-25	8.2	8.6	5.7	6.7	10.7	10.5
<b>26 or Older</b>	3.9	4.0	2.7	2.9	4.9	5.1
26-49	5.0	5.3	3.5	4.0	6.5	6.5
26-29	6.4	7.6	4.3	6.0	8.4	9.1
30-34	4.8	5.7	3.5	4.4	6.2	7.0
35-39	4.9	4.5	3.2	3.5	6.5	5.5
40-44	5.1	4.6	3.4	3.6	6.7	5.7
45-49	4.2	4.3	3.2	2.8	5.2	5.8
50 or Older	2.8	2.9	2.0	1.8	3.5	3.8
50-54	3.5	4.8	3.1	2.9	3.9 <sup>a</sup>	6.6
55-59	4.6	4.0	3.0	3.1	6.1	4.7
60-64	3.1	2.9	2.3	1.3	4.0	4.6
65 or Older	1.5	1.6	0.9	1.0	2.0	2.2

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

<sup>a</sup> The difference between this estimate and the 2018 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2017 and 2018.