


2016-2017 National Survey on Drug Use and Health National Maps of Prevalence Estimates, by State

The 116 national maps presented here show the 2016-2017 National Survey on Drug Use and Health (NSDUH) estimates for 30 substance use and mental health outcomes, by age group, for 50 states and the District of Columbia. Each web link in the table of contents can be used to navigate to all of the maps (for all of the age groups) for the outcome listed in that row. The color of each state on the U.S. maps indicates how the state ranks relative to other states for each measure. States could fall into one of five groups according to their ranking by quintiles. Because 51 states were ranked for each measure, the middle quintile was assigned to 11 states, and the remaining quintiles were assigned 10 states each. In some cases, a "quintile" could have more or fewer states than desired because two (or more) states had the same estimate (to two decimal places). When such ties occurred at the "boundary" between two quintiles, all of the states with the same estimate were conservatively assigned to the lower quintile. Those states with the highest rates for a given measure are in dark red, with the exception of the perceptions of risk measures, for which the lowest perceptions of great risk are in dark red. Those states with the lowest estimates are in dark blue, with the exception of the perceptions of risk measures, for which the highest perceptions of great risk are in dark blue. The upper and lower limits of each quintile shown in the map legend collectively define a continuum and are not necessarily the actual values of a particular state. For example, in [Figure 1a](#), the values on the boundary in the lowest quintile correspond to Texas (7.64 percent) and New Jersey (8.45 percent) and are displayed in the legend. In the next to lowest quintile, South Dakota (8.58 percent) and Mississippi (9.27 percent) are the states with the lowest and highest values; however, in the continuum of the legend, the lower limit was assigned a value of 8.46 percent because the upper limit of the quintile below it is 8.45 percent.


Tables containing specific estimates for these states are available on the 2016-2017 NSDUH web page at <https://www.samhsa.gov/data/>.

Figure 1a *Illicit Drug Use in the Past Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 1b *Illicit Drug Use in the Past Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 1c *Illicit Drug Use in the Past Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 1d *Illicit Drug Use in the Past Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 2a *Marijuana Use in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 2b *Marijuana Use in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 2c *Marijuana Use in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 2d *Marijuana Use in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 3a *Marijuana Use in the Past Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 3b *Marijuana Use in the Past Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 3c *Marijuana Use in the Past Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 3d *Marijuana Use in the Past Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 4a *Perceptions of Great Risk from Smoking Marijuana Once a Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 4b *Perceptions of Great Risk from Smoking Marijuana Once a Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 4c *Perceptions of Great Risk from Smoking Marijuana Once a Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 4d *Perceptions of Great Risk from Smoking Marijuana Once a Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 5a *First Use of Marijuana among Individuals Aged 12 or Older, by State: Average Annual Initiation Estimates (Expressed as Percentages of the At-Risk Population) Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 5b *First Use of Marijuana among Youths Aged 12 to 17, by State: Average Annual Initiation Estimates (Expressed as Percentages of the At-Risk Population) Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 5c *First Use of Marijuana among Adults Aged 18 to 25, by State: Average Annual Initiation Estimates (Expressed as Percentages of the At-Risk Population) Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 5d *First Use of Marijuana among Adults Aged 26 or Older, by State: Average Annual Initiation Estimates (Expressed as Percentages of the At-Risk Population) Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 6a *Illicit Drug Use Other Than Marijuana in the Past Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 6b *Illicit Drug Use Other Than Marijuana in the Past Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 6c *Illicit Drug Use Other Than Marijuana in the Past Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 6d *Illicit Drug Use Other Than Marijuana in the Past Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 7a *Cocaine Use in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 7b *Cocaine Use in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 7c *Cocaine Use in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 7d *Cocaine Use in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 8a *Perceptions of Great Risk from Using Cocaine Once a Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 8b *Perceptions of Great Risk from Using Cocaine Once a Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 8c *Perceptions of Great Risk from Using Cocaine Once a Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 8d *Perceptions of Great Risk from Using Cocaine Once a Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 9a *Heroin Use in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 9b *Heroin Use in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 9c *Heroin Use in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 10a *Perceptions of Great Risk from Trying Heroin Once or Twice among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 10b *Perceptions of Great Risk from Trying Heroin Once or Twice among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 10c *Perceptions of Great Risk from Trying Heroin Once or Twice among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 10d *Perceptions of Great Risk from Trying Heroin Once or Twice among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 11a *Methamphetamine Use in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 11b *Methamphetamine Use in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 11c *Methamphetamine Use in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 11d *Methamphetamine Use in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 12a *Pain Reliever Misuse in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 12b *Pain Reliever Misuse in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 12c *Pain Reliever Misuse in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 12d *Pain Reliever Misuse in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 13a *Alcohol Use in the Past Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 13b *Alcohol Use in the Past Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 13c *Alcohol Use in the Past Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 13d *Alcohol Use in the Past Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 14a *Binge Alcohol Use in the Past Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 14b *Binge Alcohol Use in the Past Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 14c *Binge Alcohol Use in the Past Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 14d *Binge Alcohol Use in the Past Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 15a *Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 15b *Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 15c *Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 15d *Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 16a *Alcohol Use in the Past Month among Individuals Aged 12 to 20, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 16b *Binge Alcohol Use in the Past Month among Individuals Aged 12 to 20, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 17a *Tobacco Product Use in the Past Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 17b *Tobacco Product Use in the Past Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 17c *Tobacco Product Use in the Past Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 17d *Tobacco Product Use in the Past Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 18a *Cigarette Use in the Past Month among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 18b *Cigarette Use in the Past Month among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 18c *Cigarette Use in the Past Month among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 18d *Cigarette Use in the Past Month among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 19a *Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 19b *Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 19c *Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 19d *Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 20a *Illicit Drug Use Disorder in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 20b *Illicit Drug Use Disorder in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 20c *Illicit Drug Use Disorder in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 20d *Illicit Drug Use Disorder in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 21a *Pain Reliever Use Disorder in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 21b *Pain Reliever Use Disorder in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 21c *Pain Reliever Use Disorder in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 21d *Pain Reliever Use Disorder in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 22a *Alcohol Use Disorder in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 22b *Alcohol Use Disorder in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 22c *Alcohol Use Disorder in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 22d *Alcohol Use Disorder in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 23a *Substance Use Disorder in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 23b *Substance Use Disorder in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 23c *Substance Use Disorder in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 23d *Substance Use Disorder in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 24a *Needing But Not Receiving Treatment at a Specialty Facility for Illicit Drug Use in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 24b *Needing But Not Receiving Treatment at a Specialty Facility for Illicit Drug Use in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 24c *Needing But Not Receiving Treatment at a Specialty Facility for Illicit Drug Use in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 24d *Needing But Not Receiving Treatment at a Specialty Facility for Illicit Drug Use in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 25a *Needing But Not Receiving Treatment at a Specialty Facility for Alcohol Use in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 25b *Needing But Not Receiving Treatment at a Specialty Facility for Alcohol Use in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 25c *Needing But Not Receiving Treatment at a Specialty Facility for Alcohol Use in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 25d *Needing But Not Receiving Treatment at a Specialty Facility for Alcohol Use in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 26a *Needing But Not Receiving Treatment at a Specialty Facility for Substance Use in the Past Year among Individuals Aged 12 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 26b *Needing But Not Receiving Treatment at a Specialty Facility for Substance Use in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 26c *Needing But Not Receiving Treatment at a Specialty Facility for Substance Use in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 26d *Needing But Not Receiving Treatment at a Specialty Facility for Substance Use in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 27a *Serious Mental Illness in the Past Year among Adults Aged 18 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 27b *Serious Mental Illness in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 28a *Any Mental Illness in the Past Year among Adults Aged 18 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 28b *Any Mental Illness in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 28c *Any Mental Illness in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 29a *Received Mental Health Services in the Past Year among Adults Aged 18 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 29b *Received Mental Health Services in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 30a *Had Serious Thoughts of Suicide in the Past Year among Adults Aged 18 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 30b *Had Serious Thoughts of Suicide in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 30c *Had Serious Thoughts of Suicide in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 31a *Major Depressive Episode in the Past Year among Adults Aged 18 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 31b *Major Depressive Episode in the Past Year among Youths Aged 12 to 17, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*




Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 31c *Major Depressive Episode in the Past Year among Adults Aged 18 to 25, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*



Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.

Figure 31d *Major Depressive Episode in the Past Year among Adults Aged 26 or Older, by State: Percentages, Annual Averages Based on 2016 and 2017 NSDUHs*



Source: SAMHSA, Center for Behavioral Health Statistics and Quality, NSDUH, 2016 and 2017.