

FLORIDA

Table 29 Selected Drug Use, Perceptions of Great Risk, Past Year Substance Use Disorder and Treatment, and Past Year Mental Health Measures in Florida, by Age Group: Estimated Numbers (in Thousands), Annual Averages Based on 2015-2016 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLICIT DRUGS					
Past Month Illicit Drug Use ^{1,2}	1,696	121	431	1,145	1,575
Past Year Marijuana Use	2,275	179	664	1,432	2,096
Past Month Marijuana Use	1,331	99	391	842	1,232
Perceptions of Great Risk from Smoking Marijuana Once a Month	5,664	374	328	4,961	5,290
First Use of Marijuana ^{3,4}	171	70	75	26	101
Past Month Illicit Drug Use Other Than Marijuana ^{1,2}	595	40	125	430	555
Past Year Cocaine Use	283	6	105	171	277
Perceptions of Great Risk from Using Cocaine Once a Month	13,128	834	1,359	10,935	12,294
Past Year Heroin Use	41	3	11	27	38
Perceptions of Great Risk from Trying Heroin Once or Twice	15,256	956	1,661	12,639	14,300
Past Year Misuse of Pain Relievers ^{2,5}	706	48	126	532	658
ALCOHOL					
Past Month Alcohol Use	9,301	129	1,134	8,038	9,172
Past Month Binge Alcohol Use ⁶	4,410	69	735	3,606	4,341
Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week	8,206	633	801	6,772	7,573
Past Month Alcohol Use ⁷ (Individuals Aged 12 to 20)	443	--	--	--	--
Past Month Binge Alcohol Use ^{6,7} (Individuals Aged 12 to 20)	263	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁸	3,583	62	520	3,000	3,520
Past Month Cigarette Use	2,892	41	401	2,450	2,851
Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day	13,442	1,003	1,414	11,025	12,439
PAST YEAR SUBSTANCE USE DISORDER AND TREATMENT					
Illicit Drug Use Disorder ^{1,2,9}	427	47	136	244	381
Pain Reliever Use Disorder ^{2,5,9}	117	8	20	88	108
Alcohol Use Disorder ⁹	980	31	196	753	949
Substance Use Disorder ^{1,2,9}	1,237	60	291	885	1,176
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,2,10}	394	46	126	222	348
Needing But Not Receiving Treatment for Alcohol Use ¹⁰	910	30	188	691	880
Needing But Not Receiving Treatment for Substance Use ^{1,2,10}	1,168	60	274	834	1,108
PAST YEAR MENTAL HEALTH ISSUES					
Serious Mental Illness ^{4,11}	--	--	100	470	570
Any Mental Illness ^{4,11}	--	--	406	2,363	2,769
Received Mental Health Services ¹²	--	--	199	1,840	2,039
Had Serious Thoughts of Suicide ¹³	--	--	160	375	535
Major Depressive Episode ^{4,14}	--	180	205	774	979

-- Not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach.

NOTE: Estimated numbers appearing as 0 in this table mean that the estimate is greater than 0 but less than 500 because estimated numbers are shown in thousands.

¹ Illicit Drug Use includes the misuse of prescription psychotherapeutics or the use of marijuana, cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine. Illicit Drug Use Other Than Marijuana includes the misuse of prescription psychotherapeutics or the use of cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine.² Misuse of prescription psychotherapeutics is defined as use in any way not directed by a doctor, including use without a prescription of one's own; use in greater amounts, more often, or longer than told; or use in any other way not directed by a doctor. Prescription psychotherapeutics do not include over-the-counter drugs.³ First Use of Marijuana is defined as follows: *Average annual number of marijuana initiates* = $X_1 \div 2$, where X_1 is the number of marijuana initiates in the past 24 months.⁴ For details, see Section B of the "2015-2016 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at <https://www.samhsa.gov/data/>.⁵ Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.⁶ Binge Alcohol Use is defined as drinking five or more drinks (for males) or four or more drinks (for females) on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. In 2015, the definition for females changed from five to four drinks.⁷ Underage drinking is defined for individuals aged 12 to 20; therefore, the "12+" estimate reflects that age group and not individuals aged 12 or older.⁸ Tobacco Products include cigarettes, smokeless tobacco (i.e., snuff, dip, chewing tobacco, or "snus"), cigars, or pipe tobacco.⁹ Substance Use Disorder is defined as meeting criteria for illicit drug or alcohol dependence or abuse. Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).¹⁰ Respondents were classified as needing treatment for a substance use problem if they met the criteria for a substance use disorder as defined in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) or received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).¹¹ Mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, assessed by the Mental Health Surveillance Study (MHSS) *Structured Clinical Interview for the Diagnostic and Statistical Manual of Mental Disorders—Fourth Edition—Research Version—Axis I Disorders* (MHSS-SCID), which is based on the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Any mental illness (AMI) includes adults with any mental disorder regardless of whether their disorder resulted in functional impairment. Serious mental illness (SMI) includes adults with any mental disorder that resulted in serious functional impairment.¹² Mental health services are defined as having received inpatient treatment/counseling or outpatient treatment/counseling or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use.¹³ Respondents were asked, "At any time in the past 12 months, did you seriously think about trying to kill yourself?" If they answered "Yes," they were categorized as having serious thoughts of suicide in the past year.¹⁴ Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when an individual experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from adults aged 18 or older to produce an estimate for those aged 12 or older.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015 and 2016.

FLORIDA

Table 30 Selected Drug Use, Perceptions of Great Risk, Past Year Substance Use Disorder and Treatment, and Past Year Mental Health Measures in Florida, by Age Group: Percentages, Annual Averages Based on 2015-2016 NSDUHs

Measure	12+	12-17	18-25	26+	18+
ILLICIT DRUGS					
Past Month Illicit Drug Use ^{1,2}	9.75	8.58	21.84	8.16	9.85
Past Year Marijuana Use	13.07	12.72	33.66	10.21	13.10
Past Month Marijuana Use	7.65	7.03	19.82	6.00	7.70
Perceptions of Great Risk from Smoking Marijuana Once a Month	32.52	26.58	16.66	35.37	33.04
First Use of Marijuana ^{3,4}	1.66	5.62	7.31	0.32	1.12
Past Month Illicit Drug Use Other Than Marijuana ^{1,2}	3.42	2.86	6.35	3.06	3.47
Past Year Cocaine Use	1.63	0.45	5.35	1.22	1.73
Perceptions of Great Risk from Using Cocaine Once a Month	75.42	59.31	68.92	77.95	76.82
Past Year Heroin Use	0.24	0.22	0.58	0.19	0.24
Perceptions of Great Risk from Trying Heroin Once or Twice	87.67	67.97	84.27	90.09	89.37
Past Year Misuse of Pain Relievers ^{2,5}	4.06	3.42	6.40	3.79	4.11
ALCOHOL					
Past Month Alcohol Use	53.43	9.15	57.50	57.30	57.33
Past Month Binge Alcohol Use ⁶	25.34	4.93	37.28	25.70	27.14
Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week	47.14	45.02	40.64	48.27	47.33
Past Month Alcohol Use ⁷ (Individuals Aged 12 to 20)	20.64	--	--	--	--
Past Month Binge Alcohol Use ^{6,7} (Individuals Aged 12 to 20)	12.26	--	--	--	--
TOBACCO PRODUCTS					
Past Month Tobacco Product Use ⁸	20.59	4.44	26.37	21.39	22.00
Past Month Cigarette Use	16.62	2.93	20.36	17.46	17.82
Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day	77.23	71.33	71.73	78.59	77.74
PAST YEAR SUBSTANCE USE DISORDER AND TREATMENT					
Illicit Drug Use Disorder ^{1,2,9}	2.45	3.31	6.90	1.74	2.38
Pain Reliever Use Disorder ^{2,5,9}	0.67	0.60	1.04	0.62	0.68
Alcohol Use Disorder ⁹	5.63	2.20	9.94	5.36	5.93
Substance Use Disorder ^{1,2,9}	7.11	4.30	14.76	6.31	7.35
Needing But Not Receiving Treatment for Illicit Drug Use ^{1,2,10}	2.27	3.29	6.40	1.58	2.17
Needing But Not Receiving Treatment for Alcohol Use ¹⁰	5.23	2.14	9.56	4.93	5.50
Needing But Not Receiving Treatment for Substance Use ^{1,2,10}	6.71	4.29	13.90	5.94	6.93
PAST YEAR MENTAL HEALTH ISSUES					
Serious Mental Illness ^{4,11}	--	--	5.06	3.35	3.56
Any Mental Illness ^{4,11}	--	--	20.57	16.84	17.30
Received Mental Health Services ¹²	--	--	10.09	13.12	12.75
Had Serious Thoughts of Suicide ¹³	--	--	8.12	2.67	3.34
Major Depressive Episode ^{4,14}	--	12.84	10.40	5.52	6.12

-- Not available.

NOTE: Estimates are based on a survey-weighted hierarchical Bayes estimation approach.

¹ Illicit Drug Use includes the misuse of prescription psychotherapeutics or the use of marijuana, cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine. Illicit Drug Use Other Than Marijuana includes the misuse of prescription psychotherapeutics or the use of cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine.

² Misuse of prescription psychotherapeutics is defined as use in any way not directed by a doctor, including use without a prescription of one's own; use in greater amounts, more often, or longer than told; or use in any other way not directed by a doctor. Prescription psychotherapeutics do not include over-the-counter drugs.

³ First Use of Marijuana is defined as follows: $Average\ annual\ initiation\ of\ marijuana\ (\%) = 100 * \{ [X_1 + (0.5 * X_1 + X_2)] \div 2 \}$, where X_1 is the number of marijuana initiates in the past 24 months and X_2 is the number of individuals who never used marijuana (with the at-risk population defined as $0.5 * X_1 + X_2$). Both of the computation components, X_1 and X_2 , are based on a survey-weighted hierarchical Bayes estimation approach. The age group shown is based on a respondent's age at the time of the interview, not his or her age at first use.

⁴ For details, see Section B of the "2015-2016 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at <https://www.samhsa.gov/data/>.

⁵ Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and 2016.

⁶ Binge Alcohol Use is defined as drinking five or more drinks (for males) or four or more drinks (for females) on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. In 2015, the definition for females changed from five to four drinks.

⁷ Underage drinking is defined for individuals aged 12 to 20; therefore, the "12+" estimate reflects that age group and not individuals aged 12 or older.

⁸ Tobacco Products include cigarettes, smokeless tobacco (i.e., snuff, dip, chewing tobacco, or "snus"), cigars, or pipe tobacco.

⁹ Substance Use Disorder is defined as meeting criteria for illicit drug or alcohol dependence or abuse. Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

¹⁰ Respondents were classified as needing treatment for a substance use problem if they met the criteria for a substance use disorder as defined in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) or received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center).

¹¹ Mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, assessed by the Mental Health Surveillance Study (MHSS) *Structured Clinical Interview for the Diagnostic and Statistical Manual of Mental Disorders—Fourth Edition—Research Version—Axis I Disorders* (MHSS-SCID), which is based on the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Any mental illness (AMI) includes adults with any mental disorder regardless of whether their disorder resulted in functional impairment. Serious mental illness (SMI) includes adults with any mental disorder that resulted in serious functional impairment.

¹² Mental health services are defined as having received inpatient treatment/counseling or outpatient treatment/counseling or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use.

¹³ Respondents were asked, "At any time in the past 12 months, did you seriously think about trying to kill yourself?" If they answered "Yes," they were categorized as having serious thoughts of suicide in the past year.

¹⁴ Major depressive episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when an individual experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. There are minor wording differences in the questions in the adult and adolescent MDE modules. Therefore, data from youths aged 12 to 17 were not combined with data from adults aged 18 or older to produce an estimate for those aged 12 or older.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2015 and 2016.