

Table 10.1 *Illicit Drug Use in the Past Month and Illicit Drug Use Other Than Marijuana in the Past Month in Florida among Individuals Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs*

State/Substate Region	Illicit Drug Use in the Past Month		Illicit Drug Use Other Than Marijuana in the Past Month	
	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Florida	10.37	(9.62 - 11.17)	3.12	(2.76 - 3.53)
Broward (Circuit 17)	9.71	(8.01 - 11.73)	3.04	(2.26 - 4.08)
Central I	10.51	(9.24 - 11.92)	2.90	(2.34 - 3.59)
Circuit 9	10.32	(8.80 - 12.08)	2.75	(2.07 - 3.65)
Circuit 18	10.79	(8.81 - 13.16)	3.12	(2.32 - 4.19)
Central II	10.88	(9.76 - 12.10)	3.11	(2.63 - 3.68)
Circuit 6	11.74	(9.73 - 14.09)	3.07	(2.28 - 4.13)
Circuit 10	9.64	(7.63 - 12.11)	2.99	(2.17 - 4.12)
Circuit 12	11.96	(9.40 - 15.11)	3.17	(2.28 - 4.39)
Circuit 13 (Hillsborough)	12.22	(10.30 - 14.44)	3.52	(2.69 - 4.60)
Circuit 20	8.52	(6.72 - 10.75)	2.75	(2.00 - 3.77)
Northeast	11.04	(9.84 - 12.36)	3.25	(2.71 - 3.88)
Circuit 4	11.31	(9.31 - 13.67)	3.50	(2.59 - 4.70)
Circuit 5	9.25	(7.39 - 11.51)	2.73	(2.00 - 3.73)
Circuit 7	12.61	(10.23 - 15.45)	3.38	(2.47 - 4.62)
Circuit 8 plus Columbia, Dixie, Hamilton, Lafayette, and Suwannee	11.51	(9.10 - 14.46)	3.55	(2.57 - 4.89)
Northwest	12.98	(11.27 - 14.91)	4.01	(3.23 - 4.97)
Circuit 1	12.19	(9.86 - 14.99)	4.12	(3.03 - 5.59)
Circuit 2 plus Madison and Taylor	15.07	(12.13 - 18.58)	3.96	(2.88 - 5.42)
Circuit 14	11.76	(9.16 - 14.97)	3.82	(2.70 - 5.37)
South (Circuits 11 and 16)	7.74	(6.51 - 9.19)	2.77	(2.14 - 3.59)
Southeast	9.86	(8.32 - 11.64)	3.12	(2.46 - 3.94)
Circuit 15 (Palm Beach)	9.94	(8.11 - 12.12)	3.25	(2.43 - 4.34)
Circuit 19	9.68	(7.48 - 12.44)	2.82	(2.00 - 3.95)

NOTE: For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

NOTE: Illicit Drug Use includes the misuse of prescription psychotherapeutics or the use of marijuana, cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine. Illicit Drug Use Other Than Marijuana includes the misuse of prescription psychotherapeutics or the use of cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine. Misuse of prescription psychotherapeutics is defined as use in any way not directed by a doctor, including use without a prescription of one's own; use in greater amounts, more often, or longer than told; or use in any other way not directed by a doctor. Prescription psychotherapeutics do not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.

Table 10.2 Marijuana Use in the Past Year, Marijuana Use in the Past Month, Average Annual Initiation Estimates (Expressed as Percentages of the At-Risk Population) of First Use of Marijuana, and Perceptions of Great Risk from Smoking Marijuana Once a Month in Florida among Individuals Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs

State/Substate Region	Marijuana Use in the Past Year		Marijuana Use in the Past Month		First Use of Marijuana ¹		Perceptions of Great Risk from Smoking Marijuana Once a Month	
	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Florida	13.89	(13.06-14.76)	8.83	(8.16-9.55)	1.71	(1.57-1.87)	30.75	(29.47-32.07)
Broward (Circuit 17)	12.90	(10.81-15.33)	7.57	(6.11-9.34)	1.59	(1.31-1.94)	31.91	(28.35-35.69)
Central I	13.71	(12.18-15.40)	9.19	(7.97-10.58)	1.76	(1.52-2.04)	31.05	(28.58-33.63)
Circuit 9	13.42	(11.57-15.51)	8.99	(7.52-10.72)	1.81	(1.51-2.16)	33.74	(30.64-36.99)
Circuit 18	14.15	(11.71-17.01)	9.49	(7.59-11.80)	1.68	(1.36-2.08)	27.02	(23.41-30.98)
Central II	14.33	(13.05-15.70)	9.30	(8.26-10.46)	1.70	(1.52-1.91)	28.73	(26.76-30.79)
Circuit 6	15.45	(12.95-18.33)	9.61	(7.76-11.84)	1.86	(1.53-2.27)	25.25	(21.86-28.99)
Circuit 10	12.16	(9.90-14.84)	8.30	(6.35-10.78)	1.61	(1.29-2.01)	32.56	(28.43-36.98)
Circuit 12	15.47	(12.44-19.07)	10.41	(7.97-13.48)	1.81	(1.41-2.32)	26.19	(22.14-30.69)
Circuit 13 (Hillsborough)	17.08	(14.74-19.72)	10.98	(9.19-13.08)	2.04	(1.67-2.49)	28.57	(25.21-32.20)
Circuit 20	10.67	(8.54-13.25)	7.06	(5.42-9.15)	1.22	(0.98-1.53)	32.25	(28.21-36.57)
Northeast	15.23	(13.69-16.91)	9.28	(8.15-10.56)	1.92	(1.67-2.20)	26.10	(23.98-28.34)
Circuit 4	16.25	(13.59-19.33)	9.52	(7.61-11.84)	2.03	(1.64-2.51)	27.57	(24.13-31.30)
Circuit 5	12.61	(10.21-15.49)	7.76	(6.03-9.95)	1.44	(1.14-1.83)	28.15	(24.20-32.47)
Circuit 7	17.00	(14.12-20.33)	10.55	(8.30-13.33)	2.28	(1.78-2.91)	23.37	(19.69-27.51)
Circuit 8 plus Columbia, Dixie, Hamilton, Lafayette, and Suwannee	15.49	(12.61-18.87)	9.79	(7.52-12.64)	2.20	(1.71-2.84)	23.37	(19.52-27.72)
Northwest	17.57	(15.48-19.88)	11.28	(9.67-13.12)	2.73	(2.30-3.23)	24.74	(22.20-27.47)
Circuit 1	17.38	(14.25-21.03)	10.59	(8.37-13.30)	3.07	(2.43-3.87)	23.73	(20.04-27.86)
Circuit 2 plus Madison and Taylor	20.36	(16.94-24.26)	13.73	(10.89-17.17)	2.78	(2.16-3.56)	23.75	(19.88-28.10)
Circuit 14	13.91	(10.80-17.73)	9.31	(6.91-12.43)	1.93	(1.47-2.53)	28.65	(23.98-33.83)
South (Circuits 11 and 16)	10.24	(8.78-11.90)	6.72	(5.60-8.05)	1.24	(1.02-1.50)	44.05	(41.01-47.13)
Southeast	13.65	(11.63-15.96)	8.50	(7.06-10.21)	1.58	(1.32-1.89)	29.56	(26.59-32.70)
Circuit 15 (Palm Beach)	13.81	(11.40-16.64)	8.47	(6.75-10.58)	1.47	(1.17-1.83)	29.31	(25.65-33.27)
Circuit 19	13.29	(10.28-17.01)	8.58	(6.43-11.38)	1.84	(1.42-2.38)	30.10	(25.61-34.99)

NOTE: For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹ First Use of Marijuana is defined as follows: $Average\ annual\ initiation\ of\ marijuana\ (\%) = 100 * \{[X_1 \div (0.5 * X_1 + X_2)] \div 2\}$, where X_1 is the number of marijuana initiates in the past 24 months and X_2 is the number of individuals who never used marijuana (with the at-risk population defined as $0.5 * X_1 + X_2$). Both of the computation components, X_1 and X_2 , are based on a survey-weighted hierarchical Bayes estimation approach. The age group is based on a respondent's age at the time of the interview, not his or her age at first use. The Bayesian confidence (credible) intervals presented for Average Annual Estimates of First Use of Marijuana use a simultaneous solution for the First Use of Marijuana and Never Used Marijuana outcomes, which had age group-specific random effects that were allowed to be correlated across the two outcomes. The associated Markov Chain Monte Carlo chains were used to calculate the posterior variance. For details, see Section B of the "2017-2018 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.

Table 10.3 Cocaine Use in the Past Year, Perceptions of Great Risk from Using Cocaine Once a Month, Heroin Use in the Past Year, and Perceptions of Great Risk from Trying Heroin Once or Twice in Florida among Individuals Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs

State/Substate Region	Cocaine Use in the Past Year		Perceptions of Great Risk from Using Cocaine Once a Month		Heroin Use in the Past Year		Perceptions of Great Risk from Trying Heroin Once or Twice	
	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Florida	1.93	(1.66-2.23)	76.26	(75.19 - 77.29)	0.32	(0.22 - 0.47)	89.03	(88.24 - 89.78)
Broward (Circuit 17)	1.42	(0.93-2.17)	75.72	(72.84 - 78.39)	0.26	(0.15 - 0.48)	89.07	(87.04 - 90.81)
Central I	1.98	(1.53-2.55)	75.73	(73.62 - 77.73)	0.30	(0.18 - 0.49)	88.14	(86.73 - 89.43)
Circuit 9	2.00	(1.46-2.73)	76.31	(73.69 - 78.74)	0.28	(0.16 - 0.52)	88.12	(86.31 - 89.72)
Circuit 18	1.94	(1.33-2.84)	74.88	(71.44 - 78.02)	0.31	(0.17 - 0.57)	88.18	(85.90 - 90.14)
Central II	1.82	(1.49-2.23)	76.43	(74.78 - 78.01)	0.31	(0.21 - 0.48)	89.01	(87.87 - 90.06)
Circuit 6	1.88	(1.31-2.68)	75.80	(72.60 - 78.73)	0.34	(0.19 - 0.60)	89.18	(87.08 - 90.97)
Circuit 10	1.72	(1.12-2.62)	79.10	(75.96 - 81.93)	0.30	(0.16 - 0.55)	89.68	(87.66 - 91.41)
Circuit 12	1.61	(1.04-2.49)	77.14	(73.46 - 80.45)	0.30	(0.17 - 0.53)	89.35	(86.97 - 91.34)
Circuit 13 (Hillsborough)	2.34	(1.67-3.28)	74.85	(71.86 - 77.62)	0.32	(0.18 - 0.57)	88.27	(86.27 - 90.01)
Circuit 20	1.41	(0.95-2.10)	76.81	(73.40 - 79.90)	0.31	(0.17 - 0.56)	88.99	(86.75 - 90.89)
Northeast	2.11	(1.66-2.67)	75.16	(73.26 - 76.96)	0.37	(0.24 - 0.56)	88.77	(87.53 - 89.89)
Circuit 4	2.26	(1.52-3.33)	73.58	(70.21 - 76.71)	0.40	(0.23 - 0.70)	88.11	(85.99 - 89.95)
Circuit 5	1.60	(1.07-2.40)	77.24	(74.01 - 80.18)	0.33	(0.18 - 0.60)	89.98	(87.84 - 91.77)
Circuit 7	2.58	(1.74-3.81)	75.61	(72.14 - 78.78)	0.35	(0.20 - 0.61)	88.83	(86.61 - 90.72)
Circuit 8 plus Columbia, Dixie, Hamilton, Lafayette, and Suwannee	2.05	(1.34-3.14)	73.42	(69.61 - 76.91)	0.38	(0.21 - 0.70)	87.56	(85.24 - 89.57)
Northwest	2.62	(1.99-3.45)	73.70	(71.17 - 76.08)	0.45	(0.29 - 0.72)	87.58	(85.95 - 89.05)
Circuit 1	2.68	(1.77-4.04)	73.16	(69.16 - 76.81)	0.50	(0.28 - 0.90)	87.14	(84.61 - 89.30)
Circuit 2 plus Madison and Taylor	3.00	(1.99-4.49)	73.13	(69.47 - 76.51)	0.35	(0.19 - 0.66)	87.62	(85.18 - 89.71)
Circuit 14	1.93	(1.21-3.05)	75.84	(71.86 - 79.42)	0.49	(0.27 - 0.87)	88.60	(86.05 - 90.73)
South (Circuits 11 and 16)	1.59	(1.12-2.25)	79.89	(77.63 - 81.98)	0.25	(0.13 - 0.46)	90.77	(89.35 - 92.03)
Southeast	2.24	(1.67-3.01)	75.77	(73.11 - 78.24)	0.36	(0.22 - 0.60)	89.28	(87.58 - 90.77)
Circuit 15 (Palm Beach)	2.55	(1.80-3.60)	75.18	(71.84 - 78.24)	0.40	(0.22 - 0.72)	89.07	(86.93 - 90.90)
Circuit 19	1.56	(0.98-2.48)	77.06	(73.14 - 80.57)	0.28	(0.15 - 0.52)	89.74	(87.41 - 91.69)

NOTE: For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.

Table 10.4 Methamphetamine Use in the Past Year and Pain Reliever Misuse in the Past Year in Florida among Individuals Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs

State/Substate Region	Methamphetamine Use in the Past Year		Pain Reliever Misuse in the Past Year ¹	
	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Florida	0.42	(0.29 - 0.60)	4.14	(3.69 - 4.65)
Broward (Circuit 17)	0.30	(0.16 - 0.58)	3.76	(2.95 - 4.79)
Central I	0.38	(0.23 - 0.62)	3.90	(3.25 - 4.69)
Circuit 9	0.33	(0.18 - 0.61)	3.63	(2.86 - 4.61)
Circuit 18	0.44	(0.24 - 0.81)	4.31	(3.37 - 5.50)
Central II	0.47	(0.30 - 0.71)	4.26	(3.68 - 4.93)
Circuit 6	0.46	(0.25 - 0.82)	4.38	(3.42 - 5.60)
Circuit 10	0.60	(0.33 - 1.09)	4.30	(3.33 - 5.53)
Circuit 12	0.43	(0.23 - 0.80)	4.26	(3.22 - 5.63)
Circuit 13 (Hillsborough)	0.43	(0.22 - 0.81)	4.11	(3.26 - 5.16)
Circuit 20	0.46	(0.25 - 0.84)	4.27	(3.28 - 5.54)
Northeast	0.48	(0.31 - 0.72)	4.41	(3.75 - 5.20)
Circuit 4	0.42	(0.22 - 0.77)	4.21	(3.28 - 5.39)
Circuit 5	0.51	(0.28 - 0.91)	4.68	(3.58 - 6.09)
Circuit 7	0.45	(0.25 - 0.82)	4.25	(3.27 - 5.52)
Circuit 8 plus Columbia, Dixie, Hamilton, Lafayette, and Suwannee	0.58	(0.31 - 1.08)	4.58	(3.54 - 5.89)
Northwest	0.56	(0.34 - 0.91)	4.78	(4.00 - 5.71)
Circuit 1	0.66	(0.35 - 1.26)	4.91	(3.77 - 6.37)
Circuit 2 plus Madison and Taylor	0.36	(0.19 - 0.70)	4.54	(3.54 - 5.80)
Circuit 14	0.61	(0.32 - 1.16)	4.84	(3.67 - 6.35)
South (Circuits 11 and 16)	0.31	(0.16 - 0.59)	3.56	(2.80 - 4.52)
Southeast	0.37	(0.22 - 0.61)	4.32	(3.53 - 5.29)
Circuit 15 (Palm Beach)	0.33	(0.17 - 0.61)	4.48	(3.50 - 5.71)
Circuit 19	0.45	(0.24 - 0.85)	3.98	(3.02 - 5.24)

NOTE: For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹ Misuse of prescription psychotherapeutics is defined as use in any way not directed by a doctor, including use without a prescription of one's own; use in greater amounts, more often, or longer than told; or use in any other way not directed by a doctor. Prescription psychotherapeutics do not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.

Table 10.5 Alcohol Use in the Past Month, Binge Alcohol Use in the Past Month, and Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week in Florida among Individuals Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs

State/Substate Region	Alcohol Use in the Past Month		Binge Alcohol Use in the Past Month ¹		Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week	
	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Florida	51.53	(50.18 - 52.87)	23.68	(22.55 - 24.84)	47.86	(46.54 - 49.18)
Broward (Circuit 17)	49.27	(45.36 - 53.19)	21.08	(18.33 - 24.12)	52.48	(48.98 - 55.96)
Central I	51.19	(48.50 - 53.88)	24.75	(22.54 - 27.10)	47.39	(44.90 - 49.90)
Circuit 9	50.52	(47.24 - 53.79)	26.64	(23.86 - 29.61)	49.79	(46.67 - 52.90)
Circuit 18	52.21	(47.93 - 56.46)	21.93	(18.77 - 25.45)	43.82	(40.18 - 47.51)
Central II	53.25	(51.16 - 55.33)	23.26	(21.57 - 25.05)	45.72	(43.78 - 47.68)
Circuit 6	55.09	(51.01 - 59.10)	22.27	(19.34 - 25.52)	44.95	(41.39 - 48.57)
Circuit 10	45.09	(40.33 - 49.95)	24.40	(20.88 - 28.30)	48.52	(44.43 - 52.62)
Circuit 12	56.88	(51.86 - 61.76)	23.02	(19.29 - 27.22)	45.31	(40.96 - 49.73)
Circuit 13 (Hillsborough)	53.08	(49.33 - 56.80)	23.43	(20.65 - 26.46)	45.69	(42.41 - 49.01)
Circuit 20	53.85	(49.39 - 58.25)	23.72	(20.36 - 27.43)	45.25	(41.35 - 49.20)
Northeast	50.93	(48.43 - 53.41)	23.63	(21.74 - 25.62)	44.49	(42.30 - 46.70)
Circuit 4	50.96	(46.72 - 55.19)	24.94	(21.60 - 28.60)	44.59	(40.96 - 48.27)
Circuit 5	46.81	(42.12 - 51.55)	21.25	(18.05 - 24.86)	46.16	(42.20 - 50.17)
Circuit 7	58.02	(53.35 - 62.56)	24.62	(21.22 - 28.36)	42.39	(38.60 - 46.27)
Circuit 8 plus Columbia, Dixie, Hamilton, Lafayette, and Suwannee	47.38	(42.40 - 52.41)	24.06	(20.37 - 28.19)	44.37	(40.18 - 48.64)
Northwest	51.45	(48.18 - 54.71)	25.53	(23.09 - 28.13)	43.79	(41.13 - 46.48)
Circuit 1	50.32	(45.34 - 55.29)	24.33	(20.84 - 28.19)	44.29	(40.31 - 48.34)
Circuit 2 plus Madison and Taylor	56.84	(51.79 - 61.75)	29.51	(25.47 - 33.90)	43.25	(39.31 - 47.27)
Circuit 14	46.20	(40.36 - 52.14)	22.50	(18.67 - 26.85)	43.39	(39.00 - 47.89)
South (Circuits 11 and 16)	49.52	(46.31 - 52.73)	25.04	(22.44 - 27.82)	56.53	(53.69 - 59.32)
Southeast	53.15	(49.69 - 56.57)	22.77	(20.17 - 25.60)	47.35	(44.43 - 50.28)
Circuit 15 (Palm Beach)	54.15	(49.83 - 58.40)	23.64	(20.38 - 27.25)	47.85	(44.11 - 51.61)
Circuit 19	50.94	(45.58 - 56.29)	20.85	(17.27 - 24.96)	46.25	(42.00 - 50.54)

NOTE: For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹ Binge Alcohol Use is defined as drinking five or more drinks (for males) or four or more drinks (for females) on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.

Table 10.6 Alcohol Use in the Past Month and Binge Alcohol Use in the Past Month among Individuals Aged 12 to 20 in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs

State/Substate Region	Alcohol Use in the Past Month among Individuals Aged 12 to 20		Binge Alcohol Use in the Past Month among Individuals Aged 12 to 20 ¹	
	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Florida	19.57	(18.27 - 20.93)	11.10	(10.11 - 12.18)
Broward (Circuit 17)	16.05	(13.16 - 19.44)	8.04	(6.22 - 10.33)
Central I	19.34	(17.12 - 21.79)	12.06	(10.25 - 14.15)
Circuit 9	19.50	(16.67 - 22.68)	12.51	(10.23 - 15.23)
Circuit 18	19.07	(15.58 - 23.14)	11.26	(8.71 - 14.44)
Central II	19.61	(17.74 - 21.63)	10.47	(9.17 - 11.93)
Circuit 6	20.13	(16.64 - 24.13)	10.17	(7.93 - 12.95)
Circuit 10	17.05	(13.42 - 21.42)	11.45	(8.74 - 14.87)
Circuit 12	22.04	(18.02 - 26.67)	11.57	(8.76 - 15.12)
Circuit 13 (Hillsborough)	20.20	(17.01 - 23.81)	9.87	(7.71 - 12.56)
Circuit 20	18.62	(15.08 - 22.77)	10.25	(7.84 - 13.30)
Northeast	20.61	(18.35 - 23.08)	11.65	(10.05 - 13.46)
Circuit 4	19.33	(15.84 - 23.39)	11.48	(8.94 - 14.63)
Circuit 5	17.24	(13.66 - 21.53)	9.39	(7.09 - 12.33)
Circuit 7	24.48	(20.20 - 29.34)	12.18	(9.32 - 15.77)
Circuit 8 plus Columbia, Dixie, Hamilton, Lafayette, and Suwannee	22.75	(18.00 - 28.32)	14.62	(11.17 - 18.90)
Northwest	24.80	(21.75 - 28.13)	16.13	(13.61 - 19.02)
Circuit 1	22.92	(18.63 - 27.85)	14.80	(11.45 - 18.93)
Circuit 2 plus Madison and Taylor	30.48	(25.15 - 36.40)	21.45	(16.88 - 26.86)
Circuit 14	19.49	(15.30 - 24.49)	10.03	(7.44 - 13.39)
South (Circuits 11 and 16)	18.40	(15.64 - 21.54)	10.73	(8.67 - 13.20)
Southeast	18.42	(15.55 - 21.68)	9.67	(7.80 - 11.94)
Circuit 15 (Palm Beach)	18.41	(15.00 - 22.40)	10.02	(7.69 - 12.96)
Circuit 19	18.42	(14.24 - 23.49)	8.91	(6.43 - 12.22)

NOTE: For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹ Binge Alcohol Use is defined as drinking five or more drinks (for males) or four or more drinks (for females) on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.

Table 10.7 Tobacco Product Use in the Past Month, Cigarette Use in the Past Month, and Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day in Florida among Individuals Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs

State/Substate Region	Tobacco Product Use in the Past Month ¹		Cigarette Use in the Past Month		Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day	
	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Florida	20.09	(19.04 - 21.18)	16.39	(15.45 - 17.37)	76.42	(75.22 - 77.58)
Broward (Circuit 17)	15.03	(12.90 - 17.43)	12.47	(10.46 - 14.79)	79.64	(77.25 - 81.85)
Central I	18.76	(16.99 - 20.67)	15.52	(13.84 - 17.37)	76.60	(74.76 - 78.35)
Circuit 9	16.72	(14.62 - 19.06)	14.08	(12.12 - 16.29)	77.74	(75.54 - 79.80)
Circuit 18	21.82	(18.98 - 24.94)	17.69	(15.04 - 20.70)	74.89	(72.08 - 77.51)
Central II	21.86	(20.27 - 23.53)	17.54	(16.11 - 19.06)	75.73	(74.12 - 77.26)
Circuit 6	23.59	(20.67 - 26.79)	18.37	(15.70 - 21.37)	74.58	(71.68 - 77.27)
Circuit 10	24.39	(21.19 - 27.89)	20.52	(17.32 - 24.14)	74.38	(71.47 - 77.09)
Circuit 12	21.68	(18.54 - 25.19)	17.59	(14.67 - 20.96)	76.14	(73.08 - 78.95)
Circuit 13 (Hillsborough)	19.64	(17.15 - 22.39)	15.64	(13.37 - 18.20)	76.50	(74.06 - 78.77)
Circuit 20	20.81	(17.87 - 24.10)	16.78	(14.06 - 19.89)	76.77	(73.82 - 79.49)
Northeast	23.88	(22.02 - 25.85)	19.65	(18.01 - 21.41)	73.66	(71.81 - 75.42)
Circuit 4	23.35	(20.30 - 26.69)	18.67	(15.99 - 21.67)	72.99	(69.97 - 75.80)
Circuit 5	24.91	(21.73 - 28.39)	20.59	(17.56 - 23.99)	74.54	(71.46 - 77.39)
Circuit 7	22.51	(19.36 - 26.01)	18.76	(15.88 - 22.02)	75.09	(72.21 - 77.76)
Circuit 8 plus Columbia, Dixie, Hamilton, Lafayette, and Suwannee	25.20	(21.75 - 28.98)	21.30	(17.91 - 25.14)	70.83	(67.60 - 73.87)
Northwest	25.55	(23.27 - 27.97)	20.49	(18.33 - 22.84)	72.34	(70.20 - 74.39)
Circuit 1	24.81	(21.53 - 28.40)	19.67	(16.59 - 23.16)	71.76	(68.65 - 74.68)
Circuit 2 plus Madison and Taylor	24.95	(21.56 - 28.67)	20.26	(17.01 - 23.96)	74.31	(71.40 - 77.02)
Circuit 14	28.23	(24.17 - 32.68)	22.80	(18.99 - 27.11)	70.83	(67.34 - 74.09)
South (Circuits 11 and 16)	13.87	(12.12 - 15.84)	11.57	(9.96 - 13.40)	80.37	(78.44 - 82.17)
Southeast	19.09	(17.02 - 21.36)	15.56	(13.64 - 17.70)	77.77	(75.60 - 79.79)
Circuit 15 (Palm Beach)	17.00	(14.63 - 19.65)	13.85	(11.69 - 16.35)	78.56	(75.88 - 81.01)
Circuit 19	23.71	(20.03 - 27.84)	19.33	(15.83 - 23.40)	76.02	(72.97 - 78.83)

NOTE: For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., snuff, dip, chewing tobacco, or snus), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.

Table 10.8 Illicit Drug Use Disorder in the Past Year, Pain Reliever Use Disorder in the Past Year, Alcohol Use Disorder in the Past Year, and Substance Use Disorder in the Past Year in Florida among Individuals Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs

State/Substate Region	Illicit Drug Use Disorder in the Past Year ^{1,2}		Pain Reliever Use Disorder in the Past Year ²		Alcohol Use Disorder in the Past Year		Substance Use Disorder in the Past Year ^{1,2}	
	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Florida	2.70	(2.38-3.07)	0.71	(0.56-0.91)	4.85	(4.34-5.41)	6.60	(6.04-7.20)
Broward (Circuit 17)	2.67	(2.04-3.49)	0.66	(0.39-1.14)	4.30	(3.26-5.65)	6.10	(4.86-7.62)
Central I	2.66	(2.17-3.27)	0.64	(0.43-0.94)	4.47	(3.68-5.43)	6.24	(5.32-7.30)
Circuit 9	2.73	(2.13-3.50)	0.61	(0.37-1.01)	4.22	(3.31-5.37)	6.19	(5.09-7.50)
Circuit 18	2.55	(1.94-3.36)	0.68	(0.40-1.14)	4.85	(3.66-6.40)	6.32	(4.99-7.97)
Central II	2.55	(2.16-3.00)	0.74	(0.54-1.00)	5.01	(4.27-5.88)	6.60	(5.84-7.46)
Circuit 6	2.54	(1.93-3.32)	0.79	(0.47-1.33)	4.98	(3.73-6.62)	6.56	(5.30-8.09)
Circuit 10	2.52	(1.87-3.39)	0.73	(0.41-1.29)	4.53	(3.34-6.12)	6.19	(4.84-7.90)
Circuit 12	2.34	(1.74-3.16)	0.75	(0.42-1.33)	4.97	(3.58-6.85)	6.24	(4.85-7.99)
Circuit 13 (Hillsborough)	2.96	(2.30-3.80)	0.62	(0.38-1.02)	5.29	(4.12-6.78)	7.21	(5.92-8.75)
Circuit 20	2.26	(1.69-3.03)	0.78	(0.45-1.37)	5.05	(3.73-6.82)	6.48	(5.07-8.26)
Northeast	2.90	(2.42-3.46)	0.80	(0.57-1.11)	5.19	(4.36-6.17)	7.13	(6.22-8.16)
Circuit 4	2.88	(2.17-3.82)	0.76	(0.46-1.26)	5.61	(4.25-7.36)	7.42	(5.93-9.24)
Circuit 5	2.67	(2.01-3.55)	0.80	(0.47-1.37)	4.72	(3.49-6.36)	6.60	(5.19-8.37)
Circuit 7	2.96	(2.22-3.93)	0.83	(0.48-1.43)	4.99	(3.68-6.74)	6.95	(5.48-8.77)
Circuit 8 plus Columbia, Dixie, Hamilton, Lafayette, and Suwannee	3.28	(2.43-4.43)	0.80	(0.47-1.36)	5.61	(4.15-7.54)	7.90	(6.18-10.06)
Northwest	3.44	(2.81-4.21)	0.84	(0.58-1.21)	5.86	(4.79-7.16)	8.24	(7.02-9.64)
Circuit 1	3.22	(2.39-4.32)	0.90	(0.53-1.53)	5.57	(4.12-7.49)	7.66	(6.04-9.67)
Circuit 2 plus Madison and Taylor	4.00	(2.98-5.34)	0.73	(0.43-1.24)	6.99	(5.26-9.22)	10.04	(7.95-12.61)
Circuit 14	3.13	(2.28-4.29)	0.83	(0.48-1.44)	4.90	(3.51-6.79)	6.95	(5.32-9.03)
South (Circuits 11 and 16)	2.56	(2.00-3.27)	0.63	(0.38-1.03)	4.22	(3.35-5.31)	5.87	(4.89-7.05)
Southeast	2.53	(2.00-3.19)	0.67	(0.44-1.04)	4.84	(3.85-6.06)	6.34	(5.23-7.67)
Circuit 15 (Palm Beach)	2.66	(2.01-3.51)	0.68	(0.39-1.18)	4.79	(3.59-6.36)	6.37	(5.05-8.02)
Circuit 19	2.24	(1.62-3.11)	0.65	(0.37-1.15)	4.95	(3.59-6.80)	6.28	(4.77-8.22)

NOTE: For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

NOTE: Substance Use Disorder is defined as meeting criteria for illicit drug or alcohol dependence or abuse. Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

¹ Illicit Drug Use includes the misuse of prescription psychotherapeutics or the use of marijuana, cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine.

² Misuse of prescription psychotherapeutics is defined as use in any way not directed by a doctor, including use without a prescription of one's own; use in greater amounts, more often, or longer than told; or use in any other way not directed by a doctor. Prescription psychotherapeutics do not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.

Table 10.9 *Needing But Not Receiving Treatment at a Specialty Facility for Illicit Drug Use in the Past Year, Needing But Not Receiving Treatment at a Specialty Facility for Alcohol Use in the Past Year, and Needing But Not Receiving Treatment at a Specialty Facility for Substance Use in the Past Year in Florida among Individuals Aged 12 or Older, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs*

State/Substate Region	Needing But Not Receiving Treatment at a Specialty Facility for Illicit Drug Use in the Past Year ¹		Needing But Not Receiving Treatment at a Specialty Facility for Alcohol Use in the Past Year		Needing But Not Receiving Treatment at a Specialty Facility for Substance Use in the Past Year ¹	
	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Florida	2.42	(2.13 - 2.74)	4.65	(4.17 - 5.18)	6.27	(5.74 - 6.84)
Broward (Circuit 17)	2.32	(1.75 - 3.07)	4.06	(3.11 - 5.29)	5.66	(4.47 - 7.14)
Central I	2.43	(1.98 - 2.98)	4.39	(3.64 - 5.28)	5.99	(5.06 - 7.07)
Circuit 9	2.55	(1.98 - 3.28)	4.11	(3.24 - 5.21)	5.97	(4.86 - 7.32)
Circuit 18	2.24	(1.68 - 2.98)	4.80	(3.65 - 6.29)	6.01	(4.64 - 7.75)
Central II	2.28	(1.94 - 2.68)	4.72	(4.05 - 5.49)	6.20	(5.47 - 7.03)
Circuit 6	2.14	(1.60 - 2.86)	4.67	(3.57 - 6.09)	5.96	(4.73 - 7.50)
Circuit 10	2.37	(1.74 - 3.22)	4.23	(3.16 - 5.63)	5.79	(4.49 - 7.43)
Circuit 12	2.23	(1.64 - 3.03)	4.68	(3.46 - 6.31)	6.01	(4.62 - 7.78)
Circuit 13 (Hillsborough)	2.77	(2.15 - 3.57)	4.76	(3.70 - 6.10)	6.86	(5.56 - 8.45)
Circuit 20	1.90	(1.42 - 2.53)	5.04	(3.77 - 6.72)	6.14	(4.73 - 7.94)
Northeast	2.50	(2.08 - 3.00)	4.97	(4.21 - 5.86)	6.72	(5.84 - 7.71)
Circuit 4	2.54	(1.89 - 3.40)	5.10	(3.86 - 6.72)	6.88	(5.41 - 8.71)
Circuit 5	2.31	(1.70 - 3.14)	4.59	(3.43 - 6.13)	6.41	(4.94 - 8.27)
Circuit 7	2.48	(1.85 - 3.33)	4.91	(3.68 - 6.51)	6.42	(4.99 - 8.21)
Circuit 8 plus Columbia, Dixie, Hamilton, Lafayette, and Suwannee	2.85	(2.10 - 3.86)	5.57	(4.23 - 7.30)	7.51	(5.80 - 9.67)
Northwest	3.08	(2.49 - 3.79)	5.77	(4.78 - 6.96)	8.04	(6.80 - 9.47)
Circuit 1	2.90	(2.13 - 3.93)	5.68	(4.27 - 7.53)	7.41	(5.77 - 9.45)
Circuit 2 plus Madison and Taylor	3.67	(2.69 - 4.98)	6.64	(5.08 - 8.64)	9.90	(7.73 - 12.59)
Circuit 14	2.63	(1.91 - 3.62)	4.70	(3.48 - 6.33)	6.79	(5.08 - 9.03)
South (Circuits 11 and 16)	2.37	(1.85 - 3.02)	4.09	(3.25 - 5.15)	5.66	(4.65 - 6.86)
Southeast	2.33	(1.82 - 2.97)	4.67	(3.73 - 5.83)	6.07	(5.02 - 7.33)
Circuit 15 (Palm Beach)	2.45	(1.83 - 3.28)	4.63	(3.49 - 6.12)	6.05	(4.78 - 7.63)
Circuit 19	2.06	(1.47 - 2.87)	4.77	(3.53 - 6.42)	6.12	(4.61 - 8.08)

NOTE: For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

NOTE: Respondents were classified as needing substance use treatment if they met the criteria for an illicit drug or alcohol use disorder as defined in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* or received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Needing But Not Receiving Substance Use Treatment refers to respondents who are classified as needing illicit drug or alcohol treatment, but who did not receive illicit drug or alcohol treatment at a specialty facility.

¹ Illicit Drug Use includes the misuse of prescription psychotherapeutics or the use of marijuana, cocaine (including crack), heroin, hallucinogens, inhalants, or methamphetamine. Misuse of prescription psychotherapeutics is defined as use in any way not directed by a doctor, including use without a prescription of one's own; use in greater amounts, more often, or longer than told; or use in any other way not directed by a doctor. Prescription psychotherapeutics do not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.

Table 10.10 Serious Mental Illness in the Past Year, Any Mental Illness in the Past Year, and Received Mental Health Services in the Past Year in Florida among Adults Aged 18 or Older, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs

State/Substate Region	Serious Mental Illness in the Past Year ¹		Any Mental Illness in the Past Year ¹		Received Mental Health Services in the Past Year ²	
	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Florida	3.82	(3.37 - 4.32)	17.28	(16.22 - 18.40)	12.53	(11.60 - 13.52)
Broward (Circuit 17)	3.15	(2.45 - 4.04)	15.44	(13.20 - 17.98)	10.85	(9.14 - 12.83)
Central I	3.85	(3.22 - 4.59)	17.67	(15.94 - 19.54)	12.34	(10.90 - 13.94)
Circuit 9	3.77	(3.01 - 4.70)	17.67	(15.54 - 20.02)	11.34	(9.69 - 13.23)
Circuit 18	3.96	(3.13 - 5.01)	17.66	(15.20 - 20.43)	13.83	(11.68 - 16.31)
Central II	4.00	(3.43 - 4.65)	18.00	(16.54 - 19.55)	13.74	(12.42 - 15.19)
Circuit 6	4.16	(3.26 - 5.30)	18.52	(15.92 - 21.44)	14.96	(12.71 - 17.52)
Circuit 10	4.07	(3.15 - 5.25)	16.77	(14.17 - 19.73)	12.33	(10.24 - 14.77)
Circuit 12	3.73	(2.88 - 4.81)	16.99	(14.33 - 20.04)	14.54	(12.05 - 17.45)
Circuit 13 (Hillsborough)	4.31	(3.43 - 5.41)	18.98	(16.51 - 21.71)	13.66	(11.59 - 16.03)
Circuit 20	3.60	(2.82 - 4.60)	17.75	(15.16 - 20.66)	12.75	(10.57 - 15.31)
Northeast	4.22	(3.59 - 4.94)	18.06	(16.47 - 19.77)	13.97	(12.61 - 15.45)
Circuit 4	4.41	(3.48 - 5.57)	17.57	(15.07 - 20.38)	14.09	(11.88 - 16.63)
Circuit 5	3.89	(3.03 - 4.98)	18.75	(16.04 - 21.80)	13.67	(11.51 - 16.16)
Circuit 7	3.96	(3.06 - 5.10)	16.92	(14.31 - 19.90)	14.10	(11.84 - 16.71)
Circuit 8 plus Columbia, Dixie, Hamilton, Lafayette, and Suwannee	4.92	(3.89 - 6.21)	19.58	(16.71 - 22.81)	14.11	(11.82 - 16.77)
Northwest	4.76	(3.99 - 5.67)	20.17	(18.12 - 22.40)	14.67	(12.95 - 16.58)
Circuit 1	4.82	(3.79 - 6.12)	19.89	(16.99 - 23.15)	15.26	(12.77 - 18.14)
Circuit 2 plus Madison and Taylor	4.71	(3.73 - 5.95)	20.93	(17.90 - 24.31)	13.56	(11.37 - 16.11)
Circuit 14	4.68	(3.60 - 6.07)	19.72	(16.61 - 23.24)	14.91	(12.42 - 17.79)
South (Circuits 11 and 16)	3.13	(2.48 - 3.94)	14.56	(12.61 - 16.75)	8.57	(7.21 - 10.15)
Southeast	3.41	(2.78 - 4.18)	16.72	(14.67 - 19.00)	12.17	(10.54 - 14.00)
Circuit 15 (Palm Beach)	3.23	(2.50 - 4.16)	16.58	(14.08 - 19.43)	11.85	(9.91 - 14.11)
Circuit 19	3.80	(2.94 - 4.91)	17.04	(14.21 - 20.29)	12.88	(10.63 - 15.51)

NOTE: For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹ Mental illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, assessed by the Mental Health Surveillance Study (MHSS) *Structured Clinical Interview for the Diagnostic and Statistical Manual of Mental Disorders—Fourth Edition—Research Version—Axis I Disorders* (MHSS-SCID), which is based on the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Any mental illness (AMI) includes adults with any mental disorder regardless of whether their disorder resulted in functional impairment. Serious mental illness (SMI) includes adults with any mental disorder that resulted in serious functional impairment. As such, these estimates are based on indicators of AMI and SMI rather than direct measures of diagnostic status. For details, see Section B of the "2017-2018 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at <https://www.samhsa.gov/data/>.

² Mental health services are defined as having received inpatient treatment/counseling or outpatient treatment/counseling or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.

Table 10.11 Had Serious Thoughts of Suicide in the Past Year and Major Depressive Episode in the Past Year in Florida among Adults Aged 18 or Older, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs

State/Substate Region	Had Serious Thoughts of Suicide in the Past Year ¹		Major Depressive Episode in the Past Year ²	
	Estimate	95% Confidence Interval	Estimate	95% Confidence Interval
Florida	3.47	(3.05 - 3.95)	6.13	(5.53 - 6.79)
Broward (Circuit 17)	3.17	(2.54 - 3.96)	5.33	(4.32 - 6.55)
Central I	3.49	(2.92 - 4.15)	6.57	(5.65 - 7.63)
Circuit 9	3.44	(2.79 - 4.24)	6.51	(5.43 - 7.80)
Circuit 18	3.55	(2.80 - 4.50)	6.66	(5.37 - 8.23)
Central II	3.61	(3.08 - 4.21)	6.27	(5.52 - 7.13)
Circuit 6	3.66	(2.88 - 4.65)	6.49	(5.24 - 8.03)
Circuit 10	3.58	(2.79 - 4.58)	5.86	(4.66 - 7.36)
Circuit 12	3.54	(2.73 - 4.58)	6.08	(4.80 - 7.66)
Circuit 13 (Hillsborough)	3.73	(3.04 - 4.57)	6.80	(5.55 - 8.30)
Circuit 20	3.46	(2.72 - 4.41)	5.84	(4.68 - 7.26)
Northeast	3.59	(3.05 - 4.22)	6.55	(5.68 - 7.55)
Circuit 4	3.56	(2.82 - 4.48)	6.61	(5.28 - 8.25)
Circuit 5	3.36	(2.61 - 4.30)	6.16	(4.88 - 7.74)
Circuit 7	3.49	(2.76 - 4.41)	6.32	(4.98 - 7.99)
Circuit 8 plus Columbia, Dixie, Hamilton, Lafayette, and Suwannee	4.31	(3.43 - 5.40)	7.62	(6.16 - 9.40)
Northwest	4.28	(3.59 - 5.09)	7.38	(6.30 - 8.63)
Circuit 1	4.25	(3.36 - 5.36)	7.63	(6.11 - 9.49)
Circuit 2 plus Madison and Taylor	4.53	(3.61 - 5.67)	7.41	(5.97 - 9.17)
Circuit 14	3.97	(3.11 - 5.06)	6.75	(5.30 - 8.55)
South (Circuits 11 and 16)	3.05	(2.45 - 3.79)	5.08	(4.19 - 6.14)
Southeast	3.16	(2.60 - 3.84)	5.69	(4.74 - 6.82)
Circuit 15 (Palm Beach)	3.06	(2.42 - 3.86)	5.51	(4.39 - 6.90)
Circuit 19	3.39	(2.63 - 4.36)	6.09	(4.78 - 7.72)

NOTE: For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.


NOTE: Estimates along with the 95 percent Bayesian confidence (credible) intervals are based on a survey-weighted hierarchical Bayes estimation approach and generated by Markov Chain Monte Carlo techniques.

¹ Respondents were asked, "At any time in the past 12 months, did you seriously think about trying to kill yourself?" If they answered "Yes," they were categorized as having serious thoughts of suicide in the past year.

² Major depressive episode (MDE) is defined as in the 5th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-5), which specifies a period of at least 2 weeks when an individual experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. For details, see Section B of the "2017-2018 NSDUH: Guide to State Tables and Summary of Small Area Estimation Methodology" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.1 Illicit Drug Use in the Past Month among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.2 Marijuana Use in the Past Year among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.3 Marijuana Use in the Past Month among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.4 Perceptions of Great Risk from Smoking Marijuana Once a Month among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.5 First Use of Marijuana among Individuals Aged 12 or Older in Florida, by Substate Region: Average Annual Initiation Estimates (Expressed as Percentages of the At-Risk Population) Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.6 Illicit Drug Use Other Than Marijuana in the Past Month among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.7 Cocaine Use in the Past Year among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.8 Perceptions of Great Risk from Using Cocaine Once a Month among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.9 Heroin Use in the Past Year among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.10 *Perceptions of Great Risk from Trying Heroin Once or Twice among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs*



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.11 Methamphetamine Use in the Past Year among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.12 Pain Reliever Misuse in the Past Year among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.13 Alcohol Use in the Past Month among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.14 Binge Alcohol Use in the Past Month among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.15 *Perceptions of Great Risk from Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs*



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.16 Alcohol Use in the Past Month among Individuals Aged 12 to 20 in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.17 Binge Alcohol Use in the Past Month among Individuals Aged 12 to 20 in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.18 Tobacco Product Use in the Past Month among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.19 Cigarette Use in the Past Month among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.20 *Perceptions of Great Risk from Smoking One or More Packs of Cigarettes per Day among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs*



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.21 Illicit Drug Use Disorder in the Past Year among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.22 Pain Reliever Use Disorder in the Past Year among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.23 Alcohol Use Disorder in the Past Year among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.24 Substance Use Disorder in the Past Year among Individuals Aged 12 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.25 Serious Mental Illness in the Past Year among Adults Aged 18 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.26 Any Mental Illness in the Past Year among Adults Aged 18 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.27 Received Mental Health Services in the Past Year among Adults Aged 18 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.


Figure 10.28 Had Serious Thoughts of Suicide in the Past Year among Adults Aged 18 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.

Figure 10.29 Major Depressive Episode in the Past Year among Adults Aged 18 or Older in Florida, by Substate Region: Percentages, Annual Averages Based on 2016, 2017, and 2018 NSDUHs



NOTE: The legend's ranges were created by dividing 395 substate regions, nationally, into 7 groups based on the magnitude of their percentages. For substate region definitions, see the "2016-2018 National Survey on Drug Use and Health Substate Region Definitions" at <https://www.samhsa.gov/data/>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2016, 2017, and 2018.