

Table 1.103A Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 or Older: Numbers in Thousands, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	18,008	18,290	5,109	4,929	5,109	4,929
Amphetamine or Methylphenidate Products	13,749	14,294	4,835	4,711	4,835	4,711
Amphetamine Products	12,215	12,647	4,615	4,486	4,615	4,486
Methylphenidate Products	3,143	3,383	747	704	747	704
Anorectic (Weight-Loss) Stimulants	2,506	2,220	147	207	147	207
Provigil®	415	536	125	87	*	*
Any Other Prescription Stimulant ¹	2,626	2,594	84	31	84	31

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.103B Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 or Older: Percentages, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	6.6	6.6	1.9	1.8	28.4	26.9
Amphetamine or Methylphenidate Products	5.0	5.2	1.8	1.7	35.2	33.0
Amphetamine Products	4.5	4.6	1.7	1.6	37.8	35.5
Methylphenidate Products	1.1	1.2	0.3	0.3	23.8	20.8
Anorectic (Weight-Loss) Stimulants	0.9	0.8	0.1	0.1	5.9	9.3
Provigil®	0.2	0.2	0.0	0.0	*	*
Any Other Prescription Stimulant ¹	1.0	0.9	0.0	0.0	3.2	1.2

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.104A Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 to 17 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 to 17: Numbers in Thousands, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	1,752	1,837	369	430	369	430
Amphetamine or Methylphenidate Products	1,572	1,643	350	407	350	407
Amphetamine Products	1,223	1,301	320	373	320	373
Methylphenidate Products	561	590	74	80	74	80
Anorectic (Weight-Loss) Stimulants	51	72	7	15	*	*
Provigil®	19	9	3	2	*	*
Any Other Prescription Stimulant ¹	286	360	12	17	12	17

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.104B Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 12 to 17 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 12 to 17: Percentages, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	7.0	7.4	1.5	1.7	21.1	23.4
Amphetamine or Methylphenidate Products	6.3	6.6	1.4	1.6	22.3	24.8
Amphetamine Products	4.9	5.2	1.3	1.5	26.1	28.6
Methylphenidate Products	2.3	2.4	0.3	0.3	13.2	13.6
Anorectic (Weight-Loss) Stimulants	0.2	0.3	0.0	0.1	*	*
Provigil®	0.1	0.0	0.0	0.0	*	*
Any Other Prescription Stimulant ¹	1.2	1.4	0.0	0.1	4.1	4.7

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.105A Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 or Older: Numbers in Thousands, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	16,256	16,453	4,739	4,499	4,739	4,499
Amphetamine or Methylphenidate Products	12,177	12,650	4,486	4,304	4,486	4,304
Amphetamine Products	10,992	11,346	4,295	4,114	4,295	4,114
Methylphenidate Products	2,583	2,792	673	623	673	623
Anorectic (Weight-Loss) Stimulants	2,455	2,148	140	193	140	193
Provigil®	396	527	121	85	*	*
Any Other Prescription Stimulant ¹	2,340	2,235	73 ^a	14	73 ^a	14

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.105B Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 or Older: Percentages, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	6.5	6.6	1.9	1.8	29.2	27.3
Amphetamine or Methylphenidate Products	4.9	5.1	1.8	1.7	36.8	34.0
Amphetamine Products	4.4	4.5	1.7	1.6	39.1	36.3
Methylphenidate Products	1.0	1.1	0.3	0.2	26.1	22.3
Anorectic (Weight-Loss) Stimulants	1.0	0.9	0.1	0.1	5.7	9.0
Provigil®	0.2	0.2	0.0	0.0	*	*
Any Other Prescription Stimulant ¹	0.9	0.9	0.0 ^a	0.0	3.1 ^a	0.6

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.106A Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 to 25 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 to 25: Numbers in Thousands, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	4,473	4,320	2,216 ^a	1,951	2,216 ^a	1,951
Amphetamine or Methylphenidate Products	4,120	3,998	2,175 ^a	1,926	2,175	1,926
Amphetamine Products	3,740	3,615	2,092 ^a	1,845	2,092	1,845
Methylphenidate Products	1,058	1,011	397	300	397	300
Anorectic (Weight-Loss) Stimulants	195	189	16	27	*	*
Provigil [®]	48	47	26	10	*	*
Any Other Prescription Stimulant ¹	435	361	13	14	13	14

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.106B Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 18 to 25 and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 18 to 25: Percentages, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	13.1	12.8	6.5	5.8	49.5 ^a	45.2
Amphetamine or Methylphenidate Products	12.1	11.9	6.4	5.7	52.8 ^a	48.2
Amphetamine Products	11.0	10.7	6.1	5.5	55.9 ^a	51.0
Methylphenidate Products	3.1	3.0	1.2	0.9	37.5	29.7
Anorectic (Weight-Loss) Stimulants	0.6	0.6	0.0	0.1	*	*
Provigil [®]	0.1	0.1	0.1	0.0	*	*
Any Other Prescription Stimulant ¹	1.3	1.1	0.0	0.0	3.0	4.0

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.107A Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 26 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 26 or Older: Numbers in Thousands, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse in Past Year among Total Population (2018)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
ANY PRESCRIPTION STIMULANT	11,783	12,133	2,524	2,548	2,524	2,548
Amphetamine or Methylphenidate Products	8,057	8,652	2,310	2,378	2,310	2,378
Amphetamine Products	7,252	7,732	2,204	2,269	2,204	2,269
Methylphenidate Products	1,524	1,781	276	323	276	323
Anorectic (Weight-Loss) Stimulants	2,260	1,959	124	165	124	165
Provigil®	348	480	95	76	*	*
Any Other Prescription Stimulant ¹	1,905	1,874	59	*	59	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.

Table 1.107B Any Use and Misuse of Stimulant Subtypes in Past Year among Persons Aged 26 or Older and Misuse of Stimulant Subtypes in Past Year among Past Year Any Users of Stimulant Subtypes Aged 26 or Older: Percentages, 2018 and 2019

Stimulant Subtype	Any Use in Past Year (2018)	Any Use in Past Year (2019)	Misuse		Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2018)	Misuse in Past Year among Past Year Any Users of Stimulant Subtypes (2019)
			in Past Year among Total Population (2018)	in Past Year among Total Population (2019)		
ANY PRESCRIPTION STIMULANT	5.5	5.6	1.2	1.2	21.4	21.0
Amphetamine or Methylphenidate Products	3.8	4.0	1.1	1.1	28.7	27.5
Amphetamine Products	3.4	3.6	1.0	1.0	30.4	29.3
Methylphenidate Products	0.7	0.8	0.1	0.1	18.1	18.1
Anorectic (Weight-Loss) Stimulants	1.1	0.9	0.1	0.1	5.5	8.4
Provigil®	0.2	0.2	0.0	0.0	*	*
Any Other Prescription Stimulant ¹	0.9	0.9	0.0	*	3.1	*

* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

^a The difference between this estimate and the 2019 estimate is statistically significant at the .05 level. Rounding may make the estimates appear identical.

¹ Includes use or misuse of stimulants containing other active ingredients. Reports of misuse of "any other prescription stimulant" that correspond only to the specific stimulant categories shown in the table are excluded from estimates for Any Other Prescription Stimulant and are included instead in the relevant stimulant category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2018 and 2019.